

# Regular sequences of pulsation overtones in young intermediate-mass stars

Timothy R. Bedding<sup>1,2</sup>, Simon J. Murphy<sup>1,2</sup>, Daniel R. Hey<sup>1,2</sup>, Daniel Huber<sup>3,2</sup>,  
Tanda Li<sup>1,2</sup>, Gang Li<sup>1,2</sup>, Yaguang Li<sup>4</sup>, Barry Smalley<sup>5</sup>, Dennis Stello<sup>6,2</sup>, William J. Chaplin<sup>7,2</sup>,  
Isabel L. Colman<sup>1,2</sup>, Jim Fuller<sup>8</sup>, Eric Gaidos<sup>9</sup>, Daniel R. Harbeck<sup>10</sup>, J. J. Hermes<sup>11</sup>,  
Andrew W. Mann<sup>12</sup>, Daniel R. Reese<sup>13</sup>, Sanjay Sekaran<sup>14</sup>, Timothy R. White<sup>1,2,15</sup>, Jie Yu<sup>16</sup>,  
Victoria Antoci<sup>2</sup>, Timothy M. Brown<sup>10</sup>, Andrew W. Howard<sup>8</sup>, Howard Isaacson<sup>17</sup>,  
Curtis McCully<sup>10</sup> & Markus Rabus<sup>10,18</sup>

<sup>1</sup>*Sydney Institute for Astronomy (SIfA), School of Physics, University of Sydney, 2006, Australia*

<sup>2</sup>*Stellar Astrophysics Centre, Department of Physics and Astronomy, Aarhus University, 8000 Aarhus C, Denmark*

<sup>3</sup>*Institute for Astronomy, University of Hawai‘i, 2680 Woodlawn Drive, Honolulu, HI 96822, USA*

<sup>4</sup>*Department of Astronomy, Beijing Normal University, Beijing 100875, China*

<sup>5</sup>*Astrophysics Group, Lennard-Jones Laboratories, Keele University, Staffordshire ST5 5BG, United Kingdom*

<sup>6</sup>*School of Physics, University of New South Wales, 2052, Australia*

<sup>7</sup>*School of Physics and Astronomy, University of Birmingham, Birmingham B15 2TT, UK*

<sup>8</sup>*TAPIR, Mailcode 350-17, California Institute of Technology, Pasadena, CA 91125, USA*

<sup>9</sup>*Department of Earth Sciences, University of Hawai‘i, Honolulu HI 96822, USA*

<sup>10</sup>*Las Cumbres Observatory Global Telescope, 6740 Cortona Dr., Suite 102, Goleta, CA 93111, USA*

<sup>11</sup>*Department of Astronomy, Boston University, 725 Commonwealth Ave., Boston, MA 02215, USA*

<sup>12</sup>*Department of Physics and Astronomy, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599-3255, USA*

<sup>13</sup>*LESIA, Observatoire de Paris, PSL Research University, CNRS, Sorbonne Universités, UPMC Univ. Paris 06, Univ. Paris Diderot, Sorbonne Paris Cité, 5 place Jules Janssen, 92195 Meudon, France*

<sup>14</sup>*Instituut voor Sterrenkunde (IvS), KU Leuven, Celestijnenlaan 200D, 3001 Leuven, Belgium*

<sup>15</sup>*Research School of Astronomy and Astrophysics, Mount Stromlo Observatory, The Australian National University, Canberra, ACT 2611, Australia*

<sup>16</sup>*Max-Planck-Institut für Sonnensystemforschung, Justus-von-Liebig-Weg 3, 37077 Göttingen, Germany*

<sup>17</sup>*Department of Astronomy, University of California at Berkeley, Berkeley, CA 94720-3411, USA*

<sup>18</sup>*Department of Physics, University of California, Santa Barbara, CA 93106-9530, USA*

**Asteroseismology is a powerful tool for probing the internal structures of stars by using their natural pulsation frequencies<sup>1</sup>. It relies on identifying sequences of pulsation modes that can be compared with theoretical models, which has been done with great success for many classes of pulsators including low-mass solar-type stars<sup>2</sup>, red giants<sup>3</sup>, high-mass stars<sup>4</sup> and white dwarfs<sup>5</sup>. However, a large group of pulsating stars of intermediate mass—the so-called  $\delta$  Scuti stars—have rich pulsation spectra for which systematic mode identification has not been possible<sup>1,6–8</sup>. This arises because only a seemingly random subset of possible modes are excited, and because rapid rotation tends to spoil the regular patterns<sup>9,10</sup>. Here we report the detection of remarkably regular sequences of pulsation modes in 60 intermediate-mass main-sequence stars, opening up a new regime in which the tools of asteroseismology can be applied. As an example, some of these stars have space motions that indicate they are members of young moving groups, and their pulsation spectra confirm that these stars are indeed young.**

The  $\delta$  Scuti variables are main-sequence stars of intermediate mass ( $1.5\text{--}2.5 M_{\odot}$ ) that pulsate in low-order pressure modes<sup>1,6–8</sup>. Observations have suggested that some  $\delta$  Scuti stars have regular frequency spacings in their Fourier amplitude spectra<sup>11–24</sup>, but the lack of a large sample has prevented a systematic and unambiguous mode identification. Each pulsation mode in a non-rotating star is identified by two integers: the radial order ( $n$ ) and the degree ( $l$ )<sup>1</sup>. We expect the strongest observable modes to be of low degree ( $l = 0, 1$  and  $2$ ), since higher degrees have greatly reduced amplitudes due to cancellation in disk-integrated light. In the so-called asymptotic regime ( $n \gg l$ ), modes with a given degree  $l$  are approximately equally spaced, with a frequency separation ( $\Delta\nu$ ) that depends on the time taken for sound waves to travel through the star, and is approximately proportional to the square root of the mean stellar density.

The patterns are more complex in a rotating star, with the mode frequencies also depending on the azimuthal order ( $m$ ). Each nonradial ( $l \geq 1$ ) mode in the Fourier spectrum is split into  $2l + 1$  components, where  $m$  ranges from  $-l$  to  $l$ . The relative amplitudes of these components depend on the inclination of the rotation axis to the line of sight. For example, if a star is seen at low inclination (close to pole-on) then the zonal ( $m = 0$ ) mode in each multiplet will dominate, leading to a simpler Fourier spectrum. In very rapidly rotating stars, the oblateness alters the pulsation cavity and further complicates the pattern. However, for rotation rates less than  $\sim 50\%$  of Keplerian break-up, the radial modes ( $l = 0$ ) and the zonal dipolar modes ( $l = 1, m = 0$ ) are still expected to follow a regular spacing<sup>25</sup>.

To search for regular patterns we have used observations from the *Transiting Exoplanet Survey Satellite* (*TESS*)<sup>26</sup>, which for the first time provides light curves of many thousands of  $\delta$  Scuti stars at rapid cadence (120 s sampling). We used the first nine 27-day sectors *TESS* data and focussed on identifying  $\delta$  Scuti stars that pulsate at high frequencies (above about  $30 \text{ d}^{-1}$ )<sup>27–29</sup>. We also examined stars not previously known to pulsate by calculating the amplitude spectra of *TESS* short-cadence light curves, measuring the skewness of the distribution of peak heights<sup>30</sup> above  $30 \text{ d}^{-1}$  as a way to flag likely detections, and inspecting the most prominent peaks for regularity

using échelle diagrams (described below). In addition, we used data from the *Kepler* spacecraft, which observed about 300  $\delta$  Scuti stars at short cadence (60 s sampling) during its four-year nominal mission<sup>24,30,31</sup>. Stars observed in *Kepler*’s long-cadence mode (29.4-min sampling) were not considered because the Nyquist frequency of  $24 \text{ d}^{-1}$  makes it very difficult to identify patterns in high-frequency pulsators.

We discovered 60 stars with regular frequency spacings (Table S1 in the Supplementary Information), which define a new group of  $\delta$  Scuti stars for which mode identification is possible. Figure 1 shows some of the amplitude spectra, which have remarkably regular patterns of peaks. Many stars in our sample (e.g., bottom half of Fig. 1) show a strong peak at very low frequency, which is likely to be the fundamental radial pressure mode ( $n = 1, l = 0$ ). This identification is strengthened by the fact that these peaks agree with the established period–luminosity relation for the fundamental radial mode in  $\delta$  Scuti stars<sup>29,32</sup>. In addition, several stars show a mode that is a factor  $\sim 0.78^{-1}$  higher in frequency, consistent with being the first radial overtone ( $n = 2, l = 0$ )<sup>33</sup>.

Figure 2 shows the pulsation spectra of several  $\delta$  Scuti stars in échelle format, where the spectra have been divided into equal segments that are stacked vertically so that the peaks with the same degree fall along ridges. The regularity of the patterns is striking, similar to échelle diagrams of solar-like oscillators<sup>1,2</sup>, but at much lower radial orders. Comparison with pulsation frequencies calculated from theoretical models (red symbols in the top row) enables an unambiguous mode identification, with ridges corresponding to sequences of radial modes ( $l = 0$ ) and dipolar modes ( $l = 1$ ), as shown. The ridge corresponding to  $l = 2$  modes does not appear to be present in these stars.

We have placed our sample in the Hertzsprung–Russell (H–R) diagram using effective temperatures and luminosities derived from broadband colors and *Gaia* parallaxes (Fig. 3). The  $\delta$  Scuti stars with regular frequency spacings tend to be located near the zero-age main-sequence, with masses between  $1.5$  and  $1.8 M_{\odot}$ . The fact that these stars are relatively young helps to explain their regular pulsation spectra. In more evolved stars, the nonradial modes are expected to be “bumped” from their regular spacings when they undergo avoided crossings due to coupling with gravity (buoyancy) modes in the core<sup>1,3</sup>. For young stars, this mode bumping only occurs at the lowest frequencies, as can be seen from the models of  $l = 1$  modes in Fig. 2 (top row; red triangles at low frequencies).

The large frequency separation,  $\Delta\nu$ , scales approximately as the square root of the mean stellar density<sup>1,19,20,22</sup>. However, the mode spacings of stars are not completely regular, even in the asymptotic regime, meaning that  $\Delta\nu$  varies with frequency. We used theoretical models to calculate  $\Delta\nu$  for  $\delta$  Scuti stars in the same way that we measured it, namely from radial modes with orders in the range  $n = 4$  to  $8$  (details in the Supplementary Information). We found that  $\Delta\nu$  in the models was typically 15% lower than would be obtained by scaling from the density of the

Sun. Figure 3b compares the observed large separations of our sample with the densities derived from fitting to evolutionary tracks in the H–R diagram. The results confirm there is a correlation, with most stars lying between the values based on the standard scaling relation (solid red curve) and those from the model calculations (dashed red curve). Some of the spread is probably due to the range of metallicities of the sample, and some will be due to rotation. For example, if a star is oblate then the inclination of its rotation axis affects the observed position in the H–R diagram (and hence the density).

High-resolution spectroscopy can be used to measure the projected rotational velocity of a star, and most intermediate-mass stars have  $v \sin i$  values<sup>34</sup> in the range 50–220 km s<sup>-1</sup>. Measurements of  $v \sin i$  are available for about one third of our sample, with some being newly measured and others coming from the literature (see Table S1). More than half are below 70 km s<sup>-1</sup>, indicating that this group of  $\delta$  Scuti stars have unusually low projected rotational velocities. This is consistent with the idea that regular frequency spacings tend to occur in stars seen at high inclinations (close to pole-on).

Some échelle diagrams show more complicated patterns, with additional ridges at various angles that indicate sequences with slightly different spacings (Fig. 4). The rotation axes of these stars are presumably at lower inclinations than those with simpler pulsation spectra, which would lead us to expect one  $l = 0$  ridge and three  $l = 1$  ridges. Beyond the usual rotational splitting of  $l = 1$  modes, slightly different frequency spacings are expected for each  $m$  in an oblate star. This is because modes with different  $m$  propagate along different paths through the star, giving different values for the sound-speed crossing time and hence for  $\Delta\nu$ . In stars with even more ridges, the additional sequences could correspond to modes with higher degrees ( $l \geq 2$ )—where coupling between modes with different degree may also be important—and perhaps also to chaotic modes<sup>10,35,36</sup>.


The detection of regular pulsation frequency patterns in intermediate-mass stars will expand the reach of asteroseismology to new frontiers, such as the ages of young moving groups, which are often uncertain by up to a factor of two<sup>37</sup>. Spectroscopic radial velocities and *Gaia* astrometry show that several stars in our sample are members of nearby young moving groups, including the Octans association<sup>38,39</sup> (HD 44930, HD 29783, HD 42915), the Carina association<sup>40</sup> (HD 89263), the Columba association<sup>41</sup> (HD 37286), and the recently-discovered Pisces–Eridanus stellar stream<sup>42</sup> (HD 31901). For the latter, gyrochronology yielded an age similar to the Pleiades ( $\sim 130$  Myr)<sup>43</sup>, in contrast to the initial  $\sim 1$  Gyr age determination from suspected evolved moving group members. Asteroseismic modelling of HD 31901 (Fig. 2c) clearly confirms a young age for this member of the Pisces–Eridanus group, and similar age determinations might be possible for other groups containing intermediate-mass stars.

Four stars in our sample (HD 28548, HD 34282, TYC 5945-497-1, V1790 Ori) exhibit excess emission in the WISE passbands, indicating a circumstellar dust disk. One of these (HD 34282) has


a disk that has been resolved by ALMA, showing it to be inclined  $60^\circ \pm 1^\circ$  to the line of sight<sup>44</sup>. The constraints on age and inclination of this host star provided by an analysis of its pulsations could illuminate the origin of stellar obliquity<sup>45</sup> and the pace of disk evolution<sup>46</sup>.

Finally, we note that six stars in our sample have been classified spectroscopically as  $\lambda$  Bootis stars (references given in Table S1), meaning that their surface chemical abundances show evidence for accretion from circumstellar material. Given that  $\lambda$  Bootis stars are rare, making up only about 2% of A stars<sup>47</sup>, the relatively high occurrence rate in our sample lends support to the hypothesis that  $\lambda$  Bootis stars tend to be young, with circumstellar material accreting from a proto-planetary disk.


Future *TESS* observations should reveal more examples of  $\delta$  Scuti stars that have clear sequences of pulsation overtones. We also suggest that these patterns can be used as a guide for mode identification in the much larger number of  $\delta$  Scuti stars whose pulsation spectra are not as regular.




**Figure 1 | Amplitude spectra of 15 high-frequency  $\delta$  Scuti stars observed with *TESS*.** The insets for some stars expand the vertical axis by a factor of 4 to make weaker peaks more visible.



**Figure 2 | Mode identification in  $\delta$  Scuti stars.** The amplitude spectra are shown in échelle format, with segments of equal length being stacked vertically. The vertical dashed line shows the value of  $\Delta\nu$  used in each case, with a repeated overlap region added on the right for clarity. The greyscale shows the observed amplitude spectrum, which in most cases is calculated from one 27-day sector of data from the *TESS* spacecraft. The exception is HD 187547, for which observations were made over 960 d with the *Kepler* spacecraft<sup>48</sup>. Smoothing was applied to the observed amplitude spectra before plotting. In the top row, the red symbols show mode frequencies calculated from theoretical models of non-rotating stars, chosen to match the observed modes reasonably well (see Supplementary Information for details of the models). These allow mode identifications in other stars as shown in the bottom row, where the red stripes mark overtones sequences of  $l = 0$  and  $l = 1$  modes. The parameters of the models shown in the top row are as follows (while noting that other values of the parameters also give fits of similar quality): **a**, HD 99506 Model parameters: mass  $1.69 M_{\odot}$ , metallicity  $[\text{Fe}/\text{H}] = -0.01$ , age 221 Myr, effective temperature 8038 K and radius  $1.51 R_{\odot}$ . **b**, HD 28548 Model parameters: mass  $1.59 M_{\odot}$ , metallicity  $[\text{Fe}/\text{H}] = -0.2$ , age 270 Myr, effective temperature 8202 K and radius  $1.41 R_{\odot}$ . **c**, HD 31901 Model parameters: mass  $1.77 M_{\odot}$ , metallicity  $[\text{Fe}/\text{H}] = 0.08$ , age 102 Myr, effective temperature 8083 K and radius  $1.51 R_{\odot}$ .



**Figure 3 | Physical properties of high-frequency  $\delta$  Scuti stars.** **a**, Location in the H–R diagram, colour-coded by the measured large-frequency separation. The small points show  $\delta$  Scuti stars observed by the *Kepler* Mission<sup>30</sup> and the red curves (labelled by mass in solar units) are evolutionary tracks calculated for solar metallicity (details in the Supplementary Information). The solid blue lines show the edges of the theoretical  $\delta$  Scuti instability strip<sup>49</sup> and the dashed magenta lines show the observed instability strip based on *Kepler* stars<sup>30</sup>. **b**, Large-frequency separation versus mean stellar density as determined from observations (symbols), as predicted from the standard scaling relation (solid red line) and from stellar models (red dashed line). Stars with spectroscopic observations are colour-coded by the measured projected rotational velocity ( $v \sin i$ ). The following stars are not shown: one star with  $A_V > 1.0$  mag (HD 290750), 7 stars with  $RUWE > 2.0$  (HD 29783, TYC 6515-7-1, HD 67688, HD 70510, HD 75040, HD 220811, HD 222496), and two stars that have companions within  $1.5''$  in the Washington Double Star Catalog (HD 25248, HR 4043).



**Figure 4 | Examples of more complicated échelle diagrams of  $\delta$  Scuti pulsations.** There are several sets of ridges at a range angles, indicating slightly different spacings. An intermediate value of  $\Delta\nu$  was chosen for these diagrams.

1. Aerts, C., Christensen-Dalsgaard, J. & Kurtz, D. W. *Asteroseismology* (Springer, 2010).
2. Garcia, R. A. & Ballot, J. Asteroseismology of solar-type stars. *arXiv e-prints* arXiv:1906.12262 (2019).
3. Hekker, S. & Christensen-Dalsgaard, J. Giant star seismology. *Astronomy and Astrophysics Review* **25**, 1 (2017).
4. Aerts, C. Massive Star Asteroseismology in Action. In Meynet, G., Georgy, C., Groh, J. & Stee, P. (eds.) *New Windows on Massive Stars*, vol. 307 of *IAU Symposium*, 154–164 (2015).
5. Córscico, A. H., Althaus, L. G., Miller Bertolami, M. M. & Kepler, S. O. Pulsating white dwarfs: new insights. *arXiv e-prints* arXiv:1907.00115 (2019).
6. Breger, M.  $\delta$  Scuti stars (Review). In *ASP Conf. Series, Vol. 210*, vol. 210 of *ASP Conf. Series*, 3 (2000).
7. Goupil, M. J. *et al.* Asteroseismology of  $\delta$  Scuti Stars: Problems and Prospects. *Journal of Astrophysics and Astronomy* **26**, 249 (2005).
8. Handler, G. Delta Scuti Variables. In Guzik, J. A. & Bradley, P. A. (eds.) *American Institute of Physics Conference Series*, vol. 1170, 403–409 (2009).
9. Ouazzani, R. M., Roxburgh, I. W. & Dupret, M. A. Pulsations of rapidly rotating stars. II. Realistic modelling for intermediate-mass stars. *Astron. Astrophys.* **579**, A116 (2015).
10. Reese, D. R. *et al.* Frequency regularities of acoustic modes and multi-colour mode identification in rapidly rotating stars. *Astron. Astrophys.* **601**, A130 (2017).
11. García Hernández, A. *et al.* Asteroseismic analysis of the CoRoT  $\delta$  Scuti star HD 174936. *Astron. Astrophys.* **506**, 79–83 (2009).
12. Breger, M. *et al.* Regularities in frequency spacings of  $\delta$  Scuti stars: the Kepler star KIC 9700322. *Mon. Not. R. Astron. Soc.* **414**, 1721–1731 (2011).
13. Antoci, V. *et al.* The excitation of solar-like oscillations in a  $\delta$ Sct star by efficient envelope convection. *Nature* **477**, 570–573 (2011).
14. Zwintz, K. *et al.* Regular frequency patterns in the classical  $\delta$  Scuti star HD 144277 observed by the MOST satellite. *Astron. Astrophys.* **533**, A133 (2011).
15. Zwintz, K. *et al.* Regular frequency patterns in the young  $\delta$  Scuti star HD 261711 observed by the CoRoT and MOST satellites. *Astron. Astrophys.* **552**, A68 (2013).
16. Páparó, M. *et al.* CoRoT 102749568: mode identification in a  $\delta$  Scuti star based on regular spacings. *Astron. Astrophys.* **557**, A27 (2013).
17. Casey, M. P. *et al.* MOST observations of the Herbig Ae  $\delta$ -Scuti star HD 34282. *Mon. Not. R. Astron. Soc.* **428**, 2596–2604 (2013).
18. García Hernández, A. *et al.* An in-depth study of HD 174966 with CoRoT photometry and HARPS spectroscopy. Large separation as a new observable for  $\delta$  Scuti stars. *Astron. Astrophys.* **559**, A63 (2013).

19. Suárez, J. C. *et al.* Measuring mean densities of  $\delta$  Scuti stars with asteroseismology. Theoretical properties of large separations using TOUCAN. *Astron. Astrophys.* **563**, A7 (2014).
20. García Hernández, A. *et al.* Observational  $\Delta\nu$ - $\rho$  Relation for  $\delta$  Sct Stars using Eclipsing Binaries and Space Photometry. *Astrophys. J.* **811**, L29 (2015).
21. Páparó, M., Benkő, J. M., Hareter, M. & Guzik, J. A. Unexpected Series of Regular Frequency Spacing of  $\delta$  Scuti Stars in the Non-asymptotic Regime. I. The Methodology. *Astrophys. J.* **822**, 100 (2016).
22. Páparó, M., Benkő, J. M., Hareter, M. & Guzik, J. A. Unexpected Series of Regular Frequency Spacing of  $\delta$  Scuti Stars in the Non-asymptotic Regime. II. Sample-Echelle Diagrams and Rotation. *The Astrophysical Journal Supplement Series* **224**, 41 (2016).
23. Michel, E. *et al.* What CoRoT tells us about  $\delta$  Scuti stars. Existence of a regular pattern and seismic indices to characterize stars. In *EPJWC*, vol. 160 of *EPJWC*, 03001 (2017).
24. Bowman, D. M. & Kurtz, D. W. Characterizing the observational properties of  $\delta$  Sct stars in the era of space photometry from the Kepler mission. *Mon. Not. R. Astron. Soc.* **476**, 3169–3184 (2018).
25. Reese, D., Lignières, F. & Rieutord, M. Acoustic oscillations of rapidly rotating polytropic stars. II. Effects of the Coriolis and centrifugal accelerations. *Astron. Astrophys.* **455**, 621–637 (2006).
26. Ricker, G. R. *et al.* Transiting Exoplanet Survey Satellite (TESS). *Journal of Astronomical Telescopes, Instruments, and Systems* **1**, 014003 (2015).
27. Rodríguez, E., López-González, M. J. & López de Coca, P. A revised catalogue of delta Sct stars. *Astron. Astrophys. Suppl. Ser.* **144**, 469–474 (2000).
28. Holdsworth, D. L. *et al.* High-frequency A-type pulsators discovered using SuperWASP. *Mon. Not. R. Astron. Soc.* **439**, 2078–2095 (2014).
29. Ziaali, E., Bedding, T. R., Murphy, S. J., Van Reeth, T. & Hey, D. R. The period-luminosity relation for  $\delta$  Scuti stars using Gaia DR2 parallaxes. *Mon. Not. R. Astron. Soc.* **486**, 4348–4353 (2019).
30. Murphy, S. J., Hey, D., Van Reeth, T. & Bedding, T. R. Gaia-derived luminosities of Kepler A/F stars and the pulsator fraction across the  $\delta$  Scuti instability strip. *Mon. Not. R. Astron. Soc.* **485**, 2380–2400 (2019).
31. Balona, L. A., Daszyńska-Daszkiewicz, J. & Pamyatnykh, A. A. Pulsation frequency distribution in  $\delta$  Scuti stars. *Mon. Not. R. Astron. Soc.* **452**, 3073–3084 (2015).
32. McNamara, D. H. Delta Scuti, SX Phoenicis, and RR Lyrae Stars in Galaxies and Globular Clusters. *Astron. J.* **142**, 110 (2011).
33. Petersen, J. O. & Christensen-Dalsgaard, J. Pulsation models of  $\delta$  Scuti variables. I. The high-amplitude double-mode stars. *Astron. Astrophys.* **312**, 463–474 (1996).

34. Zorec, J. & Royer, F. Rotational velocities of A-type stars. IV. Evolution of rotational velocities. *Astron. Astrophys.* **537**, A120 (2012).
35. Lignières, F. & Georgeot, B. Wave chaos in rapidly rotating stars. *Phys. Rev. E* **78**, 016215 (2008).
36. Lignières, F. & Georgeot, B. Asymptotic analysis of high-frequency acoustic modes in rapidly rotating stars. *Astron. Astrophys.* **500**, 1173–1192 (2009).
37. Mamajek, E. E. & Bell, C. P. M. On the age of the  $\beta$  Pictoris moving group. *Mon. Not. R. Astron. Soc.* **445**, 2169–2180 (2014).
38. Torres, C. A. O. *et al.* Search for associations containing young stars (SACY). I. Sample and searching method. *Astron. Astrophys.* **460**, 695–708 (2006).
39. Murphy, S. J. & Lawson, W. A. New low-mass members of the Octans stellar association and an updated 30-40 Myr lithium age. *Mon. Not. R. Astron. Soc.* **447**, 1267–1281 (2015).
40. Bell, C. P. M., Mamajek, E. E. & Naylor, T. A self-consistent, absolute isochronal age scale for young moving groups in the solar neighbourhood. *Mon. Not. R. Astron. Soc.* **454**, 593–614 (2015).
41. Zuckerman, B., Rhee, J. H., Song, I. & Bessell, M. S. The Tucana/Horologium, Columba, AB Doradus, and Argus Associations: New Members and Dusty Debris Disks. *Astrophys. J.* **732**, 61 (2011).
42. Meingast, S., Alves, J. & Fürnkranz, V. Extended stellar systems in the solar neighborhood . II. Discovery of a nearby 120 degree stellar stream in Gaia DR2. *Astron. Astrophys.* **622**, L13 (2019).
43. Curtis, J. L., Agüeros, M. A., Mamajek, E. E., Wright, J. T. & Cummings, J. D. TESS reveals that the nearby Pisces-Eridanus stellar stream is only 120 Myr old. *arXiv e-prints* arXiv:1905.10588 (2019).
44. van der Plas, G. *et al.* An 80 au cavity in the disk around HD 34282. *Astron. Astrophys.* **607**, A55 (2017).
45. Lai, D. Star-disc-binary interactions in protoplanetary disc systems and primordial spin-orbit misalignments. *Mon. Not. R. Astron. Soc.* **440**, 3532–3544 (2014).
46. Williams, J. P. & Cieza, L. A. Protoplanetary Disks and Their Evolution. *Ann. Rev. Astron. Astrophys.* **49**, 67–117 (2011).
47. Paunzen, E. A spectroscopic survey for lambda Bootis stars. III. Final results. *Astron. Astrophys.* **373**, 633–640 (2001).
48. Antoci, V. *et al.* The Role of Turbulent Pressure as a Coherent Pulsational Driving Mechanism: The Case of the  $\delta$  Scuti Star HD 187547. *Astrophys. J.* **796**, 118 (2014).
49. Dupret, M.-A., Grigahcène, A., Garrido, R., Gabriel, M. & Scuflaire, R. Theoretical instability strips for  $\delta$  Scuti and  $\gamma$  Doradus stars. *Astron. Astrophys.* **414**, L17–L20 (2004).

**Acknowledgements** We gratefully acknowledge support from the Australian Research Council (grant DE 180101104), and from the Danish National Research Foundation (Grant DNRF106) through its funding for the Stellar Astrophysics Center (SAC). D.H. acknowledges support by the National Aeronautics and Space Administration under Grant 80NSSC18K1585 awarded through the TESS Guest Investigator Program and by the National Science Foundation (AST-1717000). Y.L. acknowledges the Joint Research Fund in Astronomy (U1631236) under cooperative agreement between the National Natural Science Foundation of China (NSFC) and Chinese Academy of Sciences (CAS). We also acknowledge the *TESS* and *Kepler* teams, whose efforts made these results possible. This research was partially conducted during the Exostar19 program at the Kavli Institute for Theoretical Physics at UC Santa Barbara, which was supported in part by the National Science Foundation under Grant No. NSF PHY-1748958. We thank colleagues in that program, especially Rich Townsend, for many stimulating discussions. This work makes use of observations from the LCOGT network. The authors wish to recognize and acknowledge the very significant cultural role and reverence that the summit of Mauna Kea has always had within the indigenous Hawaiian community. We are most fortunate to have the opportunity to conduct observations from this mountain.