

Detection and characterisation of oscillating red giants: first results from the TESS satellite

VÍCTOR SILVA AGUIRRE,¹ DENNIS STELLO,^{2,3,1} AMALIE STOKHOLM,¹ JAKOB R. MOSUMGAARD,¹ WARRICK H. BALL,^{4,1} SARBANI BASU,⁵ DIEGO BOSSINI,⁶ LISA BUGNET,^{7,8} DEREK BUZASI,⁹ TIAGO L. CAMPANTE,^{6,10} LINDSEY CARBONEAU,⁹ WILLIAM J. CHAPLIN,^{4,1} ENRICO CORSARO,¹¹ GUY R. DAVIES,^{4,1} YVONNE ELSWORTH,^{4,1} RAFAEL A. GARCÍA,^{7,8} PATRICK GAULME,^{12,13} OLIVER J. HALL,^{4,1} RASMUS HANDBERG,¹ MARC HON,² THOMAS KALLINGER,¹⁴ LIU KANG,¹⁵ MIKKEL N. LUND,¹ SAVITA MATHUR,^{16,17} ALEXEY MINTS,¹⁸ BENOIT MOSSER,¹⁹ ZEYNEP ÇELİK ORHAN,²⁰ THAÍSE S. RODRIGUES,²¹ MATHIEU VRARD,^{6,22} MUTLU YILDIZ,²⁰ JOEL C. ZINN,^{2,23,22} SIBEL ÖRTEL,²⁰ PAUL G. BECK,^{24,16,17} KEATON J. BELL,^{25,26} ZHAO GUO,²⁷ CHEN JIANG,²⁸ JAMES S. KUSZLEWICZ,¹² CHARLES A. KUEHN,²⁹ TANDA LI,^{3,1,4} MIA S. LUNDKVIST,¹ MARC PINSONNEAULT,²² JAMIE TAYAR,^{30,31} MARGARIDA S. CUNHA,^{6,4} SASKIA HEKKER,^{12,1} DANIEL HUBER,³⁰ ANDREA MIGLIO,^{4,1} MARIO J. P. F. G. MONTEIRO,^{6,10} DITTE SLUMSTRUP,^{1,32} MARK L. WINTHER,¹ GEORGE ANGELOU,³³ OTHMAN BENOMAR,^{34,35} ATTILA BÓDI,^{36,37} BRUNO L. DE MOURA,³⁸ SÉBASTIEN DEHEUVELS,³⁹ ALIZ DEREKAS,^{40,41,36} MARIA PIA DI MAURO,⁴² MARC-ANTOINE DUPRET,⁴³ ANTONIO JIMÉNEZ,^{16,17} YVELINE LEBRETON,^{19,44} JAYMIE MATTHEWS,⁴⁵ NICOLAS NARDETTO,⁴⁶ JOSE D. DO NASCIMENTO, JR.,^{47,48} FILIPE PEREIRA,^{6,10} LUISA F. RODRÍGUEZ DÍAZ,¹ ALDO M. SERENELLI,^{49,50} EMANUELE SPITONI,¹ EDITA STONKUTÉ,⁵¹ JUAN CARLOS SUÁREZ,^{52,53} ROBERT SZABÓ,^{36,37} VINCENT VAN EYLEN,⁵⁴ RITA VENTURA,¹¹ KULDEEP VERMA,¹ ACHIM WEISS,³³ TAO WU,^{55,56,57} THOMAS BARCLAY,^{58,59} JØRGEN CHRISTENSEN-DALSGAARD,¹ JON M. JENKINS,⁶⁰ HANS KJELDEN,¹ GEORGE R. RICKER,⁶¹ SARA SEAGER,^{61,62,63} AND ROLAND VANDERSPEK⁶¹

¹*Stellar Astrophysics Centre (SAC), Department of Physics and Astronomy, Aarhus University, Ny Munkegade 120, 8000 Aarhus C, Denmark*

²*School of Physics, The University of New South Wales, Sydney NSW 2052, Australia*

³*Sydney Institute for Astronomy (SfA), School of Physics, University of Sydney, NSW 2006, Australia*

⁴*School of Physics and Astronomy, University of Birmingham, Birmingham, B15 2TT, United Kingdom*

⁵*Department of Astronomy, Yale University, New Haven, CT 06520, USA*

⁶*Instituto de Astrofísica e Ciências do Espaço, Universidade do Porto, Rua das Estrelas, 4150-762 Porto, Portugal*

⁷*IRFU, CEA, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France*

⁸*AIM, CEA, CNRS, Université Paris-Saclay, Université Paris Diderot, Sorbonne Paris Cité, F-91191 Gif-sur-Yvette, France*

⁹*Dept. of Chemistry and Physics, Florida Gulf Coast University, 10501 FGCU Blvd. S., Fort Myers, FL 33965 USA*

¹⁰*Departamento de Física e Astronomia, Faculdade de Ciências da Universidade do Porto, Rua do Campo Alegre, s/n, 4169-007 Porto, Portugal*

¹¹*INAF Osservatorio Astrofisico di Catania, Via S. Sofia, 78, 95123 Catania, Italy*

¹²*Max Planck Institute for Solar System Research, Justus-von-Liebig Weg 3, D-37077 Göttingen, Germany*

¹³*Department of Astronomy, New Mexico State University, P.O. Box 30001, MSC 4500, Las Cruces, NM 88003-8001, USA*

¹⁴*Institut für Astrophysik, Universität Wien, Türkenschanzstrasse 17, 1180 Vienna, Austria*

¹⁵*Department of Astronomy, Beijing Normal University, 100875 Beijing, PR China*

¹⁶*Instituto de Astrofísica de Canarias, E-38200 La Laguna, Tenerife, Spain*

¹⁷*Departamento de Astrofísica, Universidad de La Laguna, E-38206 La Laguna, Tenerife, Spain*

¹⁸*Leibniz-Institut für Astrophysik Potsdam (AIP), An der Sternwarte 16, 14482 Potsdam, Germany*

¹⁹*LESIA, Observatoire de Paris, Université PSL, CNRS, Sorbonne Université, Université de Paris, 92195 Meudon, France*

²⁰*Department of Astronomy and Space Sciences, Science Faculty, Ege University, 35100, Bornova, İzmir, Turkey*

²¹*Osservatorio Astronomico di Padova – INAF, Vicolo dell’Osservatorio 5, I-35122 Padova, Italy*

²²*Department of Astronomy, The Ohio State University, 140 West 18th Avenue, Columbus OH 43210, USA*

²³*Kavli Institute for Theoretical Physics, University of California, Santa Barbara, CA 93106, USA*

²⁴*Institute of Physics, Karl-Franzens University of Graz, NAWI Graz, Universitätsplatz 5/II, 8010 Graz, Austria*

²⁵*DIRAC Institute, Department of Astronomy, University of Washington, Seattle, WA 98195-1580, USA*

²⁶*NSF Astronomy and Astrophysics Postdoctoral Fellow and DIRAC Fellow*

²⁷*Center for Exoplanets and Habitable Worlds, Department of Astronomy and Astrophysics, 525 Davey Laboratory, The Pennsylvania State University, University Park, PA16802, USA*

²⁸*School of Physics and Astronomy, Sun Yat-Sen University, No. 135, Xingang Xi Road, Guangzhou, 510275, P. R. China*

²⁹*Department of Physics and Astronomy, University of Northern Colorado, Greeley, CO 80639, USA*

³⁰*Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, Hawaii 96822, USA*

³¹ *Hubble Fellow*³² *European Southern Observatory, Alonso de Córdova 3107, Vitacura, Santiago, Chile*³³ *Max-Planck-Institut für Astrophysics, Karl Schwarzschild Strasse 1, 85748, Garching, Germany*³⁴ *Solar Science Observatory, NAOJ, Mitaka, Japan*³⁵ *Center for Space Science, New York University Abu Dhabi, UAE*³⁶ *Konkoly Observatory, Research Centre for Astronomy and Earth Sciences, H-1121 Budapest, Konkoly Thege M. út 15-17, Hungary*³⁷ *MTA CSFK Lendület Near-Field Cosmology Research Group*³⁸ *Instituto Federal do R. G. do Norte - IFRN, Brazil*³⁹ *IRAP, Université de Toulouse, CNRS, CNES, UPS, (Toulouse), France*⁴⁰ *ELTE Eötvös Loránd University, Gothard Astrophysical Observatory, Szombathely, Hungary*⁴¹ *MTA-ELTE Exoplanet Research Group, 9700 Szombathely, Szent Imre h. u. 112, Hungary*⁴² *INAF-IAPS, Istituto di Astrofisica e Planetologia Spaziali, Via del Fosso del Cavaliere 100, 00133 Roma, Italy*⁴³ *STAR Institute, University of Lige, 19C Alle du 6 Aot, B-4000 Lige, Belgium*⁴⁴ *Univ Rennes, CNRS, IPR (Institut de Physique de Rennes) - UMR 6251, F-35000 Rennes, France*⁴⁵ *Department of Physics and Astronomy, University of British Columbia, Vancouver, Canada*⁴⁶ *Université Côte d'Azur, Observatoire de la Côte d'Azur, CNRS, Laboratoire Lagrange, France*⁴⁷ *Universidade Federal do Rio Grande do Norte, UFRN, Departamento de Física, 59078-970, Natal, RN, Brazil*⁴⁸ *Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138, USA*⁴⁹ *Instituto de Ciencias del Espacio (ICE, CSIC), Campus UAB, Carrer de Can Magrans, s/n, 08193 Cerdanyola del Valles, Spain*⁵⁰ *Institut d'Estudis Espacials de Catalunya (IEEC), Gran Capita 4, E-08034, Barcelona, Spain*⁵¹ *Institute of Theoretical Physics and Astronomy, Vilnius University, Saulėtekio al. 3, 10257 Vilnius, Lithuania*⁵² *Dept. Física Teórica y del Cosmos. Universidad de Granada. 18006 Granada, Spain*⁵³ *Instituto de Astrofísica de Andalucía (CSIC). Glorieta de la Astronomía s/n. 18008. Granada, Spain*⁵⁴ *Mullard Space Science Laboratory, University College London, Holmbury St Mary, Dorking RH5 6NT, UK*⁵⁵ *Yunnan Observatories, Chinese Academy of Sciences, 396 Yangfangwang, Guandu District, Kunming, 650216, P. R. China*⁵⁶ *Key Laboratory for the Structure and Evolution of Celestial Objects, Chinese Academy of Sciences, 396 Yangfangwang, Guandu District, Kunming, 650216, P. R. China*⁵⁷ *Center for Astronomical Mega-Science, Chinese Academy of Sciences, 20A Datun Road, Chaoyang District, Beijing, 100012, P. R. China*⁵⁸ *NASA Goddard Space Flight Center, 8800 Greenbelt Rd, Greenbelt, MD 20771, USA*⁵⁹ *University of Maryland, Baltimore County, 1000 Hilltop Cir, Baltimore, MD 21250, USA*⁶⁰ *NASA Ames Research Center, Moffett Field, CA, 94035, USA*⁶¹ *Department of Physics and Kavli Institute for Astrophysics and Space Research, Massachusetts Institute of Technology, Cambridge, MA 02139, USA*⁶² *Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, MA 02139, USA*⁶³ *Department of Aeronautics and Astronautics, MIT, 77 Massachusetts Avenue, Cambridge, MA 02139, USA*

(Received January 1, 2018; Revised January 7, 2018; Accepted January 27, 2020)

Submitted to ApJL

ABSTRACT

Since the onset of the ‘space revolution’ of high-precision high-cadence photometry, asteroseismology has been demonstrated as a powerful tool for informing Galactic archaeology investigations. The launch of the NASA TESS mission has enabled seismic-based inferences to go full sky – providing a clear advantage for large ensemble studies of the different Milky Way components. Here we demonstrate its potential for investigating the Galaxy by carrying out the first asteroseismic ensemble study of red giant stars observed by TESS. We use a sample of 25 stars for which we measure their global asteroseismic observables and estimate their fundamental stellar properties, such as radius, mass, and age. Significant improvements are seen in the uncertainties of our estimates when combining seismic observables from TESS with astrometric measurements from the Gaia mission compared to when the seismology and astrometry are applied separately. Specifically, when combined we show that stellar radii can be determined to a precision of a few percent, masses to 5-10% and ages to the 20% level. This is comparable to the precision typically obtained using end-of-mission *Kepler* data.

Keywords: asteroseismology — stars: fundamental parameters — techniques: photometric

1. INTRODUCTION

Asteroseismology of red giant stars has been one of the major successes of the CoRoT and *Kepler* missions.

The unambiguous detection of non-radial oscillations has fundamentally widened our understanding of the inner workings of red giants, including the conditions in their core (e.g., Bedding et al. 2011). The observed frequency spectra have allowed the determination of the physical properties of thousands of red giants to an unprecedented level of precision (e.g., Miglio et al. 2013), paving the way for the emergence of asteroseismology as a powerful tool for Milky Way studies and Galactic archaeology (e.g., Miglio et al. 2009; Casagrande et al. 2016; Anders et al. 2017; Silva Aguirre et al. 2018; Sharma et al. 2019). The *Transiting Exoplanet Survey Satellite* (TESS, Ricker et al. 2015) is on the path of continuing this legacy with its all-sky survey that is expected to increase the number of detected oscillating red giants by an order of magnitude compared to the tens of thousands reported by its predecessors *CoRoT* and *Kepler*.

In the nominal TESS mission, the ecliptic northern and southern hemispheres are each observed during thirteen 27-day-long sectors, and most (92%) of the surveyed sky will be monitored for just 1-2 sectors. Except for the 20,000 targets pre-selected in each sector for 2-min cadence observations, all stars are observed as part of the full frame images obtained in 30-min cadence, similar to the long cadence sampling of the *Kepler* satellite. The length of the observations sets the lower limit on the oscillation frequencies one can resolve, and the sampling sets the upper frequency limit. We know from previous *Kepler* observations that one month of 30-min cadence data should be well suited to detect oscillations in the low red-giant branch and sufficient to measure the global oscillation properties characterising the frequency spectrum, in particular, its frequency of maximum power, ν_{\max} , and the frequency separation between overtone modes, $\Delta\nu$ (Bedding et al. 2010). These in turn can be used in combination with complementary data such as the effective temperature, T_{eff} , the relative iron abundance, $[\text{Fe}/\text{H}]$, and parallax, to obtain precise stellar properties (including ages) when applying asteroseismic-based grid modelling approaches (see e.g., Rodrigues et al. 2017; Pinsonneault et al. 2018).

Due to the large sky coverage, approximately 97% of asteroseismic detections in red giants from the TESS nominal mission data are expected to come from stars observed for only one or two sectors¹. Here we set out to explore the capability of TESS to detect the oscillations in giants ranging from the base of the red giant branch to the red clump, determine their stellar properties, and use that to assess the prospects for Galactic archaeology studies using one to two sectors of TESS data.

2. TARGET SELECTION




Figure 1. ‘Asteroseismic HR diagram’ showing (predicted) ν_{\max} instead of luminosity. Red dots show the selected targets inside the black selection box. For reference, the Sun is shown as well as all *Hipparcos* stars brighter than 6th magnitude (grey dots). Solar metallicity MESA tracks from Stello et al. (2013) are shown to guide the eye with masses in solar units indicated (pre- and post-helium core-ignition phases are shown separately).

Our goal is to have a representative sample of giants including the types of stars in which we can expect to detect oscillations from one sector 30-min cadence TESS data. We selected red-giant candidates observed during sectors 1 and/or 2 that were deemed viable for asteroseismic detections according to their predicted properties based on the *Hipparcos* catalogue (Van Leeuwen 2007). We first estimated the stellar T_{eff} and luminosity using $B - V$ color, V -band, and *Hipparcos* parallax, and the color-temperature and bolometric correction relations of Flower (1996). We then obtained a prediction of ν_{\max} ($\propto T_{\text{eff}}^{3.5} M/L$; solar scaled, e.g. Yu et al. 2018) for each star assuming a mass of $1.2 M_{\odot}$, which is representative of a typical red giant as observed by *Kepler* (and unlikely to be more than a factor of two from the true value of each star, e.g. Yu et al. 2018). We note that one of our targets (TIC 129649472) is a known exoplanet host star recently analysed by Campante et al. (2019).

To ensure that the selected targets were amenable to asteroseismic detection from one sector of 30-min cadence data, we required that they would have an expected ν_{\max} in the range $30\text{--}220\mu\text{Hz}$ and T_{eff} in the typical range of red giants of $4500\text{--}5200\text{ K}$. In addition, we applied a narrower T_{eff} range of $4500\text{--}4700\text{ K}$ for the stars with ν_{\max} between $30\mu\text{Hz}$ and $70\mu\text{Hz}$, to avoid having red clump stars dominating our sample. The resulting

¹ Based on a preliminary simulation of the full TESS sky (TESS GI Proposal No G011188).

sample of stars span evolutionary phases from the base of the red giant branch to the red giant branch bump, as well as some clump stars.

From this sample, we selected the 25 brightest targets for light curve extraction and asteroseismic analysis. The faintest stars in our sample turned out to be ~ 6 -7th magnitude in V band (see Table 1). Under the assumption that the photometric performance of TESS is similar to *Kepler*'s, apart from its smaller aperture, this magnitude limit is equivalent to 11-12th magnitude for *Kepler*. Because single-quarter observations from *Kepler*'s second life, K2, showed no oscillation detection bias for red giants brighter than around 12th magnitude (Stello et al. 2017) we would expect to detect oscillations in all 25 giants with TESS.

Figure 1 illustrates the location of the selected stars in the HR-diagram and the applied selection criteria. We confirmed that the stars were in sectors 1-2 using the Web TESS Viewing tool (WTV)².

3. DATA PROCESSING AND ASTEROSEISMIC ANALYSIS

The stars selected were included in an early release of processed data from the TASOC pipeline³. The calibrated full frame images were produced by the TESS Science Processing Operations Center (SPOC) at NASA Ames Research Center (Jenkins et al. 2016), and processed by combining the methodology from the K2P2 pipeline (Lund et al. 2015) for extracting the flux from target pixel data with the KASOC filter for systematics correction (Handberg & Lund 2014). The resulting TASOC light curves were high-pass filtered using a filter width of 4 days, corresponding to a cut-off frequency of approximately $3\mu\text{Hz}$, and 4σ outliers were removed. Finally, we used linear interpolation to fill gaps that lasted up to three consecutive cadences and derived the Fourier transforms (power frequency spectra) of each light curve.

The light curves for the seven stars observed in both sectors were merged. To follow the approach anticipated for the millions of light curves from the TESS full frame images in the future, we first applied the neural network-based detection algorithm by Hon et al. (2018) resulting in detection of oscillations in the power spectra of all stars except one. The non-detection (TIC 204314449) is listed as an A2 dwarf and a 'Visual Double' in the University of Michigan Catalogue of two-dimensional spectral types for the HD stars (Houk 1994), and hence possibly too hot to show solar-like oscillations, or potentially contaminated. For the current test case, the number of stars was small enough that we visually checked the results, which confirmed all detections and the non-




Figure 2. Power spectra sample of our targets representative of the ν_{\max} range that they cover from around the red clump (top) to the low luminosity red giant branch (bottom). *Left:* Spectra shown in log-log space (smoothed in red) showing the location of the oscillation power excess, ν_{\max} , indicated by red arrows on top of a frequency-dependent granulation background and flat white noise component. *Right:* Close-up of spectra showing locations of the roughly equally-spaced radial modes (using red equally-spaced vertical lines to guide the eye) and their average separation, $\Delta\nu$ (red horizontal arrows). In the three bottom panels multiple dipole ($l = 1$) mixed modes are resolved in between consecutive radial modes as indicated by the black brackets.

² <https://heasarc.gsfc.nasa.gov/cgi-bin/tess/webtess/wtv.py>

³ T'DA Data Release Notes - Data Release 3 for TESS Sectors 1+2 (<https://doi.org/10.5281/zenodo.2510028>)

detection. The power spectra of a representative sample of the stars are shown in Figure 2 showing clear oscillation excess power and the frequency pattern required to measure both ν_{\max} and $\Delta\nu$.

The neural network also supplies a rough estimate for ν_{\max} , which we provided as a prior to 13 independent groups analysing the power spectra to extract high-precision values of both ν_{\max} , $\Delta\nu$, and their respective uncertainties using their preferred method. These methods have been thoroughly tested and described in the literature (see e.g., Huber et al. 2009; Gaulme et al. 2009; Hekker et al. 2010; Mathur et al. 2010; Mosser et al. 2011; Kallinger et al. 2012; Corsaro & De Ridder 2014; Davies et al. 2016; Campante et al. 2017; Zinn et al. 2019).

From the 13 independent determinations of the global asteroseismic parameters we adopted as central reference value for $\Delta\nu$ and ν_{\max} the results from the pipeline by Gaulme et al. (2009), as this method was on average closest to the ensemble mean after applying a $2\text{-}\sigma$ outlier rejection. Uncertainties in the global asteroseismic parameters obtained by the selected pipeline are at the 1.9% and 2.4% level for $\Delta\nu$ and ν_{\max} , respectively. These uncertainties are of comparable magnitude to those obtained from a single campaign with the K2 mission (see appendix in Stello et al. 2017) and about twice as large as those extracted from 50 days of *Kepler* observations (see Figs. 3 and 4 in Hekker et al. 2012). We report the central values and statistical uncertainties in $\Delta\nu$ and ν_{\max} from the selected pipeline for all targets in Table 1.

For each star, we take into account the scatter across the different methods by adding in quadrature the standard deviation among the central values retained after the $2\text{-}\sigma$ outlier rejection procedure to the formal uncertainty reported by the selected reference method. This consolidation process yields median uncertainties of 3.9% in $\Delta\nu$ and 2.6% in ν_{\max} , where the individual contribution arising from this systematic component to the total uncertainty is listed in Table 1. We note that we could decrease the level of uncertainties resulting from our ‘blind’ statistical consolidation approach by for example checking the $\Delta\nu$ and ν_{\max} results against the power spectra and/or échelle diagrams (see Fig. 5 in Stello et al. 2011). However, we want to draw a realistic picture of the uncertainties one can expect when dealing with large ensembles of stars (as expected from TESS) where detailed ‘boutique’ analysis/checking on a star-by-star basis is not practically feasible. Hence, our quoted uncertainties are conservative, but representative for analysis of TESS red giants where several pipelines are involved.

4. DERIVED STELLAR PROPERTIES

We have determined stellar properties for a subsample of 17 stars that had spectroscopic measurements of effective temperature and chemical composition available

in the literature. Since one of our goals is to follow the same analysis procedure expected for large ensembles of stars, we assumed fixed uncertainties in T_{eff} and $[\text{Fe}/\text{H}]$ of 80 K and 0.08 dex, which are at the level of those provided by current large-scale spectroscopic surveys. To extract the physical properties of our sample, the atmospheric information was complemented with the asteroseismic scaling relations:

$$\left(\frac{\Delta\nu}{\Delta\nu_{\odot}}\right)^2 \simeq \frac{\bar{\rho}}{\bar{\rho}_{\odot}} \quad (1)$$

$$\left(\frac{\nu_{\max}}{\nu_{\max,\odot}}\right) \simeq \frac{M}{M_{\odot}} \left(\frac{R}{R_{\odot}}\right)^{-2} \left(\frac{T_{\text{eff}}}{T_{\text{eff},\odot}}\right)^{1/2}, \quad (2)$$

where we adopted $\Delta\nu_{\odot} = 135.5$ (μHz) and $\nu_{\max,\odot} = 3140$ (μHz) as obtained by our reference pipeline from the analysis of solar data.

Seven teams independently applied grid-based modelling pipelines based on stellar evolution models or isochrones to determine the main physical properties of the targets (see Basu et al. 2012; Silva Aguirre et al. 2015; Rodrigues et al. 2017; Mints & Hekker 2018; Yıldız et al. 2019, and references therein). When matching the models to the atmospheric properties and the global asteroseismic parameters $\Delta\nu$ and ν_{\max} the pipelines yielded median uncertainties of $\sim 6\%$ in radius, $\sim 14\%$ in mass, and $\sim 50\%$ in age. These statistical uncertainties are of the same magnitude to those obtained with the K2 mission (Sharma et al. 2019), as expected from the similar resulting errors in the global seismic parameters described in Section 3, and about a factor of two larger than what can be achieved with the full duration of the *Kepler* observations (Pinsonneault et al. 2018).

In addition to the asteroseismic information, five of the pipelines can include parallaxes from Gaia DR2 (Gaia Collaboration et al. 2018) coupled with Tycho-2 (Høg et al. 2000) observed V -magnitudes in their fitting algorithm to further constrain the stellar properties. As a consequence of having the additional constraint on stellar radius from the astrometry, the resulting uncertainties decrease to a level of $\sim 3\%$ in radius, $\sim 6\%$ in mass, and $\sim 20\%$ in age. This level of precision resembles that obtained with the use of the full length of asteroseismic observations from the nominal *Kepler* mission, and emphasizes the potential of TESS for Galactic studies using red giants given its larger sky coverage, simple and reproducible selection function, and one order of magnitude higher expected yield of asteroseismic detections than any other previous mission.

To illustrate the differences in the obtained stellar properties arising from the selection of fitted observables, Fig. 3 shows the stellar radius obtained with one of the pipelines (BASTA, Silva Aguirre et al. 2015) when fitting different combinations of input parameters. The figure uses as the reference value the case when, in addition to the atmospheric properties, only the Gaia DR2




Figure 3. Comparison of stellar radii obtained with BASTA when fitting different combinations of input parameters: Gaia DR2 parallax and V -band magnitude (ϖ), global asteroseismic parameters ($\Delta\nu, \nu_{\max}$), and all combined ($\varpi, \Delta\nu, \nu_{\max}$). Effective temperature and composition are also fitted in all cases. See text for details.

parallax and observed V -band magnitude are included in the fit. For the majority of the targets the results are consistent across the three sets within their formal statistical uncertainties. A summary of the measured and derived stellar properties for our targets can be found in Table 1, where we have listed the central values and statistical uncertainties obtained with the BASTA pipeline, and determined the systematic contribution as the standard deviation across the results reported by all pipelines.

Two targets (TIC 141280255 and TIC 149347992) present a larger disagreement between the radii obtained with parallax and the seismic set ($\Delta\nu, \nu_{\max}$). We investigated if these discrepancies were due to the quality of the astrometric data by computing the renormalised unit weight error (RUWE⁴) for our sample of stars. In the case of TIC 141280255 we obtained a RUWE=1.98, which is above the value recommended by the Gaia team as a criterion for a good astrometric solution (RUWE ≤ 1.4). Therefore, we adopt for this star the stellar properties obtained from fitting the asteroseismic input only ($\Delta\nu, \nu_{\max}$).

In the case of TIC 149347992 the discrepancy is the result of predicted evolutionary phases: while the parallax-only solution suggests that the star is in the clump phase, the asteroseismic fit favours a star in the red-giant branch. The combined fit therefore presents a bimodal distribution that encompasses these two families of solutions. A similar situation occurs in the fit




Figure 4. Distribution of fractional age uncertainties for our sample of stars determined by the BASTA pipeline fitting different combinations of available observables. The points indicate the individual values used to construct the Gaussian kernel density estimation. For better visualization we have excluded TIC 175375523 from the figure. See text for details.

of TIC 175375523, which shows agreement in the radius determined from different sets of input but has a fractional age uncertainty above unity when only ($\Delta\nu, \nu_{\max}$) are included in the fit. Its resulting age distribution is bimodal in this set as both red-giant branch and clump models can reproduce the observations, but the inclusion of parallax information favours the red giant branch solution and accounts for the $\sim 17\%$ statistical uncertainty reported in Table 1. The availability of evolutionary classifications from deep neural networks trained on short *Kepler* data (Hon et al. 2018) would further decrease the obtained uncertainties by clearly disentangling these two scenarios.

In Fig. 4 we plot the distribution of fractional age uncertainties obtained with BASTA for the three considered cases of input, showing the clear improvement in precision when asteroseismic information and parallax are simultaneously included in the fit. For visualization purposes we have excluded the target TIC 175375523 from the figure. Our stellar ages at the 20% level are significantly more precise than what is obtained by data-driven and neural-network methods trained using asteroseismic ages from *Kepler* (above the 30% level, see e.g., Mackereth et al. 2019). As a final remark, we note that asteroseismically derived properties of red giants are accurate to at least a similar level than our statistical uncertainties (below $\sim 5\%$ and $\sim 10\%$ for radii and masses, respectively. See discussion in e.g., Pinsonneault et al. 2018, and references therein). We have made emphasis on our achieved precision instead of accuracy as our results could still be affected by a systematic component arising from uncertainties in evolutionary calculations, although recent investigations quantifying these effects

⁴ see Gaia technical note GAIA-C3-TN-LU-LL-124-01 (<https://www.cosmos.esa.int/web/gaia/dr2-known-issues>)

at solar metallicity suggest that they are smaller than our statistical uncertainties (Silva Aguirre et al. 2019).

5. CONCLUSIONS

We presented the first ensemble analysis of red giants stars observed with the TESS mission. We selected a sample of 25 stars where we expected to detect oscillations based on their magnitude and parallax value, and analysed the extracted light curves in search for asteroseismic signatures in the power spectra. Our main findings can be summarized as follows:

- We detected oscillations in all the stars (except one that was likely incorrectly listed as a red giant). Despite the modest number of stars in our sample, our detection yield supports that the TESS photometric performance is similar to that of *Kepler* and K2 except shifted by about 5 magnitudes towards brighter stars due to its smaller aperture.
- Individual pipelines retrieve the global asteroseismic parameters with uncertainties at the $\sim 2\%$ level in $\Delta\nu$ and $\sim 2.5\%$ in ν_{\max} , which respectively increase to $\sim 4\%$ and $\sim 3.5\%$ when we take into account the scatter across results. We consider these uncertainties to be representative for the forthcoming ensemble analysis of TESS targets observed in 1-2 sectors, as individual validation of the results will not be feasible due to the large number of targets observed.
- Grid-based modelling techniques applying asteroseismic scaling relations were used to retrieve stellar properties for the 17 targets with spectroscopic information. Radii, masses, and ages were obtained with uncertainties at the 6%, 14%, and 50% level, and decrease to 3%, 6%, and 20% when parallax information from Gaia DR2 is included.

The expected number of red giants with detected oscillations by TESS ($\sim 500,000^5$) greatly surpasses the final yield of *Kepler* ($\sim 20,000$). In this respect, the combination of TESS observations, Gaia astrometry, and large scale spectroscopic surveys holds a great potential for studies of Galactic structure where precise stellar properties (particularly ages) are of key importance. We note that the recently approved extended TESS mission will change the 30-min cadence to 10 minutes, making it possible to detect oscillations of stars of smaller radii using the full frame images. This will enable more rigorous investigations of the asteroseismic mass scale for giants

when anchored to empirical mass determinations (e.g., from eclipsing binaries) of turn-off and subgiant stars.

This paper includes data collected by the TESS mission, which are publicly available from the Mikulski Archive for Space Telescopes (MAST). Funding for the TESS mission is provided by NASAs Science Mission directorate. Funding for the TESS Asteroseismic Science Operations Centre is provided by the Danish National Research Foundation (Grant agreement no.: DNRFF106), ESA PRODEX (PEA 4000119301) and Stellar Astrophysics Centre (SAC) at Aarhus University. VSA acknowledges support from the Independent Research Fund Denmark (Research grant 7027-00096B). DB is supported in the form of work contract FCT/MCTES through national funds and by FEDER through COMPETE2020 in connection to these grants: UID/FIS/04434/2019; PTDC/FIS-AST/30389/2017 & POCI-01-0145-FEDER-030389. LB, RAG and BM acknowledge the support from the CNES/PLATO grant. DB acknowledges NASA grant NNX16AB76G. TLC acknowledges support from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 792848 (PULSATION). This work was supported by FCT/MCTES through national funds (UID/FIS/04434/2019). EC is funded by the European Unions Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No. 664931. RH and MNL acknowledge the support of the ESA PRODEX programme. T.S.R acknowledges financial support from Premiale 2015 MITiC (PI B. Garilli). KJB is supported by the National Science Foundation under Award AST-1903828. MSL is supported by the Carlsberg Foundation (Grant agreement no.: CF17-0760). MC is funded by FCT//MCTES through national funds and by FEDER through COMPETE2020 through these grants: UID/FIS/04434/2019, PTDC/FIS-AST/30389/2017 & POCI-01-0145-FEDER-030389, CEECIND/02619/2017. The research leading to the presented results has received funding from the European Research Council under the European Community’s Seventh Framework Programme (FP7/2007-2013) / ERC grant agreement no 338251 (StellarAges). AM acknowledges support from the European Research Council Consolidator Grant funding scheme (project ASTEROCHRONOMETRY, G.A. n. 772293, <http://www.asterochronometry.eu>). AMS is partially supported by MINECO grant ESP2017-82674-R. JCS acknowledges funding support from Spanish public funds for research under projects ESP2017-87676-2-2, and from project RYC-2012-09913 under the Ramn y Cajal program of the Spanish Ministry of Science and Education. Resources supporting this work were provided by the NASA High-End Computing (HEC) Program through the NASA Advanced

⁵ Based on a preliminary simulation of the full TESS sky (TESS GI Proposal No G011188).

Supercomputing (NAS) Division at Ames Research Center for the production of the SPOC data products.

REFERENCES

- Alves, S., Benamati, L., Santos, N. C., et al. 2015, *Monthly Notices of the Royal Astronomical Society*, 448, 2749
- Anders, F., Chiappini, C., Rodrigues, T. S., et al. 2017, *A&A*, 597, A30, doi: [10.1051/0004-6361/201527204](https://doi.org/10.1051/0004-6361/201527204)
- Basu, S., Verner, G. A., Chaplin, W. J., & Elsworth, Y. 2012, *ApJ*, 746, 76
- Bedding, T. R., Huber, D., Stello, D., et al. 2010, *ApJL*, 713, L176
- Bedding, T. R., Mosser, B., Huber, D., et al. 2011, *Nature*, 471, 608
- Campante, T. L., Veras, D., North, T. S. H., et al. 2017, *Monthly Notices of the Royal Astronomical Society*, 469, 1360
- Campante, T. L., Corsaro, E., Lund, M. N., et al. 2019, arXiv e-prints, arXiv:1909.05961. <https://arxiv.org/abs/1909.05961>
- Casagrande, L., Silva Aguirre, V., Schlesinger, K. J., et al. 2016, *Monthly Notices of the Royal Astronomical Society*, 455, 987
- Corsaro, E., & De Ridder, J. 2014, *Astronomy and Astrophysics*, 571, A71
- da Silva, R., Milone, A. d. C., & Rocha-Pinto, H. J. 2015, *Astronomy and Astrophysics*, 580, A24
- Davies, G. R., Silva Aguirre, V., Bedding, T. R., et al. 2016, *Monthly Notices of the Royal Astronomical Society*, 456, 2183
- Flower, P. J. 1996, *ApJ*, 469, 355
- Gaia Collaboration, Katz, D., Antoja, T., et al. 2018, *Astronomy and Astrophysics*, 616, A11
- Gaulme, P., Appourchaux, T., & Boumier, P. 2009, *Astronomy and Astrophysics*, 506, 7
- Handberg, R., & Lund, M. N. 2014, *Monthly Notices of the Royal Astronomical Society*, 445, 2698
- Hekker, S., Broomhall, A.-M., Chaplin, W. J., et al. 2010, *MNRAS*, 402, 2049
- Hekker, S., Elsworth, Y., Mosser, B., et al. 2012, *A&A*, 544, A90
- Høg, E., Fabricius, C., Makarov, V. V., et al. 2000, *Astronomy and Astrophysics*, 355, L27
- Hon, M., Stello, D., & Yu, J. 2018, *Monthly Notices of the Royal Astronomical Society*, 476, 3233
- Houk, N. 1994, *The MK process at 50 years. A powerful tool for astrophysical insight* *Astronomical Society of the Pacific Conference Series*, 60, 285
- Huber, D., Stello, D., Bedding, T. R., et al. 2009, *Communications in Asteroseismology*, 160, 74
- Jenkins, J. M., Twicken, J. D., McCauliff, S., et al. 2016, in *Proc. SPIE*, Vol. 9913, *Software and Cyberinfrastructure for Astronomy IV*, 99133E, doi: [10.1117/12.2233418](https://doi.org/10.1117/12.2233418)
- Jofré, E., Petrucci, R., Saffe, C., et al. 2015, *Astronomy and Astrophysics*, 574, A50
- Jones, M. I., Jenkins, J. S., Rojo, P., & Melo, C. H. F. 2011, *Astronomy and Astrophysics*, 536, A71
- Kallinger, T., Hekker, S., Mosser, B., et al. 2012, *A&A*, 541, 51
- Luck, R. E. 2015, *The Astronomical Journal*, 150, 88
- Lund, M. N., Handberg, R., Davies, G. R., Chaplin, W. J., & Jones, C. D. 2015, *The Astrophysical Journal*, 806, 30
- Mackereth, J. T., Bovy, J., Leung, H. W., et al. 2019, *Monthly Notices of the Royal Astronomical Society*, 489, 176
- Mathur, S., García, R. A., Régulo, C., et al. 2010, *A&A*, 511, 46
- Meléndez, J., Asplund, M., Alves-Brito, A., et al. 2008, *Astronomy and Astrophysics*, 484, L21
- Miglio, A., Montalbán, J., Baudin, F., et al. 2009, *A&A*, 503, L21, doi: [10.1051/0004-6361/200912822](https://doi.org/10.1051/0004-6361/200912822)
- Miglio, A., Chiappini, C., Morel, T., et al. 2013, *Monthly Notices of the Royal Astronomical Society*, 429, 423
- Mints, A., & Hekker, S. 2018, *Astronomy and Astrophysics*, 618, A54
- Mosser, B., Elsworth, Y., Hekker, S., et al. 2011, *Astronomy and Astrophysics*, 537, A30
- Pinsonneault, M. H., Elsworth, Y. P., Tayar, J., et al. 2018, *The Astrophysical Journal Supplement Series*, 239, 32
- Randich, S., Gratton, R., Pallavicini, R., Pasquini, L., & Carretta, E. 1999, *Astronomy and Astrophysics*, 348, 487
- Ricker, G. R., Winn, J. N., Vanderspek, R., et al. 2015, *Journal of Astronomical Telescopes*, 1, 014003
- Rodrigues, T. S., Bossini, D., Miglio, A., et al. 2017, *Monthly Notices of the Royal Astronomical Society*, 467, 1433
- Sharma, S., Stello, D., Bland-Hawthorn, J., et al. 2019, arXiv e-prints, arXiv:1904.12444. <https://arxiv.org/abs/1904.12444>
- Silva Aguirre, V., Davies, G. R., Basu, S., et al. 2015, *Monthly Notices of the Royal Astronomical Society*, 452, 2127
- Silva Aguirre, V., Bojsen-Hansen, M., Slumstrup, D., et al. 2018, *Monthly Notices of the Royal Astronomical Society*, 475, 5487

- Silva Aguirre, V., Christensen-Dalsgaard, J., Cassisi, S., et al. 2019, arXiv e-prints, arXiv:1912.04909.
<https://arxiv.org/abs/1912.04909>
- Stello, D., Meibom, S., Gilliland, R. L., et al. 2011, *ApJ*, 739, 13
- Stello, D., Huber, D., Bedding, T. R., et al. 2013, *ApJ*, 765, L41
- Stello, D., Zinn, J., Elsworth, Y., et al. 2017, *The Astrophysical Journal*, 835, 83
- Van Leeuwen, F. 2007, *A&A*, 474, 653
- Wittenmyer, R. A., Liu, F., Wang, L., et al. 2016, *The Astronomical Journal*, 152, 19
- Yıldız, M., Çelik Orhan, Z., & Kayhan, C. 2019, *Monthly Notices of the Royal Astronomical Society*, 489, 1753
- Yu, J., Huber, D., Bedding, T. R., et al. 2018, *The Astrophysical Journal Supplement Series*, 236, 42
- Zinn, J. C., Stello, D., Huber, D., & Sharma, S. 2019, arXiv e-prints, arXiv:1909.11927.
<https://arxiv.org/abs/1909.11927>

Table 1. Measured and derived stellar properties of our targets. Observed V -magnitudes are extracted from the Tycho-2 catalogue. The global asteroseismic quantities and stellar properties include a statistical and systematic component derived as described in Sections 3 and 4, respectively. We report them here as $value \pm \sigma_{sta} \pm \sigma_{sys}$.

TIC	HIP	ν_{\max}	$\Delta\nu$	V	T_{eff}	[Fe/H]
		(μHz)	(μHz)	(mag)	(K)	(dex)
13097379	114842	$59.10 \pm 1.50 \pm 1.01$	$6.02 \pm 0.03 \pm 0.24$	6.646 ± 0.010	4634 ± 80	0.04 ± 0.08
38574220	19805	$29.40 \pm 0.90 \pm 0.72$	$4.06 \pm 0.20 \pm 0.26$	5.577 ± 0.009	–	–
38828538	21253	$189.90 \pm 1.60 \pm 0.42$	$14.90 \pm 0.10 \pm 0.13$	5.896 ± 0.009	4828 ± 80	0.11 ± 0.08
39082723	4293	$49.30 \pm 2.10 \pm 1.99$	$5.20 \pm 0.10 \pm 0.06$	5.574 ± 0.009	4706 ± 80	-0.05 ± 0.08
47424090	112612	$28.30 \pm 1.80 \pm 1.90$	$3.40 \pm 0.10 \pm 0.29$	6.930 ± 0.010	–	–
70797228	655	$31.80 \pm 1.50 \pm 0.75$	$4.37 \pm 0.20 \pm 0.37$	5.787 ± 0.009	4750 ± 80	0.12 ± 0.08
77116701	103071	$48.30 \pm 7.60 \pm 29.85$	$5.64 \pm 0.20 \pm 3.37$	8.568 ± 0.018	–	–
111750740	113148	$142.60 \pm 2.70 \pm 1.11$	$11.80 \pm 0.10 \pm 0.23$	5.658 ± 0.009	4688 ± 80	0.16 ± 0.08
115011683	103836	$58.80 \pm 1.20 \pm 0.97$	$6.10 \pm 0.10 \pm 0.01$	6.057 ± 0.010	4590 ± 80	-0.13 ± 0.08
129649472	105854	$31.80 \pm 1.20 \pm 1.39$	$4.20 \pm 0.20 \pm 0.11$	5.755 ± 0.009	4748 ± 80	0.28 ± 0.08
139756492	106566	$27.60 \pm 0.90 \pm 0.35$	$4.16 \pm 0.20 \pm 0.65$	6.819 ± 0.010	–	–
141280255	25918	$150.40 \pm 1.00 \pm 0.57$	$12.52 \pm 0.02 \pm 0.10$	5.307 ± 0.009	4630 ± 80	0.33 ± 0.08
144335025	117075	$68.50 \pm 1.60 \pm 0.64$	$7.35 \pm 0.20 \pm 0.39$	6.194 ± 0.010	–	–
149347992	26190	$165.80 \pm 4.10 \pm 16.12$	$11.10 \pm 0.40 \pm 0.60$	6.405 ± 0.010	5132 ± 80	-0.17 ± 0.08
155940286	1766	$73.20 \pm 1.30 \pm 0.32$	$7.40 \pm 0.02 \pm 0.11$	6.810 ± 0.010	4630 ± 80	0.03 ± 0.08
175375523	114775	$60.00 \pm 1.10 \pm 0.30$	$5.80 \pm 0.10 \pm 0.14$	5.899 ± 0.009	4660 ± 80	0.26 ± 0.08
183537408	117659	$57.90 \pm 1.10 \pm 0.80$	$6.20 \pm 0.20 \pm 0.42$	6.781 ± 0.010	–	–
204313960	113801	$106.00 \pm 3.30 \pm 1.47$	$9.40 \pm 0.50 \pm 0.43$	6.083 ± 0.010	4897 ± 80	-0.20 ± 0.08
220517490	12871	$117.30 \pm 1.20 \pm 0.60$	$10.87 \pm 0.02 \pm 0.15$	5.846 ± 0.009	4961 ± 80	-0.26 ± 0.08
237914586	17440	$47.00 \pm 1.40 \pm 1.33$	$5.74 \pm 0.20 \pm 0.08$	3.959 ± 0.009	–	–
270245797	109584	$72.20 \pm 1.70 \pm 0.64$	$6.60 \pm 0.10 \pm 0.27$	6.239 ± 0.009	4824 ± 80	-0.10 ± 0.08
281597433	2789	$73.30 \pm 1.00 \pm 1.37$	$7.20 \pm 0.03 \pm 0.19$	6.163 ± 0.010	4700 ± 80	-0.41 ± 0.08
439399563	343	$44.30 \pm 1.40 \pm 0.66$	$4.54 \pm 0.06 \pm 0.11$	5.892 ± 0.009	4778 ± 80	0.11 ± 0.08
441387330	102014	$46.60 \pm 0.80 \pm 0.86$	$5.25 \pm 0.10 \pm 0.25$	5.592 ± 0.009	4710 ± 80	-0.02 ± 0.08

TIC	R	M	Age	Atmospheric Properties
	(R_{\odot})	(M_{\odot})	(Gyr)	
13097379	$8.49 \pm 0.28 \pm 0.17$	$1.22 \pm 0.08 \pm 0.06$	$6.10 \pm 1.06 \pm 0.97$	Luck (2015)
38574220	–	–	–	–
38828538	$4.66 \pm 0.10 \pm 0.06$	$1.21 \pm 0.05 \pm 0.03$	$6.20 \pm 0.50 \pm 1.02$	Alves et al. (2015)
39082723	$9.30 \pm 0.27 \pm 0.17$	$1.19 \pm 0.09 \pm 0.07$	$5.90 \pm 1.20 \pm 1.37$	Alves et al. (2015)
47424090	–	–	–	–
70797228	$11.27 \pm 0.61 \pm 0.47$	$1.19 \pm 0.13 \pm 0.11$	$6.80 \pm 2.20 \pm 2.20$	Jones et al. (2011)
77116701	–	–	–	–
111750740	$5.11 \pm 0.16 \pm 0.09$	$1.06 \pm 0.07 \pm 0.05$	$10.80 \pm 1.78 \pm 1.71$	Wittenmyer et al. (2016)
115011683	$7.92 \pm 0.21 \pm 0.22$	$1.04 \pm 0.07 \pm 0.07$	$9.90 \pm 1.63 \pm 2.01$	Wittenmyer et al. (2016)
129649472	$10.85 \pm 0.60 \pm 0.24$	$1.13 \pm 0.12 \pm 0.06$	$8.50 \pm 2.76 \pm 1.87$	Jofré et al. (2015)
139756492	–	–	–	–
141280255	$4.98 \pm 0.12 \pm 0.05$	$1.07 \pm 0.06 \pm 0.03$	$11.70 \pm 2.62 \pm 2.39$	Meléndez et al. (2008)
144335025	–	–	–	–
149347992	$7.20 \pm 0.38 \pm 0.20$	$2.17 \pm 0.22 \pm 0.06$	$0.80 \pm 0.30 \pm 0.10$	Randich et al. (1999)
155940286	$6.95 \pm 0.18 \pm 0.14$	$1.01 \pm 0.06 \pm 0.04$	$12.00 \pm 1.78 \pm 36.07$	Wittenmyer et al. (2016)
175375523	$9.00 \pm 0.30 \pm 0.58$	$1.38 \pm 0.09 \pm 0.21$	$4.50 \pm 0.76 \pm 1.35$	Jones et al. (2011)
183537408	–	–	–	–
204313960	$6.50 \pm 0.18 \pm 0.13$	$1.32 \pm 0.07 \pm 0.05$	$3.80 \pm 0.57 \pm 0.57$	Randich et al. (1999)
220517490	$5.61 \pm 0.11 \pm 0.10$	$1.10 \pm 0.04 \pm 0.04$	$6.10 \pm 0.50 \pm 1.18$	Alves et al. (2015)
237914586	–	–	–	–
270245797	$8.73 \pm 0.23 \pm 0.39$	$1.60 \pm 0.09 \pm 0.13$	$2.10 \pm 0.22 \pm 0.64$	Alves et al. (2015)
281597433	$6.77 \pm 0.15 \pm 0.24$	$0.95 \pm 0.05 \pm 0.08$	$11.00 \pm 1.35 \pm 2.73$	Randich et al. (1999)
439399563	$10.69 \pm 0.34 \pm 0.40$	$1.44 \pm 0.11 \pm 0.14$	$3.50 \pm 0.71 \pm 0.78$	da Silva et al. (2015)
441387330	$10.18 \pm 0.34 \pm 0.68$	$1.40 \pm 0.09 \pm 0.20$	$3.40 \pm 0.57 \pm 1.19$	Jones et al. (2011)

NOTE—Last column gives the reference from which we retrieved the central values of T_{eff} and [Fe/H] used for the grid-based modelling. Their uncertainties have been homogenised to 80 K and 0.08 dex, respectively (see Section 4).