

A tidally tilted sectoral dipole pulsation mode in the β Lyrae star TIC 63328020

S. A. Rappaport^{1*}, D. W. Kurtz^{2,3}, G. Handler⁴, D. Jones^{5,6}, L. A. Nelson⁷,
H. Saio⁸, J. Fuller⁹, D. L. Holdsworth³, A. Vanderburg¹⁰, J. Žák^{11,5}, M. Skarka^{11,12},
J. Aiken⁷, P. F. L. Maxted¹³, D. J. Stevens¹⁴, and D. L. Feliz^{15,16}

¹*Department of Physics, and Kavli Institute for Astrophysics and Space Research, M.I.T., Cambridge, MA 02139, USA*

²*Centre for Space Research, Physics Department, North West University, Mahikeng 2745, South Africa*

³*Jeremiah Horrocks Institute, University of Central Lancashire, Preston PR1 2HE, UK*

⁴*Nicolaus Copernicus Astronomical Center, Polish Academy of Sciences, ul. Bartycka 18, 00-716, Warszawa, Poland*

⁵*Instituto de Astrofísica de Canarias, E-38205 La Laguna, Tenerife, Spain*

⁶*Departamento de Astrofísica, Universidad de La Laguna, E-38206 La Laguna, Tenerife, Spain*

⁷*Department of Physics and Astronomy, Bishop's University, 2600 College St., Sherbrooke, QC J1M 1Z7*

⁸*Astronomical Institute, Graduate School of Science, Tohoku University, Sendai 980-8578, Japan*

⁹*Division of Physics, Mathematics and Astronomy, California Institute of Technology, Pasadena, CA 91125, USA*

¹⁰*Department of Astronomy, The University of Texas at Austin, 2515 Speedway, Stop C1400, Austin, TX 78712, USA*

¹¹*Department of Theoretical Physics and Astrophysics, Masaryk University, Kotlářská 2, 60200 Brno, Czech Republic*

¹²*Astronomical Institute, Czech Academy of Sciences, Fričova 298, 25165, Ondřejov, Czech Republic*

¹³*Astrophysics Group, Keele University, Staffordshire, ST5 5BG, UK*

¹⁴*Department of Astronomy & Astrophysics and Center for Exoplanets and Habitable Worlds, PSU, 525 Davey Lab, University Park, PA 16802, USA*

¹⁵*Department of Physics and Astronomy, Vanderbilt University, Nashville, TN 37235, USA*

¹⁶*Department of Physics, Fisk University, 1000 17th Avenue North, Nashville, TN 37208, USA*

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

We report the discovery of the third tidally tilted pulsator, TIC 63328020. Observations with the *TESS* satellite reveal β Lyrae-type eclipses with an orbital period of 1.1057 d, and δ Scuti-type pulsations with a mode frequency of 21.09533 d^{-1} . This pulsation exhibits a septuplet of orbital sidelobes as well as a harmonic quintuplet. Using the oblique pulsator model, the primary oscillation is identified as a sectoral dipole mode with $l = 1, m = |1|$. We find the pulsating star to have $M_1 \simeq 2.5 M_\odot$, $R_1 \simeq 3 R_\odot$, and $T_{\text{eff},1} \simeq 8000 \text{ K}$, while the secondary has $M_2 \simeq 1.1 M_\odot$, $R_2 \simeq 2 R_\odot$, and $T_{\text{eff},2} \simeq 5600 \text{ K}$. Both stars appear to be close to filling their respective Roche lobes. The properties of this binary as well as the tidally tilted pulsations differ from the previous two tidally tilted pulsators, HD74423 and CO Cam, in important ways. We also study the prior history of this system with binary evolution models and conclude that extensive mass transfer has occurred from the current secondary to the primary.

Key words: stars: oscillations – stars: variables – stars: individual (TIC 63328020;)

1 INTRODUCTION

The single-sided or tidally-tilted pulsators are a newly recognized type of pulsating star in close binary systems. In these stars the tidal distortion caused by the companion aligns the pulsation axis of the oscillating star with the tidal axis. This has two important consequences. First, the pulsation axis corotates with the orbit and therefore the stellar pulsation modes are seen at varying aspect, leading to amplitude and phase variations with the orbital phase. Second,

the tidal distortion of the pulsating star causes an intrinsically uneven distribution of pulsation amplitude over the stellar surface.

These two facts can be used to our astrophysical advantage. Viewing the stellar pulsation over the full orbital cycle allows us to constrain the orbital inclination, i , and the obliquity of the pulsation axis, β , to the orbital axis. The required mathematical framework, the oblique pulsator model, has been developed over the past four decades, starting with Kurtz (1982). Even though it was developed for the rapidly oscillating Ap (roAp) stars, whose pulsation axes are tilted with respect to their pulsation axes due to the stellar magnetic fields, it is readily applicable to tidally tilted pulsators. Since

* E-mail: sar@mit.edu

the pulsation axes of those stars are located in the orbital plane, the analyses of the binary-induced variability and of the tidally tilted pulsation mutually constrain each other. The predominant shape of the distorted oscillations, and thus the “pulsational quantum numbers” – the spherical degree l and the azimuthal order m – can be determined from the variation of the pulsation amplitude and phase over the orbit. In reality, the tidal distortion induces coupling between modes of different spherical degree l , modifying the perturbed flux distribution at the surface, and sometimes “tidally trapping” the oscillations on one side of the star. The theoretical groundwork for this type of analysis was laid by Fuller et al. (2020, MNRAS, in press).

Two single-sided pulsators have so far been reported in the literature; both are ellipsoidal variables. The first discovery was HD 74423 (Handler et al. 2020), which contains two chemically peculiar stars of the λ Bootis type in a 1.58-d orbit that are close to filling their Roche lobes. Although the two components are almost identical, only one of them shows δ Scuti pulsation. Rather unusual for this type of pulsating star, there is only a single mode of oscillation, and it is not clear which of the two components is the pulsator. Shortly afterwards, Kurtz et al. (2020) reported the discovery of a second such system, CO Cam. The properties of this binary are different from those of HD 74423. Its orbital period is somewhat shorter (1.27 d), the secondary component is spectroscopically undetected, hence considerably less luminous than the pulsating primary, which is far from filling its Roche lobe. The pulsating star in the CO Cam system is also chemically peculiar, but it is a marginal metallic-lined A-F star (often denoted with the spectral classification “Am:”), and it pulsates in at least four tidally distorted modes.

Both HD 74423 and CO Cam were designated as “single-sided pulsators” because they show enhanced pulsation amplitude on the L_1 side of the star facing the secondary, as explained by Fuller et al. (2020, MNRAS, in press). In the present paper, we report the discovery of the third single-sided pulsator, TIC 63328020, which is different from the two systems studied earlier. With the discovery of a sectoral pulsation mode in TIC 63328020, we now refer to these stars generally as ‘tidally tilted pulsators’, where the more specific name, ‘single-sided pulsators’, can still be used for those stars that have strongly enhanced pulsation amplitude on one side of the star.

In Section 2 we discuss how this object was first noticed in Transiting Exoplanet Survey Satellite (*TESS*) data and we present a detailed analysis of the pulsations. In particular, we show that the pulsations are strongly modulated in amplitude and phase around the orbit, with the peak pulsation amplitudes coinciding in time with the maxima of the ellipsoidal light variations (‘ELVs’), hence in quadrature with the eclipses. In Section 3.1 we present the results of a study of archival data for the spectral energy distribution (‘SED’) as well as the long-term eclipse timing variations (‘ETVs’) for the system. Our radial velocity (RV) data for the system are presented and analysed in Section 3.2, while Section 3.3 utilizes the RV and SED data to analyse the system properties. As a complementary analysis of the system parameters, in Section 3.4 we combine the RV data and the *TESS* light curve via the `phoebe2` code to derive the system parameters. Finally, in Section 4 we use a series of MESA binary evolution grids to understand the formation and evolutionary history of TIC 63328020. We conclude that there was almost certainly a prior history of mass transfer in the system and that the roles of the primary and secondary stars have reversed.

2 ECLIPSES AND PULSATIONS IN TIC 63328020

TIC 63328020 = NSVS 5856840 was reported as a suspected eclipsing binary of the β Lyrae type by Hoffman et al. (2008), the only literature reference to this star. It has an apparent magnitude $V = 11.97$, and a parallax of 0.949 ± 0.035 mas, giving a distance of 1053 ± 38 pc (Luri et al. 2018; Gaia Collaboration et al. 2018) and an absolute visual magnitude of $M_V = 1.86$. The *TESS* Input Catalog (TIC) parameters are $T_{\text{eff}} = 7100$ K and $\log g = 3.6$ (cgs). There is no spectral type. The archival properties of TIC 63328020 are summarised in Section 3.1.1 below.

In addition to the eclipses, δ Scuti-type pulsations were discovered by one of us (DWK) during a visual inspection of *TESS* Sector 15 light curves. During its main two-year mission, 2018 – 2020, *TESS* observed almost the whole sky in a search for transiting extrasolar planets around bright stars ($4 < I_c < 13$) in a wide red-bandpass filter. The measurements were taken in partly overlapping $24 \times 96^\circ$ sectors around the ecliptic poles that were observed for two 13.6-d satellite orbital periods each (Ricker et al. 2015).

TIC 63328020 was observed by *TESS* in Sectors 15 and 16 in 2-min cadence. We used the pre-search data conditioned simple aperture photometry (PDCSAP) data downloaded from the Mikulski Archive for Space Telescopes (MAST)¹. The data have a time span of 51.95 d with a centre point in time of $t_0 = \text{BJD } 2458737.34462^2$, and comprise 34657 data points after some outliers were removed with inspection by eye.

Fig. 1 shows the *TESS* photometry for TIC 63328020. The top panel displays the full Sectors 15 and 16 light curve, where the eclipses and ellipsoidal light variations are obvious. The bottom panel shows a short segment of the light curve where more details can be seen. Importantly, a careful look reveals that the pulsational variations are largest on the ELV humps where the star is brightest. That, of course, is at orbital quadrature, so this is distinctly different from the first two single-sided pulsators, HD 74423 and CO Cam. The frequency analysis below bears out this first impression. First, we look at the light curve, which shows the ellipsoidal orbital variations clearly, and the amplitude modulation of the pulsations on careful inspection.

2.1 The orbital frequency

We analysed the data using the frequency analysis package PERIOD04 (Lenz & Breger 2005)³, a Discrete Fourier Transform program (Kurtz 1985) to produce amplitude spectra, and a combination of linear and nonlinear least-squares fitting to optimise frequency, amplitude and phase. The derived orbital frequency is $\nu_{\text{orb}} = 0.9043640 \pm 0.0000007 \text{ d}^{-1}$ ($P_{\text{orb}} = 1.1057495 \pm 0.0000008 \text{ d}$), where variance from the pulsations and from some low frequency artefacts have been filtered for a better estimate of the uncertainty in the frequency. A 50-harmonic fit by least-squares was done to see how this frequency fits the data, and to show that the pulsation frequencies are not orbital harmonics. In practice, the

¹ http://archive.stsci.edu/tess/all_products.html

² This t_0 was used to begin the analysis, but later changed to the time of pulsation maximum to test the oblique pulsator model. For the assessment of phase errors with nonlinear least-squares fitting, it is important that the t_0 chosen is near to the centre of the data set. Since frequency and phase are degenerately coupled in the fitting of sinusoids, when t_0 is not the centre of the data set, small changes in frequency result in very large changes in phase, since phase is referenced from t_0 .

³ <https://www.univie.ac.at/tops/Period04/>




Figure 1. Top: The full Sectors 15-16 light curve of TIC 63328020 showing the orbital variations. Bottom: A section of the light curve where close inspection shows the pulsations, particularly at orbital quadrature. The zero point of the ordinate scale is the mean.

harmonics at frequencies higher than $20 \times \nu_{\text{orb}}$ do not have statistically significant amplitudes.

After pre-whitening the data by the 50-harmonic fit, some low frequency artefacts from the data reduction were removed with a high-pass filter, which was a simple consecutive pre-whitening of low frequency peaks extracted by Fourier analysis until the noise level was reached in the frequency range $0 - 6 \text{ d}^{-1}$. This was done to study the pulsations with white noise for the purpose of estimating the uncertainties.

2.2 The pulsation

TIC 63328020 pulsates principally in a single mode at a frequency of $\nu_1 = 21.09533 \pm 0.00014 \text{ d}^{-1}$, typical of δ Sct stars. Because this oblique nonradial pulsation mode is observed from changing aspect with the orbit of the star and its synchronous rotation, amplitude and phase modulation of the pulsation generate a frequency septuplet. There is also a harmonic at $2\nu_1$ that generates a quintuplet. These are typical of oblique pulsators. There are also two low-amplitude frequencies at 10.502 d^{-1} and 11.406 d^{-1} that are separated by the orbital frequency. It is likely that one of these is a mode frequency and the other is part of a frequency multiplet from oblique pulsation where the signal-to-noise is too low to detect the other multiplet components. Whichever of these two frequencies is the mode frequency, it is close to, but is not, a sub-harmonic of the principal mode frequency. Because of the low signal-to-noise, we do not discuss these 10.502 d^{-1} and 11.406 d^{-1} frequencies further.

The top panel in Fig. 2 shows the amplitude spectrum for the high-pass filtered data. An inspection shows that there is a septuplet centred on $\nu_1 = 21.09533 \text{ d}^{-1}$ and a quintuplet centred on $2\nu_1 = 42.19065 \text{ d}^{-1}$. The detected frequencies are listed in Table 1. Because the orbital inclination is close to 90° , and the tidal pulsation axis is inclined 90° to that, the pulsation shows amplitude maxima twice per orbit, as will be seen in the next subsection. That then generates two principal peaks in the amplitude




Figure 2. Top: The amplitude spectrum of the high pass data. There is a frequency septuplet centred on $\nu_1 = 21.09533 \text{ c/d}$, and a quintuplet centred on $2\nu_1$. The reason there are two high amplitude first sidelobes about an almost zero amplitude mode pulsation frequency is that we are seeing this pulsation inclined by 90° to the orbital axis, which itself is near to $i = 90^\circ$, thus giving two pulsation amplitude maxima per orbit.

spectrum at $\nu_1 - \nu_{\text{orb}}$ and $\nu_1 + \nu_{\text{orb}}$, along with the other sidelobes. We determined the frequencies in the multiplets by a combination of linear and nonlinear least-squares fitting, and determined that all separations in the multiplets are equal to the orbital frequency $\nu_{\text{orb}} = 0.9043640 \pm 0.0000007 \text{ d}^{-1}$ determined in the last section within 1.5σ . We took the average of those two highest amplitude sidelobe frequencies to determine the value of ν_1 , and generated the frequency multiplets from that. Thus we only give frequency uncertainties on these two highest amplitude frequencies, i.e., those for the first orbital sidelobes $\nu_1 - \nu_{\text{orb}}$ and $\nu_1 + \nu_{\text{orb}}$, which are $\pm 0.00015 \text{ d}^{-1}$ and $\pm 0.00022 \text{ d}^{-1}$, respectively.

We then forced a frequency septuplet for ν_1 and a quintuplet for $2\nu_1$, all split by exactly the orbital frequency, $\nu_{\text{orb}} = 0.904339 \pm 0.000013 \text{ d}^{-1}$. The reason for choosing this exact

Table 1. A least squares fit of the two low frequencies, the frequency septuplet for ν_1 and the frequency quintuplet for $2\nu_1$. The zero point for the phases, $t_0 = \text{BJD } 2458737.14936$, has been chosen to be a time when the two first orbital sidelobes of ν_1 have equal phase.

	frequency d^{-1}	amplitude mmag ± 0.023	phase radians
$\nu_{\text{low}} - \nu_{\text{orb}}$	10.50197	0.133	-0.708 ± 0.175
ν_{low}	11.40633	0.152	1.129 ± 0.152
$\nu_1 - 3\nu_{\text{orb}}$	18.38223	0.181	1.350 ± 0.128
$\nu_1 - 2\nu_{\text{orb}}$	19.28660	0.192	2.859 ± 0.121
$\nu_1 - \nu_{\text{orb}}$	20.19096	1.585	0.719 ± 0.015
ν_1	21.09533	0.110	0.304 ± 0.211
$\nu_1 + \nu_{\text{orb}}$	21.99969	1.080	0.719 ± 0.022
$\nu_1 + 2\nu_{\text{orb}}$	22.90405	0.154	-0.419 ± 0.151
$\nu_1 + 3\nu_{\text{orb}}$	23.80842	0.185	1.296 ± 0.125
$2\nu_1 - 2\nu_{\text{orb}}$	40.38193	0.049	-2.882 ± 0.479
$2\nu_1 - \nu_{\text{orb}}$	41.28629	0.266	1.608 ± 0.087
$2\nu_1$	42.19065	0.341	0.057 ± 0.068
$2\nu_1 + \nu_{\text{orb}}$	43.09502	0.277	-1.652 ± 0.084
$2\nu_1 + 2\nu_{\text{orb}}$	43.99938	0.133	-3.130 ± 0.175

splitting is that for oblique pulsators it is instructive to examine the pulsation phases, and those are inextricably coupled to the frequencies, as can be seen by examining the function we fitted, $(\cos 2\pi f(t - t_0) + \phi)$. Pre-whitening by that solution (and including in the fit the two low-frequencies 10.502 d^{-1} and 11.406 d^{-1}) leads to no variance in the data above noise, as can be seen in the bottom panel of Fig. 2. This shows that exact splitting by the orbital frequency about ν_1 and $2\nu_1$ fits the data, and that there are no other detectable pulsations. This star has one principal oblique pulsating mode.

Finally, we chose a time zero point, t_0 , such that the pulsation phases were equal for the two dominant peaks seen in the top panel of Fig. 2. This effectively chooses a time of pulsation amplitude maximum with the orbit and rotation of the star, thus provides the orbital phase for the time of pulsation maximum. Table 1 shows a least-squares fit of the determined frequencies.

2.3 Pulsation as a function of orbital phase

While the frequencies, amplitudes and phases determined by Fourier analysis and least-squares fitting in the last subsection contain the information to study the oblique pulsation, it is instructive and easier to see how this pulsation varies with orbital aspect by plotting the pulsation amplitude and phase as a function of orbital (rotational) phase. To do this we fitted the pulsation frequency, ν_1 , and its harmonic, $2\nu_1$, to chunks of the data that are $P_{\text{orb}}/10$ in duration. It is immediately obvious from doing this that pulsation amplitude maximum occurs in quadrature to the orbital eclipses. This is the signature of an oblique pulsation in a dipole sectoral mode.

To show this, we have set the time zero point $\frac{1}{4}$ of an orbital period prior to pulsation maximum, as determined from the fitting in the last subsection. That zero point is $t_0 = \text{BJD } 2458736.87292$. We emphasise that this time has been chosen with reference to the pulsation amplitude maximum, hence is independent of the determination of the time of primary eclipse from the study of the orbital variations.




Figure 3. ν_1 : Top and second panels: The pulsation amplitude and phase variation as a function of orbital phase taking ν_1 to be the pulsation frequency. The zero point in time, $t_0 = \text{BJD } 2458736.87292$, has been set to be $\frac{1}{4}$ of an orbital period before the time of pulsation maximum, which was determined by choosing the time when the phases of the first orbital sidelobes are equal. Third panel: The pulsation amplitude as a function of orbital phase for the second harmonic $2\nu_1$. The phase diagram for this frequency is not shown, since it is uninformative because the low amplitude results in high scatter in the phase determinations. The second harmonic peaks when the L_3 point is closest to the observer. Bottom: the orbital light variations as a function of orbital phase for comparison. For this panel the data were binned by a factor of 10. It can be seen that orbital light maximum coincides with pulsation maximum, and that the pulsation amplitude is very similar at orbital phases 0.0 and 0.5. This is currently unique.

Fig. 3 shows the results. The mode is a sectoral dipole mode with $\ell = 1, m = |1|$. Amplitude maximum coincides with orbital quadrature, and the phase reverses by π rad at the times of the eclipses when the line of sight aligns with the tidal axis. This is new and currently unique among tidally tilted pulsators. Because the mode is sectoral, it has a symmetry with respect to the tidal distortion such that the star is not strongly “single-sided”. The third panel of Fig. 3 shows that the harmonic distortion of the mode is strongest during secondary eclipse when the L_3 side of the pulsating star is closest to the observer, thus the L_1 and L_3 sides of the pulsator do differ and the star is mildly a “single-sided pulsator”.

2.4 A identification constraint on the p mode

The standard simple relation for a toy model pulsator of $P\sqrt{\frac{\rho}{\rho_\odot}} = Q$ can be rewritten in terms of observables as

$$\log Q = \log P + \frac{1}{2} \log g + \frac{1}{10} M_{\text{bol}} + \log T_{\text{eff}} - 6.454,$$

where Q is a pulsation “constant” that can be compared with mod-

Table 2. Properties of the TIC 63328020 System

Parameter	Value
RA (J2000) (h m s)	21:20:14.41
Dec (J2000) ($^{\circ}$ ' ")	51:23:41.04
T^a	11.553 ± 0.014
G^b	11.967 ± 0.009
G_{BP}^b	12.267 ± 0.026
G_{RP}^b	11.462 ± 0.024
B^a	12.619 ± 0.228
V^a	11.965 ± 0.114
J^c	10.991 ± 0.020
H^c	10.829 ± 0.019
K^c	10.744 ± 0.018
W1 ^d	10.451 ± 0.024
W2 ^d	10.476 ± 0.022
W3 ^d	10.232 ± 0.070
W4 ^d	> 9.07
$R (R_{\odot})^b$	$3.6^{+0.2}_{-0.6}$
$L (L_{\odot})^b$	13.5 ± 0.8
Orbital Period (d) ^e	1.10575
K_1 (km s ⁻¹) ^e	87.5 ± 4.5
γ (km s ⁻¹) ^e	-8.8 ± 3.6
Distance (pc) ^b	1053 ± 38
μ_{α} (mas yr ⁻¹) ^b	$+2.51 \pm 0.05$
μ_{δ} (mas yr ⁻¹) ^b	$+1.49 \pm 0.05$

Notes. (a) ExoFOP (exofop.ipac.caltech.edu/tess/index.php). (b) Gaia DR2 (Lindegren et al. 2018; Luri et al. 2018; Gaia Collaboration et al. 2018). (c) 2MASS catalog (Skrutskie et al. 2006). (d) WISE point source catalog (Cutri & et al. 2013). (e) This work; see Table 4 for details on the orbital period and Table 5 and Fig. 6 for the RV results.

els, P is the pulsation period in days and $\log g$ is in cgs units. Taking $T_{\text{eff}} = 7100$ K, $\log g = 3.6$ from the TIC, and $M_{\text{bol}} = M_V = 1.86$ from the GAIA parallax and V magnitude then gives for ν_1 a value of $Q = 0.011$, indicative of radial overtone around $n = 6 - 10$. The same calculation for the low frequency peak at 11.406 d^{-1} gives $Q = 0.021$, suggesting a second radial overtone mode.

3 SYSTEM PROPERTIES OF TIC 63328020

3.1 Archival Data

3.1.1 Magnitudes and Gaia Results

We have collected a set of archival magnitudes for TIC 63328020 and report these values in Table 2. Additionally, we list the Gaia information about this object in Table 2. There is a fainter ($G = 19.9$) neighbour star some $2.84''$ away, but there is insufficient Gaia information to tell whether that star is physically associated with TIC 63328020.

3.1.2 Spectral Energy Distribution

The spectral energy distribution (SED) points for this object are plotted in flux units in Fig. 4, and many of them are reported as magnitudes in Table 2. The SED points have all been corrected for interstellar extinction at a level⁴ of $E(g-r) \simeq 0.30 \pm 0.03$, which




Figure 4. SED data and model for TIC 63328020 spanning the blue to 20 micron region for the non-coeval case described in Section 3.3.2. The continuous green, blue and red curves represent the contributions of the secondary, the primary, and the total system flux, respectively.




Figure 5. Long-term eclipse timing variations for TIC 63328020. The data sets used are marked in red labels and include *TESS*, *ASAS-SN*, *WASP*, *KELT*, and *DASCH* (see Table 3 for references). The latter two data sets have each been divided into three subsets of 30 years and ~ 10 months, respectively. The time intervals covered by each data set and the references are given in Table 3. The error bars plotted here and used in the long-term fit are scaled from the actual uncertainties in Table 3, σ_0 , to be $\sigma \equiv 0.1\sqrt{\sigma_0}$. We did this so as not to allow the tiny error bars of the modern-epoch points to totally dominate the determination of the long-term period or its derivative. The green curve is a fit to linear and quadratic terms. See Table 4 for results.

we take to mean $A_V \simeq 1$. We then use the wavelength dependence of extinction given by Cardelli, Clayton & Mathis (1989), in particular, the fitting formulae in their Eqn. (2) and (3) for $\langle A(\lambda)/A(V) \rangle$. Also shown on the figure are fitted curves to the SED based on the contributions from both stars in the binary. These will be discussed in Section 3.3.2.

3.1.3 Archival Photometric Data

In addition to the new *TESS* photometry on TIC 63328020, we have utilized archival photometric data from *ASAS-SN*, *WASP*, *KELT*, and *DASCH* (for references see Table 3) to establish the long-term orbital ephemeris for this binary. The time intervals for the various data sets, and references to the archival data are given in Table 3.

⁴ (<http://argonaut.skymaps.info/query>)

Table 3. Eclipse Timing Variation Data for TIC 63328020

Source	Start Date ^a	End Date ^a	ETV ^b
<i>TESS</i> ^c	58711	58763	+0.0029 ± 0.0003
ASAS-SN ^d	57106	58348	+0.00002 ± 0.0044
KELT 3 ^e	56744	57021	+0.00002 ± 0.0039
KELT 2 ^e	56387	56668	+0.0006 ± 0.0033
KELT 1 ^e	56007	56303	+0.0055 ± 0.0022
WASP ^f	54070	54670	−0.0005 ± 0.0007
DASCH 3 ^g	35119	47857	−0.0303 ± 0.0077
DASCH 2 ^g	22335	35119	−0.0066 ± 0.0050
DASCH 1 ^g	10606	22335	+0.0514 ± 0.0099

Notes. (a) HJD-2 400 000. b. The fold period is 1.105 769 8 days with an epoch of HJD 2 444 000.2795. The ETV value is expressed in days. (c) This work. (d) Shappee et al. (2014) and Kochanek et al. (2017). (e) Pepper et al. (2007) and Pepper et al. (2012). (f) Collier Cameron et al. (2006); Pollacco et al. (2006). (g) Grindlay et al. (2009).

Table 4. Orbital Period Determinations for TIC 633280200

Parameter	Value	Uncertainty
<i>TESS</i> Only		
P_{orb}^a [days]	1.105 749	0.000 001
P_{orb}^b [days]	1.105 9	0.000 2
P_{orb}^c [days]	1.105 751	0.000 001
P_{orb}^d [days]	1.105 754	0.000 001
$P_{\text{orb}}/\dot{P}_{\text{orb}}^e$ [years]	−3050	500
Long-Term ETV Study		
P_{orb}^f [days]	1.105 769 8	0.000 000 3
P_{orb}^g [days]	1.105 769 8	0.000 000 3
$P_{\text{orb}}/\dot{P}_{\text{orb}}^h$ [years]	+15.7 × 10 ⁶	3 × 10 ⁶

Notes. (a) Based on the frequency analysis. (b) Spacing between the ν_1 pulsation sextuplet. The corresponding epoch time is JD 2 458 736.8729. (c) Eclipse timing analysis (‘ETV’) assuming no period changes. The reference fold epoch is JD 2 458 736.8720. (d) ETV analysis allowing for P_{orb} . (e) $P_{\text{orb}}/\dot{P}_{\text{orb}}$ from the ETV analysis. (f) From the long-term photometric ETV analysis assuming P_{orb} is a constant. The fold epoch time is JD 2 444 000.2795. (g) From the long-term photometric ETV analysis allowing for P_{orb} . (h) $P_{\text{orb}}/\dot{P}_{\text{orb}}$ from the long-term photometric ETV analysis

The DASCH data cover more than a century, but only about 1100 scanned plates for this object were available; we divided these up into three roughly 30-yr long intervals which are denoted ‘1’, ‘2’ and ‘3’. The same was done for the KELT data which spanned three observing seasons and was divided into three ~10-month segments. For each data set we derived a time of mid-primary eclipse from a fold of the data about a common reference period and epoch (see Table 3).

From this set of archival photometry we derived a long-term ETV curve for this object which is plotted in Fig. 5. While the long-term orbital period is well defined to a about a part per million with $P_{\text{orb}} = 1.105 769 8(3)$ d, it is also apparent that there are significant non-linear ETVs present. At the moment, there is insufficient information to quantify whether these are due to orbital motion induced by a distance companion or some other effect causing jitter in P_{orb} (e.g., Applegate 1992). In Table 4 we summarize all the information that we have about the orbital period, and its derivative,

Table 5. Radial Velocity Data for TIC 63328020

Epoch (BJD)	RV (Ca II K) ^a km s ^{−1}	RV (CCF) ^b km s ^{−1}	RV (average) ^c km s ^{−1}
2458816.3148	70.9 ± 4.5	67.7 ± 3	68.8 ± 4
2448816.4059	37.1 ± 4.7	30.7 ± 3	32.8 ± 4
2458817.3112	78.0 ± 4.1	77.5 ± 3	77.7 ± 4
2458817.3870	69.0 ± 4.3	70.1 ± 3	69.7 ± 4
2458818.3317	62.8 ± 4.5	67.0 ± 3	65.6 ± 4
2458818.4005	72.3 ± 6.3	77.0 ± 3	75.4 ± 4
2458819.3069	14.1 ± 4.6	12.0 ± 3	12.7 ± 4
2458819.3908	55.4 ± 4.2	58.5 ± 3	57.5 ± 4
Epoch of ϕ_0	K_1^d km s ^{−1}	γ^d km s ^{−1}	cycles after <i>TESS</i>
2458818.7042(36)	87.5 ± 4.5	−8.8 ± 3.6	82.006 ± 0.005

Notes. (a) Based on measurements of the Ca II K line only. (b) Based on a cross correlation analysis of the spectrum over the region of 3700 to 4700 Å. (c) Weighted average of the two RV analyses. (d) Based on the average of the RVs, and a simple sinusoidal fit that does not take into account the Rossiter-McLaughlin effect (but see Section 3.4 for a more complete analysis).

derived in several different ways from the various available data sets.

3.2 Radial Velocity Measurements

TIC 63328020 was observed with the Intermediate Dispersion Spectrograph (IDS) on the 2.5-m Isaac Newton Telescope (INT) between 28 November and 1 December 2019. The blue-sensitive EEV10 detector was used along with a 1'' wide slit and the R1200B grating centred at 4000 Å for an unvignetted spectral coverage of ~3600–4600 Å at a resolution of approximately 4500. In total, eight exposures were acquired with integration times between 1200 s and 1400 s, the dates of which are shown in Table 5. The spectra were wavelength calibrated against arc frames, illuminated using Copper-Argon and Copper-Neon lamps, obtained immediately after each observation in order to avoid shifts due to flexure in the instrument. Bias subtraction, wavelength calibration, sky subtraction and optimal extraction (following the routine of Horne 1986) were performed using standard STARLINK routines.

We extracted the K -velocity of the primary star via two different approaches. In the first we simply fit the deep Ca II K line (at 3934 Å) and thereby estimated RVs with the corresponding uncertainties. In the second, we did a cross-correlation analysis. For the latter, we removed the spectrograph blaze function from the spectra by breaking the spectra into 4-nm wide bins, identifying the highest 10 per cent of flux measurements within each bin (most of which are in the continuum), fitting a basis spline to the continuum points, and dividing the spectrum by the best-fit spline. We then measured radial velocities by cross-correlating each blaze-corrected spectrum with the highest signal-to-noise observation. The results are very similar to those obtained from the Ca II K line alone, but the uncertainties for the cross-correlation result are empirically somewhat smaller. We list both sets of RVs in Table 5.

We do not see any lines from the companion star. We estimate that the luminosity of the secondary is $\lesssim 10$ per cent that of the primary.

The RV data were taken over four consecutive nights, and since the orbital period is 1.1 d, there was insufficient time for the




Figure 6. Radial velocity curve for the primary in TIC 63328020. Because the orbital period is relatively close to a day, the orbital phase sampled over the four nights of observations did not cover the lower half of the RV curve. The red curve is a model fit using *Phoebe2*, yielding 89 ± 6 km/s (see Sect. 3.4).

orbital phases during the observations to drift more than $\sim 40\%$ of an orbital cycle. Nonetheless, we were able to measure the K velocity of the primary star to be 87.5 ± 4.5 km s $^{-1}$ (see Table 5). The radial velocities are plotted in Fig. 6 along with a superposed model fit which is discussed in Section 3.4.

The RV orbital phase zero comes out to be 82.006 ± 0.005 orbital cycles after the *TESS* reference phase zero, and thus is consistent to within an uncertainty of ~ 8 minutes. The RV orbital phase zero is also consistent with the orbital phase zero determined from the pulsation frequencies (see Fig. 3), with the same uncertainty. This shows how precisely the pulsation axis coincides with the tidal axis.

3.3 System Parameters From RV Data Plus SED Fitting

In order to evaluate the binary system parameters of TIC 63328020, we utilised two essentially independent approaches to the analysis (see also (Kurtz et al. 2020)). In the first, we find the stellar masses, inclination and system age that best yield a match to the existing measurements of the spectral energy distribution (SED) and the measured radial velocity of the primary star. In the second approach, we model the *TESS* light curve and simultaneously the radial velocity curve with the *phoebe2* binary light curve emulator (Prša et al. 2016). Both methods utilise a Markov chain Monte Carlo (MCMC) approach to evaluate the uncertainties in the parameters.

3.3.1 Coeval, No-Mass-Loss Assumption

The first method for finding the system parameters utilises three basic ingredients: (1) the known K velocity for the primary star (see Fig. 6); (2) the measured SED points⁵ between 0.4 and 22 μm ; and (3) the Gaia distance (Lindegren et al. 2018).

In this part of the analysis we also make use of the MIST (MESA Isochrones & Stellar Tracks; Dotter 2016; Choi et al. 2016; Paxton et al. 2011; Paxton et al. 2015) evolution tracks for stellar masses between 0.7 and $3.0 M_{\odot}$ with solar composition⁶, in steps of $0.1 M_{\odot}$. Both here and in Section 3.3.2, we utilise the Castelli

⁵ <http://viz-beta.u-strasbg.fr/vizier/sed/doc/>; see also Table 2.

⁶ We have chosen solar metallicity due to lack of information about the interior composition of the primary star in TIC 63328020.

& Kurucz (2003) model stellar atmospheres for $4000 < T_{\text{eff}} < 10,000$ K in steps of 250 K. A solar chemical composition is assumed.

Our approach follows that of Kurtz et al. (2020; and references therein), but we briefly describe our procedure here for completeness. We use an MCMC code (see, e.g., Ford 2005) that evaluates four parameters: the primary mass, M_1 , secondary mass, M_2 , system inclination angle, i , and the MIST equivalent evolutionary phase (EEP) of the primary star. The use of EEPs as a fitted parameter are described in detail in Kurtz et al. (2020).

For each step in the MCMC analysis we use the value of M_1 and the EEP value for the primary to find R_1 and $T_{\text{eff},1}$ from the corresponding MIST tracks, using interpolation for masses between those that are tabulated. That also automatically provides an age, τ , for the star. Since, in this first step of the analysis, we assume that the two stars in the binary are coeval and have experienced no mass exchange, we use the value of τ to find the EEP for the secondary. In turn, that yields the values of R_2 and $T_{\text{eff},2}$.

We check to see that neither star overfills its Roche lobe, and if one does, then that step in the MCMC chain is rejected.

The two masses and the orbital inclination angle determine what the K velocity of the primary should be. This is then compared to the measured value of 89 ± 6 km s $^{-1}$, and determines the contribution to χ^2 due to the RV evaluation.

Finally, we use R_1 and $T_{\text{eff},1}$, as well as R_2 and $T_{\text{eff},2}$, along with interpolated Kurucz model spectra, to fit the 26 available SED points. Here $\log g$ is simply fixed at 4.0. The value of χ^2 for this part of the analysis is added to the contribution from the RV match, and a decision is made in the usual way via the Metropolis-Hastings jump condition (Metropolis et al. 1953; Hastings 1970) as to whether to accept the new step or not.

This is done 10^7 times and the posterior system parameters are collected. The parameter posterior distributions are further weighted according to the derivative of the age with respect to the primary EEP number: $d\tau/d(\text{EEP})$ as described in Kurtz et al. (2020). This corrects for the unevenly spaced EEP points within a larger evolutionary category, and across their boundaries.

The results of this analysis are summarised in a single plot of distributions in Fig. 7. We show the posterior distributions for M_1 , M_2 , R_1 and R_2 , in solar units, while T_1 and T_2 are in units of 10^4 K, and R_1/R_L is dimensionless. For this coeval and no mass exchange scenario, the radius of the lower mass secondary, R_2 is much smaller than for the primary, R_1 . This results from the fact that if the more massive primary is only somewhat evolved off the ZAMS, then the secondary with a much lower mass cannot be hardly evolved at all.

These results are summarised in the second column of Table 6.

3.3.2 Relaxing the No-Mass-Loss Assumption

Here we utilised the same information as in Section 3.3.1 (viz, the K velocity of the primary and the SED points, but we relax the constraint that the two stars must be coeval and have undergone no mass exchange.

The MCMC system parameter evaluation results for this case are summarised by the distributions in Fig. 8. We find three major differences from this removal of the coeval constraint: (1) the distributions are considerably broader than in Fig. 7; (2) the mass of the primary star has shifted considerably to higher values; and (3) the radius of secondary has nearly doubled.

The system parameter results for the case where the no-prior-




Figure 7. Posterior distributions from the MCMC analysis for the case where the stars are assumed to be coeval and without any prior mass exchange. For M_1 , M_2 , R_1 , and R_2 , the x axis is labeled in solar units, while T_1 and T_2 are in units of 10^4 K, and R_1/R_L is dimensionless. Note, in particular, that R_2 is small compared to R_1 .




Figure 8. Posterior distributions from the MCMC analysis for the case where the stars are *not* assumed to be coeval and without any prior mass exchange. The axis labeling is the same as in Fig. 8. Note, that the distributions are broader than in Fig. 7, and M_1 has gotten larger, as has R_2 .

mass-exchange assumption has been relaxed are summarised in the third column of Table 6.

3.4 System Parameters From RV Data Plus Light curve Modelling

In Section 3.3.1 above we analysed the basic system parameters from an MCMC evaluation of the two masses, the inclination angle, and the evolutionary phases (EEP) of the two stars. The fitted quantities were K_1 and 26 SED points, coupled with the Gaia distance. In Section 3.3.2 we relaxed the coeval constraint on the two stars and fit independently for their masses and radii.

We now proceed to analyse the system parameters via simultaneous fitting of the *TESS* orbital light curve as well as the radial velocity curve using the next-generation Wilson-Devinney code *phoebe2* (Prša et al. 2016). First, we removed the pulsations from the light curve. In addition, a visual inspection of the orbital light curve in Fig. 1 shows that the ELV peak following the primary eclipses is lower than the preceding ELV peak. This differ-




Figure 9. The model *phoebe2* light curve (red) on the observations (black). A sinusoidal component of amplitude 0.4 per cent was removed from the *TESS* light curve to equalize the two ELV maxima before performing the fit (see text for details).

ence can be empirically removed by subtracting a simple sinusoid at the orbital frequency and of amplitude 4110 ppm. The phasing of this sinusoid would be correct for the Doppler boosting (DB) effect (Loeb & Gaudi 2003; van Kerkwijk et al. 2010) if the lower luminosity secondary were the source of the DB. From the orbital solutions already in hand (see columns 2 and 3 of Table 6), the expected DB amplitude would be ~ 1000 ppm and $\lesssim 300$ ppm for the primary and secondary, respectively. Thus, the observed amplitude is far too large to be the DB effect, and we attribute it to spots on the secondary that are corotating with the orbit. Therefore, we elected to subtract off a sinusoidal component with amplitude 4110 ppm from the light curve before carrying out the fitting with *phoebe2*.

The light-curve fitting utilized the MCMC methodology outlined in Boffin et al. (2018) and Jones et al. (2019). The component masses, radii and temperatures, and the orbital inclination were allowed to vary freely over ranges consistent with the observed SED. The only additional free parameters were the gravity brightening exponent, β_1 (where $T_{\text{eff,local}}^4 = T_{\text{eff,pole}}^4 (g_{\text{local}}/g_{\text{pole}})^\beta$), of the primary and the (Bond) bolometric albedo (Horvat et al. 2019), A_2 , of the secondary, which are critical for constraining the ELV and irradiation effect amplitudes, respectively.

The best-fitting *phoebe2* orbital light curve is shown in Fig. 9 while the corresponding model fit to the radial velocities was presented in Fig. 6. The resultant model parameters for the system are listed in the last column of Table 6.

It is clear that the *phoebe2* model provides a remarkably good fit to both the observed light and radial velocity curves, with all model variables extremely well constrained. The model variables all present with strongly Gaussian posteriors, however several are strongly correlated. For example, due to the use of a single photometric band, the posteriors of the primary and secondary temperatures show a weak positive correlation. Likewise, the primary’s radius is positively correlated with its mass, with larger primary masses necessitating larger primary radii in order to maintain the same Roche lobe filling factor and thus the same amplitude of ELV. Ultimately, further observations are required to break these correlations but, nonetheless, the current data are sufficient to strongly constrain the properties of the system (see Kurtz et al. 2020, for further discussion of the fitting of a singled-band ELV light curve for a similar case, albeit without the observed eclipses of TIC 63328020 which provide additional strong constraints).

The results for the system parameters derived from the

Table 6. Derived Parameters for the TIC 63328020 System

Input Constraints	SED + RV ^a	SED + RV ^b	Light curve + RV ^c
Period (days)	1.1057	1.1057	1.1057
K_1 (km s ⁻¹) ^d	89 ± 6	89 ± 6	89 ± 6
Spectral	26 SED points ^e	26 SED points ^e	...
Light curve modeling	TESS ^g
Distance (pc)	1054 ± 20	1054 ± 20	1054 ± 20
A_V	1.0	1.0	1.0
Derived Parameter	SED + RV ^a	SED + RV ^b	Light curve + RV ^c
M_1 (M_\odot)	2.08 ± 0.02	2.81 ± 0.36	2.50 ± 0.20
M_2 (M_\odot)	1.11 ± 0.06	1.11 ± 0.10	1.07 ± 0.06
R_1 (R_\odot)	2.83 ± 0.04	2.87 ± 0.12	3.10 ± 0.10
R_2 (R_\odot)	1.02 ± 0.08	1.97 ± 0.15	2.06 ± 0.06
$T_{\text{eff},1}$ (K)	8025 ± 80	7885 ± 120	8200 ± 450
$T_{\text{eff},2}$ (K)	6111 ± 200	4915 ± 400	5600 ± 250
i (deg)	73 ± 6	76 ± 4	79.1 ± 0.6
a (R_\odot)	6.6 ± 0.1	7.0 ± 0.4	6.9 ± 0.2
R_1/R_L	≥ 0.98	0.88 ± 0.06	≥ 0.92
K_2 (km s ⁻¹) ^h	189 ± 8	225 ± 17	208 ± 10
age (Myr)	800 ± 25
β_1 ⁱ	0.80 ± 0.06
A_2 ^j	0.68 ± 0.06

Notes. (a) MCMC fits to the measured RV amplitude plus the SED points. The assumption is made that the two stars are coeval in their evolution, and have not exchanged any mass. (b) Same as (a) except that the assumption of no prior mass exchange has been dropped. (c) `phoebe2` fit to the *TESS* orbital light curve plus the RV amplitude. (d) This work (see Sect. 3.4). (e) See Fig. 4. (f) Modelled with `phoebe2`. (g) Gaia DR2 (Lindgren et al. 2018). (h) Predicted from the MCMC parameter evaluations. (i) Gravity brightening exponent. (j) Bond bolometric albedo of the secondary.

`Phoebe` fit to the *TESS* are summarised in the fourth column of Table 6.

4 FORMATION AND EVOLUTION – PRIOR HISTORY OF MASS TRANSFER

In order to determine the formation history of TIC 63328020, we must first consider whether or not mass transfer between the binary components has had a significant effect on its evolution. The second column of Table 6 lists the inferred properties of TIC 63328020 that were derived based on the SED and RV data under the assumption of no mass transfer during the binary’s evolution. With this latter constraint relaxed, the properties of the binary deduced using (i) the SED and RV data, and (ii) the RV data in conjunction with a `phoebe2` fit to the light curve, are shown in the third and fourth columns, of Table 6, respectively.

The most glaring discrepancy between any of the predicted properties of the two stars occurs for the radius of the secondary (R_2). The inference for the radius made under the assumption of no mass transfer disagrees with the other two inferences by nearly a factor of two. Given that the last two inferences were derived independently (although they do share the same RV data) and given that no constraint on mass transfer was imposed, the relatively good agreement between these two cases seems to imply that TIC 63328020 very likely experienced mass transfer in the past. Moreover, an orbital period on the order of days is typical of many ‘Algol-like’ binaries for which mass transfer/loss occurred during their prior evolution (see, e.g., Batten 1989; Eggleton 2000).

In analysing the evolution of the progenitor binary we will therefore assume that mass transfer occurred. The next question to address is whether or not the current primary was the original

primary of the progenitor system (i.e., the more massive one) or whether a mass-ratio reversal occurred (i.e., an Algol-like evolution). If a mass-ratio reversal did not occur, then we are forced to conclude that the original primary could not have lost much mass simply because the original mass ratio (M_1/M_2) would have been so large ($\gtrsim 3$) that the binary would have undergone a dynamical instability leading to the presumed merger of the two stars (see, e.g., Webbink 1976; Soberman, Phinney & van den Heuvel 1997 for a discussion of the conditions leading to dynamical instability).

For the other scenario, the masses of the two primordial components (we will refer to them as ‘Star 1’ and ‘Star 2’) may have changed significantly during the course of the evolution. The largest uncertainty concerns the degree to which mass transfer between the components is conservative. For a fully conservative transfer, all of the mass that is lost by the more massive star (Star 1) is subsequently accreted by Star 2. On the other hand, for completely non-conservative mass transfer, the mass of Star 2 would not change as the mass of Star 1 decreased. Because we do not know how non-conservative mass transfer could have been (or the amount of angular momentum transported out of the binary), we have investigated a realistic range of possibilities.

4.1 Evolutionary Grid

To determine the properties of putative primordial binaries that could evolve to approximately match the observational properties inferred for TIC 63328020, we generated a grid of binary evolutions using version 10108 of `MESA`. Specifically, we use the binary version of `MESA` for which the evolution of both the donor and accretor stars are computed simultaneously (see Paxton et al. 2011; Paxton et al. 2015). Because of the large number of dimensions of initial parameter space, we have tried to minimize computations by




Figure 10. Representative tracks of three distinctly different primordial binary evolutions leading to the formation of TIC 63328020 are shown in the HR diagram. The evolution of the more massive component of the primordial binary (Star 1) is denoted by a solid curve and the evolution of its less massive companion (Star 2) is represented by a dashed curve. Arrows superposed on the curves denote the direction of increasing age (i.e., the direction in which the component is evolving). Different colours are used to denote each pair of the components of the primordial binary. Each component initially starts evolving from the ZAMS and those masses for all three cases are labelled. The red tracks correspond to a binary that experiences highly non-conservative mass-transfer ($\beta = 0.8$), while the blue and green tracks assume fully conservative mass-transfer. The solid dots of the same colour as the track indicate the points in the evolution when the orbital period matches the observed value (1.1057 d). The purple dots indicate the points for which (rapid) thermal-timescale mass-transfer is first initiated. All of the tracks terminate when both stars have simultaneously fill their Roche lobes.

fixing the chemical composition. Given the binary’s proximity to the mid-plane of the Galaxy, we have set the metallicity to a value of $Z = 0.03$ and have taken the hydrogen and helium abundances to be $X = 0.69$ and $Y = 0.28$, respectively.

This still leaves the initial mass values of the more massive component (Star 1) and its companion (Star 2), in addition to the primordial orbital period (P_{orb}), as unknown values. Moreover, the degree to which mass transfer is non-conservative and the mechanism describing the removal of orbital angular momentum is very uncertain. This uncertainty can be parametrized in terms of the quantities α and β (for details see Tauris & van den Heuvel 2006). In the MESA code, α is the fraction of the mass lost by the donor star that gets ejected from the binary such that the mass carries away the specific angular momentum of the donor star (i.e., fast Jeans’ ejection), and β is the fraction of the mass that is ejected from the accretor and is assumed to carry away the specific angular momentum corresponding to that star. Thus the mass gained by the accretor (secondary) can be written as:

$$\delta M_2 = -(1 - \alpha - \beta)\delta M_1. \quad (1)$$

We further assume that none of the mass that is lost from the binary forms a circumbinary torus that can extract additional orbital angular momentum during the binary’s evolution.

After extensive experimentation we found that the choices of the primordial masses and orbital period in addition to the sum of $(\alpha + \beta)$ were much more important factors than the individual choices of the values of α and β in trying to replicate the properties of TIC 63328020. For this reason our final set of models have been computed with $\alpha = 0$. Finally, orbital angular momentum dissipation was calculated based on the torques associated

with gravitational radiation and magnetic braking as described in Goliaš & Nelson (2015) and Kalomeni et al. (2016), with the magnetic braking index set equal to 3. The magnetic braking formula (Verbunt-Zwaan law; Verbunt & Zwaan 1981) was inferred from observations of low-mass main-sequence stars and thus must sometimes be extrapolated to stars that are either evolved, very low-mass, or rapidly rotating. Although the magnitude of magnetic braking torques remains uncertain, it has relatively little effect on the evolutionary tracks until after thermal timescale mass transfer has occurred.

To further optimise numerical computations, we were guided by a number of grids that were previously generated to solve for the evolution of other types of interacting binaries. Specifically, we have used the grids generated for post-Algol binaries such as MWC 882 (Zhou et al. 2018) and wide, hot subdwarf binaries (Nelson & Senhadji 2019) to try to constrain the range of possible initial conditions. After creating new grids of binary evolution tracks, we found that both conservative and non-conservative evolutions could produce the desired results given the appropriate choices of the primordial masses and the primordial period. Thus we conclude that a fine-tuning of the initial conditions is not required in order to reproduce the observations.

Possible evolutionary scenarios can be divided into two separate classes: (1) the more massive star (Star 1) loses a relatively small fraction of its initial mass while the companion (Star 2) gains some portion of that mass; or, (2) the more massive primordial star loses a large fraction of its mass leading to a mass-ratio reversal (the mass ratio being defined as $q = M_{\text{Star1}}/M_{\text{Star2}}$), thus implying that the accretor becomes more massive than the donor. For

either scenario, the evolution can be fully conservative ($\beta = 0$) or highly non-conservative ($\beta = 0.8$).

We find that the first scenario (mass ratio does not change significantly) never fully reproduces the observationally inferred properties listed in columns 2 and 3 of Table 6. Although this grouping (class) of evolutionary tracks can reproduce most of the properties of TIC 63328020, we did not find a primordial binary that could ultimately produce a secondary star with such a large radius ($\approx 2R_{\odot}$) while simultaneously matching all of the other inferred properties of both stars. The problem stems from the following physical constraints: (i) in order to bloat the accretor sufficiently, mass-transfer rates in excess of $\sim 3 \times 10^{-7} M_{\odot} \text{ yr}^{-1}$ are required for extended periods of time; and, (ii) binaries with large mass ratios tend to experience dynamical instabilities when Roche lobe overflow first commences. With respect to the latter issue, the very large mass ratio inferred for TIC 63328020 necessarily implies that the accretor could only have gained a few tenths of a solar mass during the evolution (otherwise the initial evolution would have been dynamically unstable). And given the required high mass-transfer rates and the small net accretion, this implies that mass transfer would have occurred over an extremely short interval ($\lesssim 1$ Myr), making the whole scenario less likely. Moreover, mass-transfer rates of $\sim 10^{-6} M_{\odot} \text{ yr}^{-1}$ are expected at the current epoch and there is little evidence to support such a high value (see the discussion below).

According to the second scenario, the original primary of the primordial binary (i.e., the donor) loses so much mass to its accreting companion that a mass-ratio reversal occurs (in other words, the observed low mass secondary of TIC 63328020 was originally the higher mass star). As discussed above, the largest uncertainty concerns the choice of β and we attempt to mitigate the effects of this uncertainty by creating a grid of models with the variable β taken to be one of the dimensions of parameter space. It is generally expected that the evolution of Algol-like binaries will be at least mildly non-conservative (see, e.g., Eggleton 2000) and that is why we chose to investigate the range $0 \leq \beta \leq 0.8$.

The evolution of both binary stars in the Hertzsprung-Russell diagram for three representative systems (one for the first scenario and two for the second scenario corresponding to extreme values of β) is shown in Fig. 10. For the first scenario (see the solid and dashed blue curves for the evolution of the donor and accretor, respectively), we chose a primordial binary consisting of 2.7 and $0.9 M_{\odot}$ components with an orbital period of 3.0 d. Possible solutions for TIC 63328020 are denoted by the solid blue dots. For the second scenario, this fully conservative case has components initially consisting of 2.25 and $1.85 M_{\odot}$ stars in a 2.35 -d orbit (see the green solid and dashed curves, respectively). The highly non-conservative evolution with $\beta = 0.8$ is denoted by the red curves.

The primordial binary consisted of a $3.5 M_{\odot}$ donor in a 1.706 -d orbit with a $2.09 M_{\odot}$ accretor. For these latter two sets of evolutionary tracks there are two sets of solid dots (green and red) that denote possible solutions at the observed orbital period of 1.1057 d. In each case, the latter set of dots (corresponding to a later age) better fits the inferred properties of TIC 63328020 enumerated in Table 6. The solid magenta dots indicate the onset of (rapid) thermal timescale mass transfer. For all cases, the evolutionary tracks are seen to abruptly change their trajectories in the HR diagram once mass transfer commences. The donor stars all tend to evolve towards lower luminosities and effective temperatures while the accretors immediately evolve towards higher temperatures and luminosities.

Each of the three tracks terminates once the accretor has ex-




Figure 11. The evolution of the masses of the two components of the primordial binary as a function of orbital period is illustrated. Arrows superposed on the curves denote the direction of increasing age (i.e., the direction in which the component is evolving). Star 1 is always losing mass while Star 2 is accreting mass. The colour scheme for the three cases shown in Fig. 10 is repeated here. The initial orbital period for each case is also labeled (upper left) and the observed orbital period (1.1057 d) is denoted by a vertical black bar. The solid dots of the same colour as the curves indicate the points that could be possible solutions for the properties of TIC 63328020. The cyan-shaded region represents the approximate range of possible masses of the current primary based on the analysis presented in Table 6 (columns 3 and 4). The yellow-shaded region corresponds to the possible range of masses for the current secondary. All of the curves terminate when both stars have simultaneously filled their Roche lobes.

panded sufficiently to fill its Roche lobe. A summary of the three tracks and the best fit to the inferred data enumerated in columns 3 and 4 of Table 6 is presented in Table 7. Note that the subscripts 1 and 2 denote the properties of the primary and secondary, respectively, for TIC 63328020 at the current epoch. The age is measured from the formation of the primordial binary and $\log \dot{M}$ indicates the (current) mass-transfer rate from the donor star.

In order to further elucidate the scenarios associated with our three representative tracks, the evolution of the masses of each component is shown as a function of the orbital period (P_{orb}) in Fig. 11. The colour coding and the use of solid and dashed lines in addition to the solid dots have the same meaning as that described for Fig. 10. For track #1 (blue curves), the primordial donor simply loses a few tenths of a solar mass that is then gained by the primordial secondary (accretor). Although this is one of the simplest types of evolution that could reproduce the observed masses of TIC 63328020, we were unable to find any combination of initial conditions or values of the parameters governing some of the input physics (e.g., orbital angular momentum dissipation) that reproduced the inferred radius of the secondary. Instead, both tracks #2 (green curves) and #3 (red curves) can produce reasonable facsimiles of TIC 63328020.

Both donor stars (solid curves) initially undergo thermal

Table 7. MESA Model Parameters for the TIC 63328020 System

Model Parameter	Track #1 (Blue)	Track #2 (Green)	Track #3 (Red)
$M_{\text{Star}1,0}$ (M_{\odot}) ^a	2.70	2.25	3.50
$M_{\text{Star}2,0}$ (M_{\odot}) ^a	0.90	1.85	2.09
$P_{\text{orb},0}$ (d) ^a	3.00	2.35	1.71
β ^b	0.0	0.0	0.8
q_0 ($\equiv M_{1,0}/M_{2,0}$) ^c	3.00	1.216	1.675
M_1 (M_{\odot}) ^d	2.48	2.87	2.55
M_2 (M_{\odot}) ^d	1.12	1.23	1.18
R_1 (R_{\odot}) ^d	3.10	3.07	3.06
R_2 (R_{\odot}) ^d	0.92	2.22	2.19
$T_{\text{eff},1}$ (K) ^d	6280	9490	8510
$T_{\text{eff},2}$ (K) ^d	5050	5410	5630
$L_{\text{bol},1}$ (L_{\odot}) ^d	13.4	68.7	44.2
$L_{\text{bol},2}$ (L_{\odot}) ^d	0.49	3.80	4.33
$R_1/R_{L,1}$ ^d	1.0	0.94	0.98
$R_2/R_{L,2}$ ^d	0.43	1.0	1.0
$\log \dot{M}$ ($M_{\odot} \text{ yr}^{-1}$) ^e	-5.57	-8.69	-9.00
$\dot{P}_{\text{orb}}/P_{\text{orb}}$ (yr^{-1})	-4.0E-6	+5.9E-10	-6.1E-10
system age (Myr)	485	935	545

Notes. (a) ‘Star 1’ and ‘Star 2’ refer to the original primordial primary and secondary, respectively. The subscript ‘0’ indicates the initial system parameters. (b) β is the fraction of mass that is transferred to the accretor but is ejected from the system with the specific angular momentum of that star. The parameter α (not in the Table) is fixed at 0.0 and is the fraction of mass lost by the donor star that is ejected from the system with the specific angular momentum of the donor star. (c) Initial mass ratio of the primordial binary ($M_{\text{Star},1}/M_{\text{Star},2}$). (d) These are the model parameters for the current-epoch TIC 63328020 system. (e) The total rate of mass lost by the donor star.

timescale mass transfer causing the orbital period to decrease (see Fig. 11) as both donor stars shrink due to quasi-adiabatic mass loss. However, once the mass ratio has been reduced to about unity, further mass transfer causes the orbits to expand with a concomitant increase in P_{orb} . As the donor stars approach quasi-thermal equilibrium (with a much reduced mass transfer rate), they contract forcing the orbit to shrink⁷. The accretors for both cases continuously gain mass with a resulting increase in their radii. The tracks terminate once the accretors also fill their Roche lobes.

4.2 Discussion

Based on the analysis of an extensive grid of models, we conclude that there is a wide range of initial conditions that can replicate the currently observed properties of TIC 63328020. The simplest type of evolution wherein the primordial primary loses a few tenths of a solar mass to a much less massive accretor, although appealing, cannot reproduce all of the inferred properties. But a wide range of evolutionary scenarios for which a mass-ratio reversal occurs (i.e., the primordial primary becomes the less massive secondary) can be accommodated. In particular, if the evolution is highly non-conservative, then the total mass of the primordial binary would have to be considerably more massive than the presently inferred value with an initial mass ratio of $q_0 \gtrsim 1.5$. For more conservative

evolutions, the initial total mass can be much smaller and the mass ratio much closer to unity. Thus there is a wide range of initial parameters for the primordial binary that can produce robust models of TIC 63328020.

Although there are significant uncertainties associated with systemic mass loss and the orbital angular momentum dissipated as a result of this non-conservative mass transfer, we find that the individual choices of the parameters α and β are not nearly as important as the contribution from their sum. For this reason, we parametrized the effects of non-conservative mass transfer in terms of β ($\alpha = 0$). We conclude that the values of β in the range of 0 to 0.8 can lead to plausible solutions for the properties of TIC 63328020 (see columns 3 and 4 of Table 6). However, the highly conservative models tend to produce primaries with higher effective temperatures ($\gtrsim 1000$ K higher). For this reason, we somewhat prefer models for which $\beta \simeq 0.3$.

Another important result to note is that the ‘simple’ evolutionary scenario implies that mass-transfer rates at the present epoch should be on the order of $10^{-6} M_{\odot} \text{ yr}^{-1}$. By way of contrast, the ‘mass-reversal’ scenario requires mass-transfer rates that are typically three orders of magnitude smaller (Table 7). We examined the spectrum of TIC 63328020 for P Cygni profiles and found no evidence for that feature. This would seem to imply a relatively low mass transfer rate. We also examined four WISE band observations looking for any evidence of nebulosity that might be expected due to a significant wind emanating from the binary. We could not find any hint of nebulosity in that region. There was also no sign of any NIR nebulosity from the PanSTARRS images. Although not conclusive, these results seem to hint at a relatively low mass transfer rate or one that is not highly non-conservative.

One of the very intriguing features of many of our evolutionary tracks that reproduce robust models of TIC 63328020 is that the accretor is very close to filling its Roche Lobe. Since it is relatively unlikely that we would find such a configuration based solely on the observed pulsational properties, the question arises as to whether the binary had already evolved to a point where both stars temporarily over/filled their Roche lobes before one of them contracted leading to the currently observed configuration. If one of the stars contracted then it is possible that it could remain in a detached state for at least a Kelvin time. On the other hand, it is quite possible that the binary would have merged were contact to have occurred. We are currently trying to address this and questions related to the formation and evolution of WU Ma binaries using smoothed particle hydrodynamics (SPH; S. Tripathi, L. Nelson, & T. S. Tricco 2020 [in preparation]).

5 CONCLUSIONS

In this work we report the discovery of a short-period β Lyrae binary with tidally-tilted pulsations at $\nu = 21.09533 \text{ d}^{-1}$. The pulsation amplitude varies with orbital phase and is a maximum at orbital quadrature, i.e., when the ellipsoidal light variations are at a maximum. The phase of the pulsations rapidly change by more than π radians around the time of the primary eclipse, and there is a smaller jump in phase at the secondary eclipse by about half that amount in the opposite direction. We note that the phase is not a pure π -radian jump because the mode is distorted from a pure sectoral dipole mode.

In order to help visualize how the tidally tilted pulsations would appear to an observer on an circumbinary planet orbiting TIC 63328020, we include a simulation in the form of an MP4

⁷ Note that the green track of the donor star subsequently experiences a second orbital period minimum. This is due to the thermal re-adjustment of the core from purely convective energy transport to a fully radiative mode.

video (‘TIC63328020.mp4’). The video is supplied as Supporting Information for the paper. This same video is also presented in Fuller et al. (2020, MNRAS, in press).

The pulsating star has $M_1 \simeq 2.5 M_\odot$, $R_1 \simeq 3 R_\odot$, and $T_{\text{eff},1} \simeq 8000$ K, while the secondary has $M_2 \simeq 1.1 M_\odot$, $R_2 \simeq 2 R_\odot$, and $T_{\text{eff},2} \simeq 5600$ K. Both stars appear to be close to filling their respective Roche lobes. The orbital period is constant to a part in $\sim 10^5$ over the last century. However, the period appears to vary erratically on timescales of weeks to decades. At present we have no firm explanation for this behavior.

We have carried out an investigation of the history of this system with an extensive set of binary evolution models. We conclude that the most likely scenario is that there has been a prior epoch of mass transfer which has reduced the mass of the original primary so that it is currently the low-mass secondary. By contrast, the original secondary is now the pulsating primary star. The mass transfer may still be ongoing with a low mass-transfer rate of $\sim 3 \times 10^{-9} M_\odot \text{ yr}^{-1}$.

Although the architecture and evolutionary histories of the three known tidally tilted pulsators (HD 74423, CO Cam, and TIC 63328020) are unique, they all feature heavily tidally distorted δ Sct pulsators in short-period orbits. Whereas HD 74423 and CO Cam feature axisymmetric tidally tilted pulsations trapped on the L_1 side of the star, the tidally tilted mode in TIC 63328020 exhibits very different phase and amplitude modulation, indicative of a non-axisymmetric ($|m| = 1$) mode that is mostly confined to the equator (Fuller et al. 2020). Future discoveries of tidally tilted pulsators will likely reveal more diversity amongst this new class of stars.

ACKNOWLEDGEMENTS

Based on observations made with the Isaac Newton Telescope operated by the Isaac Newton Group of Telescopes, which resides on the island of La Palma at the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias. The authors thankfully acknowledge the technical expertise and assistance provided by the Spanish Supercomputing Network (Red Española de Supercomputación), as well as the computer resources used: the LaPalma Supercomputer, located at the Instituto de Astrofísica de Canarias.

G.H. acknowledges financial support from the Polish National Science Center (NCN), grant no. 2015/18/A/ST9/00578. D.J. acknowledges support from the State Research Agency (AEI) of the Spanish Ministry of Science, Innovation and Universities (MCIU) and the European Regional Development Fund (FEDER) under grant AYA2017-83383-P. DJ also acknowledges support under grant P/308614 financed by funds transferred from the Spanish Ministry of Science, Innovation and Universities, charged to the General State Budgets and with funds transferred from the General Budgets of the Autonomous Community of the Canary Islands by the Ministry of Economy, Industry, Trade and Knowledge. This research was supported by the Erasmus+ programme of the European Union under grant number 2017-1-CZ01-KA203-035562. L.N. thanks the Natural Sciences and Engineering Research Council (Canada) for financial support through the Discovery Grants program. Some computations were carried out on the supercomputers managed by Calcul Québec and Compute Canada. The operation of these supercomputers is funded by the Canada Foundation for Innovation (CFI), NanoQuébec, Réseau de Médecine Génétique Appliquée, and the Fonds de recherche du Québec – Nature et tech-

nologies (FRQNT). J.A. thanks NSERC (Canada) for an Undergraduate Student Research Award (USRA). D.J.S. acknowledges funding support from the Eberly Research Fellowship from The Pennsylvania State University Eberly College of Science. The Center for Exoplanets and Habitable Worlds is supported by the Pennsylvania State University, the Eberly College of Science, and the Pennsylvania Space Grant Consortium.

This project utilized data from the Digital Access to a Sky Century@Harvard (‘DASCH’) project at Harvard that is partially supported from NSF grants AST-0407380, AST-0909073, and AST-1313370. This paper also makes use of the WASP data set as provided by the WASP consortium and services at the NASA Exoplanet Archive, which is operated by the California Institute of Technology, under contract with the National Aeronautics and Space Administration under the Exoplanet Exploration Program (DOI 10.26133/NEA9).

This project also makes use of data from the Kilodegree Extremely Little Telescope (‘KELT’) survey, including support from The Ohio State University, Vanderbilt University, and Lehigh University, along with the KELT follow-up collaboration.

REFERENCES

- Applegate J. H., 1992, ApJ, 385, 621
 Batten A. H., 1989, Space Sci. Rev., 50
 Boffin H. M. J. et al., 2018, A&A, 619, A84
 Cardelli J. A., Clayton G. C., Mathis J. S., 1989, ApJ, 345, 245
 Castelli F., Kurucz R. L., 2003, in IAU Symposium, Vol. 210, Modelling of Stellar Atmospheres, Piskunov N., Weiss W. W., Gray D. F., eds., p. A20
 Choi J., Dotter A., Conroy C., Cantiello M., Paxton B., Johnson B. D., 2016, ApJ, 823, 102
 Collier Cameron A. et al., 2006, MNRAS, 373, 799
 Cutri R. M., et al., 2013, VizieR Online Data Catalog, II/328
 Dotter A., 2016, ApJS, 222, 8
 Eggleton P. P., 2000, NewAR, 44, 111
 Ford E. B., 2005, AJ, 129, 1706
 Gaia Collaboration et al., 2018, A&A, 616, A1
 Goliaš J., Nelson L., 2015, ApJ, 809, 80
 Grindlay J., Tang S., Simcoe R., Laycock S., Los E., Mink D., Doane A., Champine G., 2009, Astronomical Society of the Pacific Conference Series, Vol. 410, DASCH to Measure (and preserve) the Harvard Plates: Opening the 100-year Time Domain Astronomy Window, Osborn W., Robbins L., eds., p. 101
 Handler G. et al., 2020, Nature Astronomy
 Hastings W., 1970, Biometrika, 57, 97
 Hoffman D. I., Harrison T. E., Coughlin J. L., McNamara B. J., Holtzman J. A., Taylor G. E., Vestrand W. T., 2008, AJ, 136, 1067
 Horne K., 1986, PASP, 98, 609
 Horvat M., Conroy K. E., Jones D., Prša A., 2019, ApJS, 240, 36
 Jones D., Boffin H. M. J., Sowicka P., Miszalski B., Rodríguez-Gil P., Santand er-García M., Corradi R. L. M., 2019, MNRAS, 482, L75
 Kalomeni B., Nelson L., Rappaport S., Molnar M., Quintin J., Yakut K., 2016, ApJ, 833, 83
 Kochanek C. S. et al., 2017, PASP, 129, 104502
 Kurtz D. W., 1982, MNRAS, 200, 807
 —, 1985, MNRAS, 213, 773
 Kurtz D. W. et al., 2020, MNRAS, 494, 5118

- Lenz P., Breger M., 2005, *Communications in Asteroseismology*, 146, 53
- Lindegren L. et al., 2018, *A&A*, 616, A2
- Loeb A., Gaudi B. S., 2003, *ApJ*, 588, L117
- Luri X. et al., 2018, *A&A*, 616, A9
- Metropolis N., Rosenbluth A. W., Rosenbluth M. N., Teller A. H., Teller E., 1953, *J. Chem. Phys.*, 21, 1087
- Nelson L. A., Senhadji A., 2019, in *American Astronomical Society Meeting Abstracts*, Vol. 234, *American Astronomical Society Meeting Abstracts #234*, p. 215.02
- Paxton B., Bildsten L., Dotter A., Herwig F., Lesaffre P., Timmes F., 2011, *ApJS*, 192, 3
- Paxton B. et al., 2015, *ApJS*, 220, 15
- Pepper J., Kuhn R. B., Siverd R., James D., Stassun K., 2012, *PASP*, 124, 230
- Pepper J. et al., 2007, *PASP*, 119, 923
- Pollacco D. L. et al., 2006, *PASP*, 118, 1407
- Prša A. et al., 2016, *ApJS*, 227, 29
- Ricker G. R. et al., 2015, *Journal of Astronomical Telescopes, Instruments, and Systems*, 1, 014003
- Shappee B. J. et al., 2014, *ApJ*, 788, 48
- Skrutskie M. F. et al., 2006, *AJ*, 131, 1163
- Soberman G. E., Phinney E. S., van den Heuvel E. P. J., 1997, *A&A*, 327, 620
- Tauris T. M., van den Heuvel E. P. J., 2006, *Formation and evolution of compact stellar X-ray sources*, Vol. 39, pp. 623–665
- van Kerkwijk M. H., Rappaport S. A., Breton R. P., Justham S., Podsiadlowski P., Han Z., 2010, *ApJ*, 715, 51
- Verbunt F., Zwaan C., 1981, *A&A*, 100, L7
- Webbink R. F., 1976, *ApJ*, 209, 829
- Zhou G. et al., 2018, *ApJ*, 854, 109