

# TESS Cycle 1 observations of roAp stars with 2-min cadence data

D. L. Holdsworth,<sup>1</sup>★ M. S. Cunha,<sup>2</sup> D. W. Kurtz,<sup>3,1</sup> V. Antoci,<sup>4,5</sup> D. R. Hey,<sup>6,5</sup>  
 D. M. Bowman,<sup>7</sup> O. Kobzar,<sup>8</sup> D. L. Buzasi,<sup>9</sup> E. Niemczura,<sup>10</sup> D. Ozuyar,<sup>11</sup> F. Shi,<sup>12,13</sup>  
 R. Szabó,<sup>14,15</sup> A. Samadi,<sup>16</sup> Zs. Bognár,<sup>14,15</sup> L. Fox-Machado,<sup>17</sup> V. Khalack,<sup>8</sup>  
 O. Kochukhov,<sup>18</sup> M. Lares-Martiz,<sup>19</sup> C. C. Lovekin,<sup>20</sup> P. Mikołajczyk,<sup>10</sup>  
 D. Mkrtichian,<sup>21</sup> J. Pascual-Granado,<sup>19</sup> E. Paunzen,<sup>22</sup> Á. Sódor,<sup>23</sup> J. Sikora,<sup>24</sup> T. Yang,<sup>25</sup>  
 A. David-Uraz,<sup>26</sup> A. Derekas,<sup>27</sup> A. García Hernández,<sup>28</sup> J. A. Guzik,<sup>29</sup> S. J. Murphy,<sup>6,5</sup>  
 A. Ramón Ballesta,<sup>19</sup> B. Smalley,<sup>30</sup> and I. Stateva<sup>31</sup>

Author affiliations shown in Appendix A

26 April 2021

## ABSTRACT

We present the results of a systematic search for new rapidly oscillating Ap (roAp) stars using the 2-min cadence data collected by the *Transiting Exoplanet Survey Satellite* (TESS) during its Cycle 1 observations. We identify 12 new roAp stars. Amongst these stars we find the roAp star with the longest pulsation period, another with the shortest rotation period, and six with multiperiodic variability. In addition to these new roAp stars, we present an analysis of 44 known roAp stars observed by TESS during Cycle 1, providing the first high-precision and homogeneous sample of a significant fraction of the known roAp stars. The TESS observations have shown that almost 60 per cent (33) of our sample of stars are multiperiodic, providing excellent cases to test models of roAp pulsations, and from which the most rewarding asteroseismic results can be gleaned. We report four cases of the occurrence of rotationally split frequency multiplets that imply different mode geometries for the same degree modes in the same star. This provides a conundrum in applying the oblique pulsator model to the roAp stars. Finally, we report the discovery of non-linear interactions in  $\alpha$  Cir (TIC 402546736, HD 128898) around the harmonic of the principal mode – this is only the second case of such a phenomenon.

**Key words:** asteroseismology – stars: chemically peculiar – stars: oscillations – techniques: photometric – stars: variables – stars: individual

## 1 INTRODUCTION

The rapidly oscillating Ap (roAp) stars are a rare subset of the chemically peculiar, magnetic, Ap stars. They are found at the base of the classical instability strip where it intersects the main-sequence, and range in age from the zero-age main-sequence to beyond the terminal-age main-sequence. Since their discovery (Kurtz 1978, 1982) there have been several attempts to enlarge, and study, the sample of roAp stars. These have included targeted and survey ground-based photometry (e.g., Martínez et al. 1991; Nelson & Kreidl 1993; Martínez & Kurtz 1994b; Paunzen et al. 2012; Holdsworth et al. 2014a; Joshi et al. 2016; Paunzen et al. 2018), time-resolved high resolution spectroscopic observa-

tions (e.g., Elkin et al. 2008; Freyhammer et al. 2008a; Kochukhov et al. 2013) and most recently space photometry (e.g., Gruberbauer et al. 2011; Kurtz et al. 2011; Balona et al. 2011; Weiss et al. 2016; Hey et al. 2019; Balona et al. 2019; Cunha et al. 2019; Holdsworth 2021). To date, including the results presented here, there are 88 roAp stars known in the literature.

The overarching class of Ap stars host strong, stable, global magnetic fields that can have field strengths of up to 34 kG (Babcock 1960; Mathys 2017). The magnetic field is thought to be the main factor in braking the rotation velocity of the star (Stepień 2000), such that the Ap stars show rotation period of between a few days and perhaps centuries (North 1984; Sikora et al. 2019c; Mathys et al. 2020b). The presence of the magnetic field also serves to suppress convection, allowing for the stratification of elements in the stellar atmosphere (Michaud et al. 1981; Richer et al. 2000) through the

★ e-mail: dlholdsworth@uclan.ac.uk

effects of radiative levitation and gravitational settling. In particular, the radiative levitation leads to surface inhomogeneities of elements such as Ce, Pr, Nd, Sm, Eu and Tb, which can be overabundant by over a million times the solar value (e.g., Lüftinger et al. 2010).

These chemical spots, which are either bright or dark depending on the wavelength of observation, often form around the magnetic poles of the Ap stars (e.g., Kochukhov 2011), with the magnetic axis misaligned with respect to the rotation axis. Such a geometry leads to a non-sinusoidal modulation of the light curve over the stellar rotation period allowing for precise rotation periods to be measured, and the development of the oblique rotator model (Stibbs 1950). Such a modulation is also seen in the magnetic field strength (e.g., Mathys 2017; Hubrig et al. 2018), and spectral line strength. These rotationally modulated stars are commonly known as  $\alpha^2$  CVn stars.

The pulsations in the roAp stars are high overtone ( $n \gtrsim 15$ ), low degree ( $\ell \lesssim 3$ ) pressure (p) modes with frequencies in the range 0.7 – 3.6 mHz (60 – 310 d<sup>-1</sup>;  $P = 4.7 - 23.6$  min). The pulsation axis in these stars is misaligned to both the rotation and magnetic axes, leading to the development of the oblique pulsator model (Kurtz 1982; Shibahashi & Saio 1985; Bigot & Dziembowski 2002; Bigot & Kurtz 2011). Such a configuration means that the pulsation mode is viewed from varying aspects over the rotation cycle of the star, leading to an apparent amplitude modulation of the mode. In a Fourier spectrum of a light curve, a multiplet is seen with  $2\ell + 1$  components for a non-distorted mode, with the components split from the central mode frequency by exactly the stellar rotation frequency.

The driving mechanism for the pulsations in roAp stars is still an area of active research. In many cases, it is currently thought that the pulsations are driven by the opacity ( $\kappa$ ) mechanism in the hydrogen ionisation layers in regions of the star where the magnetic field suppresses convection (Balmforth et al. 2001). However, this theory cannot reproduce observations of the highest frequency modes in some stars. In these cases Cunha et al. (2013) proposed that a mechanism linked to turbulent pressure plays a role in the excitation of these highest frequency modes.

A further conundrum linked to the excitation mechanism of the roAp star pulsations comes with the inspection of the theoretical instability strip of these stars. Cunha (2002), based on the models of Balmforth et al. (2001), calculated the extent of the theoretical instability strip for roAp stars and found a much hotter blue edge than the observations mapped, and that some known roAp stars were cooler than the red edge. It is still unclear why the theoretical models do not match the observational evidence.

Of particular scientific interest are the roAp stars which show multiple pulsation modes as these stars provide the ability to constrain structural models of Ap stars. In the absence of a magnetic field, the pulsation modes of an roAp star would form a series of alternating degree modes that are equally spaced in frequency, as is expected in the asymptotic regime for high radial-order acoustic oscillations (Shibahashi 1979; Tassoul 1980). The acoustic waves in the outer layers of the roAp stars, in the presence of the strong magnetic field, become magnetoacoustic in nature which consequently changes the mode frequencies (Cunha & Gough 2000; Cunha 2006; Saio & Gautschy 2004). In most cases, the frequency separation between consecutive modes is not significantly affected, still enabling the use of asteroseismic techniques to help constraining the stellar properties as would be done for non-magnetic stars (Cunha et al. 2003; Deal et al. 2021). Nevertheless, at some particular frequencies the coupling between the magnetic field and pulsations is optimal and the pulsation spectrum can be significantly modified.

It is the multiperiodic stars, which in some cases show significant shifts from the expected non-magnetic frequency pattern, that provide the most scientific return (e.g. Cunha 2001; Gruberbauer et al. 2008).

The *Transiting Exoplanet Survey Satellite* (TESS; Ricker et al. 2015) is surveying almost the entire sky, in 1-month long strips known as *sectors*, in the search for transiting exoplanets. As such, it is collecting high precision photometric data on millions of stars at 2-min and 30-min cadences (with a 20-s cadence added in the extended mission). The Cycle 1 observations, conducted in the first year of the primary mission, covered the southern ecliptic hemisphere, with Cycle 2 observing the northern ecliptic hemisphere. Such an observational data set provides the opportunity to perform an unbiased search for high-frequency pulsations in a significant selection of stars.

This work utilises the 2-min cadence TESS Cycle 1 observations (sectors 1 through 13) to search for high-frequency pulsations in main-sequence stars hotter than 6000 K. We present the observational results here, and will, in a future work, use these results to perform an ensemble study of the properties of the roAp stars. The paper is laid out as follows: in Section 2 we introduce our data sample; in Section 3 we provide details of spectroscopic follow up observations; in Section 4 we present the results of our search. In Section 5 we present candidate roAp stars where we have insufficient data to confirm their true nature, and in Section 6 we draw our conclusions.

## 2 DATA SAMPLE AND SEARCH STRATEGY

To construct our catalogue of target stars, we took the 2-min target list for each sector of observations from TESS<sup>1</sup> which consisted of 20 000 stars per sector. These lists were then crossmatched with TESS Input Catalogue (TIC) version 8 with 3 arcsec radius. This gave us access to the secondary information in the TIC to refine our targets. Using only the  $T_{\text{eff}}$  parameter, we selected stars with temperatures of 6000 K or hotter. This amounted to about 7500 stars per sector, with 101 847 light curves in total, corresponding to 50 703 unique stars (with 15 363 stars observed in more than one sector). The light curves for each star in all available sectors were downloaded from the Mikulski Archive for Space Telescopes (MAST) server. These data have been processed with the Science Processing Operations Center (SPOC) pipeline. In the following we used the Pre-search Data Conditioning Simple Aperture Photometry (PDC\_SAP) data unless otherwise stated.

Initially, two independent analysis techniques were implemented for the search of pulsational variability. For one team, each individual light curve was automatically prewhitened to a frequency of 0.23 mHz (20 d<sup>-1</sup>) to an amplitude limit of the approximate noise level between 2.3 – 3.3 mHz (200 – 300 d<sup>-1</sup>) to remove instrumental artefacts and any low-frequency signal whose window function affects the noise at high frequency. If multiple sectors for a target are available, the data were then combined. An amplitude spectrum of the light curve was calculated to the Nyquist frequency of 4.2 mHz (360 d<sup>-1</sup>), which also included a calculation of the false alarm probability (FAP).

From these amplitude spectra, we selected stars which showed peaks with frequencies > 0.52 mHz (> 45 d<sup>-1</sup>) and a corresponding FAP of < 0.1. The amplitude spectra of these stars were then plotted

<sup>1</sup> <https://tess.mit.edu/observations/target-lists/>

for visual inspection. This totalled 6713 stars, of which 2125 stars have multi-sector observations.

A second team used a complementary method which calculated the skewness of the amplitude spectrum of the MAST PDC\_SAP data above 0.46 mHz ( $40 \text{ d}^{-1}$ ). Where multiple sectors of data were available, they were combined to a single light curve. If the skewness was greater than 5, then all peaks in the amplitude spectrum with FAP values below 0.05 were extracted and the star flagged as variable for later human inspection. This method produced a total of 189 variable star candidates.

These two lists were then combined to produce a master list where each star and light curve was subjected to human inspection. An initial pass was made to remove false positive detections. A detection was determined to be false positive if the S/N of the high-frequency signal was below 4.0; if there was obvious contamination from a low-frequency harmonic series; or if the amplitude spectrum displayed obvious characteristics of either  $\delta$  Sct stars (with many modes in the  $\approx 0.2 - 1.0$  mHz range), sdBV stars (stars with pulsations in low and high-frequency ranges) or pulsating white dwarfs (modes with amplitudes significantly greater than those known in roAp stars). The final sample consisted of 163 stars.

This final sample was distributed amongst the members of the TESS Asteroseismic Science Consortium (TASC) Working Group 4 (WG4) for detailed analysis to confirm the presence of a positive roAp detection, and to extract rotation periods from the light curves. This process identified 12 new roAp stars previously unreported in the literature and 10 roAp stars discovered through TESS observations and reported by either Cunha et al. (2019) or Balona et al. (2019). There are also positive detections of pulsations in 31 roAp stars known prior to the launch of TESS, with 3 roAp stars where TESS did not detect pulsational variability as their low amplitude modes are below the TESS detection limit. Finally, we present 5 roAp candidate stars where there is inconclusive evidence as to whether the star is truly an roAp pulsator.

### 3 SPECTROSCOPIC OBSERVATIONS

For a subset of our sample, we have made new spectroscopic observations to confirm or detect the chemically peculiar nature of the star. Most of these observations were obtained with the SpUpNIC long-slit spectrograph (Crause et al. 2016) mounted on the 1.9-m telescope of the South African Astronomical Observatory (SAAO). The observations were made in the blue part of the spectrum ( $\sim 3900 - 5100 \text{ \AA}$ ) with grating 4 which achieves a resolution of about  $0.6 \text{ \AA}$  per pixel, with some stars also having a red spectrum ( $\sim 6100 - 7150 \text{ \AA}$ ) obtained with grating 6 with a resolution of  $1.4 \text{ \AA}$  per pixel. The data were reduced in the standard way, with wavelength calibrations provided by arc spectra obtained directly after the science observation.

Where possible, we also observed some stars with the Southern African Large Telescope (SALT; Buckley et al. 2006), utilising the High Resolution Spectrograph (HRS; Bramall et al. 2010; Crause et al. 2014). HRS is a dual-beam spectrograph with wavelength coverage of  $3700 - 5500 \text{ \AA}$  and  $5500 - 8900 \text{ \AA}$ . The observations were automatically reduced using the SALT custom pipeline, which is based on the ESO's MIDAS pipeline (Kniazev et al. 2017, 2016).

Spectral classification with these observations was made by comparison with MK standard stars, following the procedure of Gray & Corbally (2009). The observations were normalised using an automated spline fitting procedure in the iSPEC spectral analysis software (Blanco-Cuaresma et al. 2014; Blanco-Cuaresma 2019),

with the MK standards plotted for comparison. With the classification as a starting point, we computed synthetic spectra using iSPEC to estimate effective temperatures. The synthesis used the SPECTRUM code (Gray & Corbally 1994) with a MARCS Gaia-ESO survey model atmosphere (Gustafsson et al. 2008) and a VALD line list (Ryabchikova et al. 2015). Spectra were synthesised with a fixed  $\log g = 4.0$  ( $\text{cm s}^{-2}$ ), solar metallicity and zero microturbulence and macroturbulence. Although these parameters are not refined for each star, they are adequate given the low resolution of most of the data. Spectra were synthesised in temperature steps of 100 K, with the result taken as the best fit to the wings of the observed Balmer lines.

### 4 RESULTS

Here we present the results of our search and analysis of new and known roAp stars in TESS Cycle 1 data. We provide an overview of the targets to be discussed in Table 1 and divide this section, and the table, into subsections addressing new discoveries reported in this work, previous discoveries reliant on the TESS data, and TESS observations of known roAp stars. Finally, we provide a list of candidate roAp stars where the currently available data do not allow us to confirm the nature of these stars.

Throughout this section, to keep homogeneity of the results, we present the numerical results of a single team, hereafter named the reference team. These results have been corroborated by the WG4 contribution to the analysis of the target stars. Where discrepancies were identified between the reference team and the WG4 members, secondary checks were conducted to ensure accurate results are presented.

To avoid repetition, we outline here the process by which each star was analysed in detail, unless a different method is detailed in the appropriate star's subsection. Where available, we combined multi-sector observations to produce a single light curve for analysis. Using the original PDC\_SAP data, we calculated a Discrete Fourier Transform, following Kurtz (1985) in the low-frequency range (to  $0.11 \text{ mHz}$ ;  $10 \text{ d}^{-1}$ ) in the search for rotational variability. Where present, we fitted a harmonic series by non-linear least squares to determine the rotation frequency. Where we found a harmonic to have a stronger signal, we derived the rotation frequency from that signal as it can be more precisely determined.

Subsequently, with the astrophysical information extracted, we removed the rotation signal and any instrumental artefacts through iterative pre-whitening of the light curve to a noise level determine in the range  $2.3 - 3.3 \text{ mHz}$  ( $200 - 300 \text{ d}^{-1}$ ). This serves to make the noise in the amplitude spectrum white when considering the errors on pulsation signals. Any pulsation signals found were extracted through non-linear least squares fitting. Both the derived rotation period and the pulsation frequency(ies) for each star are listed in Table 1. We provide only the *pulsation mode* frequencies and not rotationally split sidelobes; any sidelobes detected are mentioned in the text, and indicated on the appropriate figure.

Since the launch of TESS, there have been two papers addressing the discovery of roAp stars in the new data set. The TESS Ap first light paper (Cunha et al. 2019) focused on the analysis of the first and second TESS sectors, while a paper by Balona et al. (2019) analysed data from at sectors 1-7. We do not present an in depth analysis of the stars addressed in those papers again, unless there are more data available, or significant differences in the results were found. We do, however, include these stars in Table 1 to provide a complete inventory of TESS observations of roAp stars in Cycle 1.

**Table 1.** Details of the stars analysed in these paper. The columns provide the TIC identifier and star name, the TIC v8.1 TESS magnitude, the spectral type, where ‘\*’ denotes a classification derived in this work, the effective temperature, as provided in the TIC, the sectors in which TESS observed the target in Cycle 1. The final three columns provide the stellar rotation period derived in this work, with a † denoting the first detection of this period, the pulsation frequency(ies) found in this work (note that sidelobes that arise from oblique pulsation are not listed), and the pulsation amplitudes seen in the TESS data. A ‘\*\*\*’ denotes rotation periods derived from rotationally split sidelobes. A ‘\*\*\*\*’ denotes a tentative detection.

TIC	HD/TYC name	TESS mag	Spectral type	$T_{\text{eff}}^{\text{TIC}}$ (K)	Sectors	$P_{\text{rot}}$ (d)	Pulsation frequency (mHz)	Pulsation amplitude (mmag)
<i>New TESS roAp stars this work</i>								
96315731	51203	10.18	Ap SrEuCr	7100	6,7	$6.6713 \pm 0.0007$	$1.91271 \pm 0.00001$	$0.125 \pm 0.011$
119327278	45698	8.08	A2 SrEu	8540	6,7	$1.08457 \pm 0.00003$	$0.20583 \pm 0.00001$ $2.43755 \pm 0.00001$	$0.036 \pm 0.004$ $0.041 \pm 0.004$
170586794	107619	8.47	F5 p EuCr*	6590	10	$10.31 \pm 0.04^{\dagger}$	$1.76151 \pm 0.00003$	$0.045 \pm 0.006$
176516923	38823	7.12	A5 SrEuCr	7660	6	$8.782 \pm 0.003$	$1.76019 \pm 0.00005$ $1.85820 \pm 0.00004$ $1.91302 \pm 0.00005$	$0.023 \pm 0.004$ $0.006 \pm 0.004$ $0.023 \pm 0.004$
178575480	55852	8.83	Ap SrEuCr	7720	7	$4.7788 \pm 0.0005$	$2.23870 \pm 0.00004$ $2.39524 \pm 0.00004$	$0.010 \pm 0.007$ $0.052 \pm 0.008$
294266638	6021-415-1	9.74	Ap SrEu*	7230	8	No signature	$1.56421 \pm 0.00002$ $1.62882 \pm 0.00002$	$0.119 \pm 0.012$ $0.149 \pm 0.012$
294769049	161423	9.17	Ap SrEu(Cr)	7760	12,13	$10.4641 \pm 0.0007^{\dagger}$	$2.26044 \pm 0.00001$	$0.054 \pm 0.006$
310817678	88507	9.60	Ap SrEu(Cr)	8230	9,10	$2.75003 \pm 0.00008^{\dagger}$	$1.170979 \pm 0.000019$ $1.205272 \pm 0.000003$ $1.229890 \pm 0.000030$	$0.049 \pm 0.032$ $0.337 \pm 0.032$ $0.133 \pm 0.032$
356088697	76460	9.53	Ap SrEuCr*	7110	9-11	No signature	$0.646584 \pm 0.000008$	$0.053 \pm 0.006$
380651050	176384	8.19	F0/F2	6490	13	$4.19 \pm 0.01^{\dagger}$	$1.85817 \pm 0.00002$ $1.87771 \pm 0.00002$	$0.059 \pm 0.004$ $0.045 \pm 0.004$
387115314	9462-347-1	9.56	A5	7660	13	$5.264 \pm 0.002^{\dagger}$	$1.319020 \pm 0.000039$ $1.328382 \pm 0.000020$ $1.356225 \pm 0.000006$	$0.051 \pm 0.009$ $0.101 \pm 0.009$ $0.375 \pm 0.009$
466260580	9087-1516-1	11.64	Ap EuCr*	6830	13	No signature	$1.384083 \pm 0.000007$ $1.340060 \pm 0.00002$	$0.274 \pm 0.009$ $0.453 \pm 0.035$
<i>roAp stars discovered by TESS in the literature</i>								
12968953	217704	10.09	Ap SrEuCr*	7880	2	No signature	$1.27047 \pm 0.00005$ $1.25875 \pm 0.00004$ $1.26961 \pm 0.00003$ $1.33782 \pm 0.00003$	$0.071 \pm 0.013$ $0.083 \pm 0.013$ $0.064 \pm 0.013$ $0.099 \pm 0.013$
17676722	63773	8.61	Ap SrEuCr*	8320	7	$1.5995 \pm 0.0003$	$1.941545 \pm 0.000002$	$0.671 \pm 0.006$
41259805	43226	8.84	Ap SrEu(Cr)*	8360	1-8,10-13	$1.714489 \pm 0.000002$	$2.311039 \pm 0.000001$ $2.326368 \pm 0.000003$	$0.025 \pm 0.002$ $0.013 \pm 0.002$
49818005	19687	9.36	FpSrEu(Cr)*	7300	4	No signature	$1.641910 \pm 0.000008$	$0.267 \pm 0.009$
152808505	216641	7.88	Fp EuCr*	6430	1	$1.877 \pm 0.006^{***}$	$1.36260 \pm 0.00005$ $1.38593 \pm 0.00002$ $1.39119 \pm 0.00002$	$0.019 \pm 0.004$ $0.054 \pm 0.004$ $0.041 \pm 0.004$
156886111	47284	9.11	Ap SrEuCr*	7540	6,7	$6.8580 \pm 0.0003$	$0.063251 \pm 0.000002$ $1.274574 \pm 0.000021$ $1.302738 \pm 0.000014$ $1.317816 \pm 0.000008$	$0.366 \pm 0.006$ $0.039 \pm 0.006$ $< 0.020$ $0.107 \pm 0.006$
259587315	30849	8.69	Ap SrCrEu	7720	4,5	$15.776 \pm 0.005$	$0.904064 \pm 0.000009$ $0.905469 \pm 0.000013$ $0.931050 \pm 0.000003$ $0.950735 \pm 0.000022$	$0.071 \pm 0.005$ $0.050 \pm 0.005$ $0.154 \pm 0.005$ $0.029 \pm 0.005$
349945078	57040	8.92	A2 EuCr	7200	6-9	$13.4256 \pm 0.0008$	$2.126381 \pm 0.000002$ $2.186645 \pm 0.000005$	$0.140 \pm 0.004$ $0.063 \pm 0.004$
350146296	63087	9.17	F0p EuCr*	7680	1-13	$2.66387 \pm 0.00002$	$3.4031468 \pm 0.0000078$ $3.4424674 \pm 0.0000021$ $3.4839854 \pm 0.0000015$ $3.5232955 \pm 0.0000005$ $3.5626041 \pm 0.0000031$	$0.005 \pm 0.002$ $0.020 \pm 0.002$ $0.026 \pm 0.002$ $0.081 \pm 0.002$ $0.013 \pm 0.002$
431380369	20880	7.82	Ap Sr(EuCr)	8230	2,6,13	$5.19716 \pm 0.00005$	$0.805252 \pm 0.000002$ $0.815837 \pm 0.000002$ $0.860412 \pm 0.000001$ $0.872662 \pm 0.000003$	$0.022 \pm 0.003$ $0.016 \pm 0.003$ $0.032 \pm 0.003$ $0.013 \pm 0.003$

Table 1 – continued

TIC	HD/TYC name	TESS mag	Spectral type	$T_{\text{eff}}^{\text{TIC}}$ (K)	Sectors	$P_{\text{rot}}$ (d)	Pulsation frequency (mHz)	Pulsation amplitude (mmag)
<i>Known roAp stars prior to TESS launch</i>								
6118924	116114	6.77	F0Vp SrCrEu	7670	10	No signature	$0.76923 \pm 0.00004$	$0.020 \pm 0.003$
33601621	42659	6.60	Ap SrCrEu	7880	6	$2.6627 \pm 0.0003$	$1.74757 \pm 0.00003$	$0.039 \pm 0.005$
35905913	132205	8.50	Ap EuSrCr	7510	11,12	$7.513 \pm 0.001$	No signal in TESS data	$< 0.025$
44827786	150562	9.48	F5Vp SrCrEu	7350	12	No signature	$1.54700 \pm 0.00002$	$0.181 \pm 0.013$
49332521	119027	9.66	Ap SrEu(Cr)	6940	11	No signature	$1.55040 \pm 0.00005$	$0.068 \pm 0.013$
							$1.57070 \pm 0.00005$	$0.053 \pm 0.011$
							$1.83521 \pm 0.00004$	$0.071 \pm 0.011$
							$1.87490 \pm 0.00005$	$0.059 \pm 0.011$
							$1.88779 \pm 0.00002$	$0.174 \pm 0.011$
							$1.91338 \pm 0.00002$	$0.160 \pm 0.011$
							$1.94029 \pm 0.00002$	$0.184 \pm 0.012$
69855370	213637	9.19	A (pEuSrCr)	6610	2	No signature	$1.94207 \pm 0.00002$	$0.142 \pm 0.012$
							$1.42245 \pm 0.00001$	$0.120 \pm 0.007$
							$1.45237 \pm 0.00001$	$0.172 \pm 0.007$
93522454	143487	9.11	A3 SrEuCr	7110	12	No signature	No signal in TESS data	$< 0.030$
125297016	69013	9.21	Ap EuSr	7010	7	No signature	$1.47519 \pm 0.00007^{***}$	$0.042 \pm 0.009$
136842396	9289	9.20	A(p) SrEuCr	7750	3	$8.660 \pm 0.006$	$1.55361 \pm 0.00005$	$< 0.026$
139191168	217522	7.18	A5 SrEuCr	6920	1	No signature	$1.57327 \pm 0.00005$	$0.059 \pm 0.009$
							$1.58496 \pm 0.00001$	$0.262 \pm 0.009$
							$1.61549 \pm 0.00003$	$< 0.038$
							$1.20088 \pm 0.00002$	$0.030 \pm 0.003$
							$1.20169 \pm 0.00003$	$0.028 \pm 0.003$
146715928	92499	8.72	A2p SrEuCr	7730	9,10	No signature	$1.20579 \pm 0.00002$	$0.032 \pm 0.003$
							$1.21513 \pm 0.00002$	$0.031 \pm 0.003$
							$1.54260 \pm 0.00002$	$0.034 \pm 0.005$
							$1.58221 \pm 0.00001$	$0.043 \pm 0.005$
167695608	8912-1407-1	11.69	F0p SrEu(Cr)	7180	1-4,6-13	No signature	$1.512238 \pm 0.000004$	$0.044 \pm 0.009$
							$1.532161 \pm 0.000001$	$0.171 \pm 0.009$
							$1.554816 \pm 0.000001$	$0.135 \pm 0.009$
							$1.557383 \pm 0.000002$	$0.074 \pm 0.009$
							$1.564767 \pm 0.000002$	$0.079 \pm 0.009$
							$1.586081 \pm 0.000003$	$0.052 \pm 0.009$
168383678	96237	9.25	A4 SrEuCr	7410	9	$\sim 21$ or $\sim 42$	$1.19610 \pm 0.00003$	$0.077 \pm 0.010$
170419024	151860	8.53	Ap SrEu(Cr)	6630	12	No signature	$0.85014 \pm 0.00002$	$0.085 \pm 0.007$
173372645	154708	8.43	Ap SrEuCr	6920	12	$5.363 \pm 0.001$	$1.36330 \pm 0.00004$	$0.045 \pm 0.007$
							No signal in TESS data	$< 0.030$
							$1.84163 \pm 0.00002^{***}$	$0.013 \pm 0.003$
189996908	75445	6.89	Ap SrEu(Cr)	7610	8,9	No signature	$2.69844 \pm 0.00005$	$0.030 \pm 0.006$
211404370	203932	8.56	Ap SrEu	7540	1	$6.44 \pm 0.01$	$2.80476 \pm 0.00002$	$0.082 \pm 0.006$
237336864	218495	9.24	Ap EuSr	8120	1	$4.2006 \pm 0.0001$	$2.09827 \pm 0.00003$	$0.056 \pm 0.008$
							$2.22073 \pm 0.00002$	$< 0.035$
							$2.24862 \pm 0.00002$	$0.103 \pm 0.008$
							$2.26123 \pm 0.00001$	$0.159 \pm 0.008$
							$1.436302 \pm 0.000006$	$0.682 \pm 0.012$
268751602	12932	9.98	Ap SrEuCr	7320	3	No signature	$2.557764 \pm 0.000043$	$0.013 \pm 0.002$
279485093	24712	5.72	A9Vp SrEuCr	7280	5	$12.578 \pm 0.008$	$2.604135 \pm 0.000034$	$0.018 \pm 0.002$
							$2.619775 \pm 0.000013$	$0.046 \pm 0.002$
							$2.652959 \pm 0.000005$	$0.113 \pm 0.002$
							$2.687440 \pm 0.000003$	$0.170 \pm 0.002$
							$2.720639 \pm 0.000003$	$0.199 \pm 0.002$
							$2.755166 \pm 0.000025$	$0.025 \pm 0.002$
							$2.791338 \pm 0.000014$	$0.040 \pm 0.002$
							$2.805580 \pm 0.000016$	$0.036 \pm 0.002$
							$1.42797 \pm 0.00001$	$0.132 \pm 0.005$
							$1.114117 \pm 0.000005$	$< 0.055$
280198016	83368	6.06	A8V SrCrEu	7710	9	$2.8517 \pm 0.0002$	$1.557657 \pm 0.000002$	$0.179 \pm 0.007$
315098995	84041	9.20	Ap SrEuCr	7780	8,9	$3.6884 \pm 0.0004$	$2.402163 \pm 0.000007$	$0.063 \pm 0.006$
322732889	99563	8.47	F0 Sr	7940	9	$2.9114 \pm 0.0002$		
326185137	6532	8.35	Ap SrEuCr	8500	3	$1.9447 \pm 0.0001$		

**Table 1** – *continued*

TIC	HD/TYC name	TESS mag	Spectral type	$T_{\text{eff}}^{\text{TIC}}$ (K)	Sectors	$P_{\text{rot}}$ (d)	Pulsation frequency (mHz)	Pulsation amplitude (mmag)
<i>Known roAp stars prior to TESS launch – continued</i>								
340006157	60435	8.72	A3 SrEu	8160	3,6-10,13	$7.6797 \pm 0.0001$	$1.29573 \pm 0.00063$ $1.31293 \pm 0.00019$ $1.32446 \pm 0.00030$ $1.35251 \pm 0.00006$ $1.38056 \pm 0.00017$ $1.40721 \pm 0.00012$ $1.43233 \pm 0.00012$ $1.45727 \pm 0.00036$	$0.012 \pm 0.003$ $0.042 \pm 0.003$ $0.025 \pm 0.003$ $0.126 \pm 0.003$ $0.043 \pm 0.003$ $0.064 \pm 0.003$ $0.064 \pm 0.003$ $0.021 \pm 0.003$
348717688	19918	9.12	A5 SrEuCr	7790	1,12,13	No signature	$1.37159815 \pm 0.00000146$ $1.37673847 \pm 0.00000098$ $1.39872517 \pm 0.00000194$ $1.44003439 \pm 0.00000199$ $1.48119405 \pm 0.00000130$ $1.51008560 \pm 0.00000006$ $1.53991119 \pm 0.00000191$	$0.039 \pm 0.005$ $0.058 \pm 0.005$ $0.029 \pm 0.005$ $0.029 \pm 0.005$ $0.044 \pm 0.005$ $0.997 \pm 0.005$ $0.030 \pm 0.005$
363716787	161459	10.01	Ap EuSrCr	7330	13	$5.966 \pm 0.001$	$1.39089 \pm 0.00003$ $1.42253 \pm 0.00004$	$0.081 \pm 0.017$ $0.092 \pm 0.017$
368866492	166473	7.73	A5p SrCrEu	7510	13	No signature	$1.884520 \pm 0.000017$ $1.888201 \pm 0.000030$ $1.891891 \pm 0.000005$	$0.070 \pm 0.005$ $0.041 \pm 0.005$ $0.274 \pm 0.005$
369845536		12.90	A7Vp Eu(Cr)	7930	13	$9.50 \pm 0.02$	$2.041601 \pm 0.000003$	$5.843 \pm 0.094$
394124612	218994	8.31	A3 Sr		1	$5.855 \pm 0.008$	$1.14433 \pm 0.00001$	$0.124 \pm 0.008$
394272819	115226	8.27	Ap Sr(Eu)	7630	11,12	$2.98827 \pm 0.00008$	$1.479334 \pm 0.000069$ $1.508904 \pm 0.000002$ $1.538487 \pm 0.000016$ $1.568072 \pm 0.000007$	$< 0.010$ $0.213 \pm 0.005$ $0.034 \pm 0.005$ $0.075 \pm 0.005$
402546736	128898	1.84	A7Vp SrCrEu		11,12	$4.4812 \pm 0.0005$	$2.2213028 \pm 0.0000133$ $2.2680569 \pm 0.0000105$ $2.3692228 \pm 0.0000117$ $2.3817256 \pm 0.0000086$ $2.4118973 \pm 0.0000013$ $2.4420726 \pm 0.0000003$ $2.4722512 \pm 0.0000010$ $2.5024429 \pm 0.0000117$ $2.5326183 \pm 0.0000205$ $2.5666565 \pm 0.0000091$	$0.009 \pm 0.001$ $0.012 \pm 0.001$ $0.011 \pm 0.001$ $0.015 \pm 0.001$ $0.100 \pm 0.001$ $0.428 \pm 0.001$ $0.118 \pm 0.001$ $0.011 \pm 0.001$ $0.006 \pm 0.001$ $0.014 \pm 0.001$
434449811	80316	7.69	Ap Sr(Eu?)	8180	8	$2.08862 \pm 0.00004$	$0.112311 \pm 0.000022$ $0.131686 \pm 0.000014$ $2.251749 \pm 0.000007$	$0.055 \pm 0.005$ $0.089 \pm 0.005$ $0.182 \pm 0.005$
469246567	86181	9.15	Ap Sr	7210	9,10	$2.05115 \pm 0.00006$	$2.663229 \pm 0.000015$ $2.694099 \pm 0.000003$ $2.728458 \pm 0.000012$	$< 0.020$ $0.276 \pm 0.007$ $0.071 \pm 0.007$
<i>Candidate roAp</i>								
1727745	113414	7.22	F7/F8	6150	10	$3.172 \pm 0.001^{\dagger}$	$2.29116 \pm 0.00004$	$0.035 \pm 0.006$
3814749	3748	9.73	A0/I IV/V	8420	3	$1.689 \pm 0.002$	$1.63100 \pm 0.00006$ $2.75700 \pm 0.00002$	$0.163 \pm 0.031$ $0.389 \pm 0.031$
158637987	10330	8.91	A9	7180	2,3	No signature	$2.72598 \pm 0.00002$ $2.74240 \pm 0.00001$	$0.022 \pm 0.005$ $0.031 \pm 0.005$
324048193	85892	7.74	Ap Si	11950	12,13	$4.2953 \pm 0.0001$	$3.15347 \pm 0.00002$	$0.020 \pm 0.004$
410163387	76276	9.50	Ap SrEuCr	7480	8	No signature	$1.43155 \pm 0.00003$	$0.075 \pm 0.011$

## 264 4.1 New TESS roAp stars


### 265 4.1.1 TIC 96315731

266 TIC 96315731 (HD 51203) is identified in the literature as an  
 267  $\alpha^2$  CVn with a spectral type of Ap SrEuCr (Houk 1982). The star  
 268 is reported to have a mean magnetic field modulus of  $7.9 \pm 0.5$  kG  
 269 (Chojnowski et al. 2019).

270 TIC 96315731 was observed by TESS during sectors 6 and

271 7, and shows clear signatures of rotation and pulsation, making  
 272 this the first report that TIC 96315731 is a new roAp star. From  
 273 the combined TESS light curve, we measure a rotation period of  
 274  $6.6713 \pm 0.0007$  d which is similar to the value of 6.675 d presented  
 275 by Netopil et al. (2017). We show a light curve phased on the rotation  
 276 period in the top panel of Fig. 1.

277 At high frequency, the star shows an obvious pulsation signal  
 278 at  $1.91271 \pm 0.00001$  mHz ( $165.258 \pm 0.001$  d $^{-1}$ ) which is flanked



**Figure 1.** Top: light curve of TIC96315731 phase folded on the derived rotation period of 6.67134 d. Bottom: the pulsation quintuple arising from a distorted quadrupole mode. The vertical dotted lines indicate the quintuplet components.


by two sidelobes with significant amplitude ( $S/N \sim 14$ ) separated from the pulsation by the rotation frequency, and two further sidelobes which have lower significance ( $S/N \sim 6$ ) separated by twice the rotation frequency. This is indicative of either a distorted dipole mode, or a quadrupole mode, although an analysis of the pulsation amplitude and phase as a function of rotation phase favours a distorted quadrupole mode. In the case of a distorted quadrupole mode, the pulsation phase does not change by the expected  $\pi$ -rad when a pulsation node crosses the line of sight, but rather results in a phase blip (see e.g., Holdsworth et al. 2016, 2018b, Shi et al. submitted). These peaks are shown in the bottom panel of Fig. 1.

#### 4.1.2 TIC 119327278

TIC 119327278 (HD 45698) was initially classified as an Ap star by Bidelman & MacConnell (1973) due to the presence of Sr absorption, with a revised spectral type of A2 SrEu provided by Renson et al. (1991). There is little information about this star in the literature other than a rotation period of 1.085 d (Netopil et al. 2017), and a lack of pulsation signal reported by Joshi et al. (2016).

TIC 119327278 was observed during sectors 6 and 7. The data show the star to have a rotation period of  $1.08457 \pm 0.00003$  d (Fig. 2) which is in agreement with the literature. Further to the harmonic series, we identify another mode that is typical of the  $\delta$  Sct stars, at a frequency of  $0.20583 \pm 0.00001$  mHz ( $17.783 \pm 0.001$  d $^{-1}$ ). Although it was not expected that Ap stars show these low-overtone modes as a result of magnetic suppression (Saio 2005), Murphy et al. (2020) have recently shown the existence of  $\delta$  Sct and roAp pulsations in the same star.

The amplitude spectrum at high frequency shows a multiplet of 4 significant components ( $\nu + 2\nu_{\text{rot}}$  is in the noise) of a presumed quintuplet (Fig. 2), with the highest peak represent-




**Figure 2.** Top: light curve of TIC 119327278 phase folded on the derived rotation period of 1.08457 d. Middle: view of the  $\delta$  Sct mode in this star. Bottom: the pulsation multiplet signature. The vertical dotted lines indicate the quintuplet components.

ing the pulsation mode at a frequency of  $2.43755 \pm 0.00001$  mHz ( $210.605 \pm 0.001$  d $^{-1}$ ). The components are split by the rotation frequency of the star as a result of oblique pulsation, thus confirming this star as a new roAp star. Given the presence of the low-overtone mode in TIC 119327278, it may be the second example of a  $\delta$  Sct-roAp hybrid star. Given the low amplitude of the pulsation mode and the relatively short rotation period, analysing the pulsation amplitude and phase over the rotation period results in inconclusive results with regards to the identification of the mode degree. A detailed study, beyond the scope of this work, is needed to resolve this problem.

#### 4.1.3 TIC 170586794

TIC 170586794 (HD 107619) was classified as a possible metallicity late-A type star by Houk (1982). The star was later defined to be an F0 type by Cannon & Pickering (1993) and an F0 type  $\delta$  Del star by Perry (1991). It is not a frequently studied object with



**Figure 3.** Top: the light curve of TIC 170586794 phased on a period of 10.305 d. Bottom: amplitude spectrum of the light curve showing a single pulsation mode.

the literature being mainly concerned with  $uvbyH\beta^2$  photometry. Among these studies, Perry & Christodoulou (1996) gave dereddened Strömgren colour indexes and metallicity of the star, while Perry (1991) provided  $(b - y)$ ,  $m_1$ ,  $c_1$  colour indexes and  $H\beta$  index of 0.303, 0.226, 0.432 and 2.682, respectively. The temperature of the star was derived as  $T_{\text{eff}} = 6504$  K by Stevens et al. (2017) and  $T_{\text{eff}} = 6564$  K by Ammons et al. (2006) from photometric investigations.

TESS observed TIC 170586794 during sector 10. The light curve shows the star to have a rotation period of  $10.305 \pm 0.041$  d, which has a double wave nature (Fig. 3). At high frequency, this star shows a single pulsation mode at  $1.76151 \pm 0.00003$  mHz ( $152.194 \pm 0.003$  d $^{-1}$ ; Fig. 3). With both the presence of a rotationally modulated light curve, and a high frequency pulsation, we revisited this star spectroscopically. Our SAAO classification spectrum (Fig. B1) shows this star to have overabundances of Eu II and Cr II, thus making it chemically peculiar. The Balmer lines in the spectrum are well matched to an F5 V star, thus making this a cool F5VpEuCr star, and confirming it as a new roAp star and making it one of the coolest roAp stars to date.

#### 4.1.4 TIC 176516923

TIC 176516923 (HD 38823) has several classifications in the literature: A5 SrEu (Vogt & Faundez 1979; Hensberge et al. 1981); a strongly magnetic A5p Sr star (Chen et al. 2017); and A5 SrEuCr (Renson & Manfroid 2009). Sikora et al. (2019a) confirmed the star as a magnetic CP star and provided several abundance measures

<sup>2</sup> Throughout we use the  $H\beta$  notation to refer to the Strömgren-Crawford index to avoid confusion between angle of magnetic obliquity which is given the symbol  $\beta$ .

and estimated the age to be  $\log t = 8.58^{+0.56}_{-0.76}$  Gyrs. Estimates of the effective temperature range from 6600 K to 7700 K, the latter of which agrees with the TIC temperate provided in Table 1.  $\log g$  values range from about 3.7 ( $\text{cm s}^{-2}$ ) to 4.6 ( $\text{cm s}^{-2}$ ), with most rotational velocity measurements suggesting a  $v \sin i \approx 22$   $\text{km s}^{-1}$ .

There are several studies related to the magnetic structure of TIC 176516923. The strong magnetic field of the star was first discovered by Romanyuk et al. (2016) after three observations made in 2009, with a  $\langle B_z \rangle$  value of 1.74 kG. They found a variable radial velocity within a small interval around  $-10$   $\text{km s}^{-1}$  which is significantly different from the result of  $1.40$   $\text{km s}^{-1}$  by Grenier et al. (1999). Based on this result, they concluded that such differences in radial velocity could be an indication of a variability on a scale of up to several years or a decade. Kudryavtsev et al. (2007) performed further spectroscopic observations and calculated radial velocities ranging from  $-5$   $\text{km s}^{-1}$  to  $-11$   $\text{km s}^{-1}$ . Therefore, they suspected that the star is in a binary system.

The rotation period of the star has been under debate for several years, with fits to both light and magnetic data being used to determine it (Hensberge et al. 1981; Kudryavtsev et al. 2006; Kudryavtsev & Romanyuk 2012). Most recently, Bernhard et al. (2020) calculated the period of the star as  $8.677 \pm 0.002$  d which is representative of the other periods reported.

TIC 176516923 was observed by TESS in sector 6 only. The data for this star provides an example where the SPOC pipeline has interfered with the astrophysical signal of the rotation signature, and injected noise to the light curve. We demonstrate this in Fig. 4 where the top panel compares the PDC\_SAP (grey) data to the SAP data (black). Using the SAP data, we extract a rotation period of  $8.782 \pm 0.003$  d, which agrees well with the literature.


The pulsation spectrum of this star consists of a quadrupole quintuplet with a missing central component, a dipole triplet also with a missing central component, and a further dipole triplet with a significant central component. Furthermore, there are two peaks with significant amplitude (labelled by the arrows in Fig. 4) that do not show any multiplet components.

The difference in the multiplet structures is puzzling in this star. Although different modes have different depth pulsation cavities, one would expect both dipole modes to show the very similar relative amplitudes since they are governed by the geometry of the star through  $i$ , the inclination angle, and  $\beta$ , the angle of magnetic obliquity. This difference may suggest different pulsation axes for the two dipole modes (as suggested for KIC 10195926; Kurtz et al. 2011). However, it is likely that we are seeing a depth dependence on the geometry which is being affected by different limb darkening weighting due to the stratified atmosphere of the star, as is proposed to be the case for HD 6532 (Kurtz & Holdsworth 2020). Another option is that the difference may result from the modes suffering different magnetic distortions in the atmosphere (Sousa & Cunha 2011; Quiral-Manosalva et al. 2018), due to their slightly different frequencies. Either way, the nature of these different multiplet structures is not clear, which clearly warrants a further, in-depth, study with additional multi-colour observations and detailed modelling.

#### 4.1.5 TIC 178575480

TIC 178575480 (HD 55852) has a spectral classification of Ap SrEuCr (Houk & Swift 1999a) and according to the TIC has an effective temperature of 7720 K. There is no additional information in the literature.


This star was observed in sector 7, allowing us to measure a



**Figure 4.** Top: comparison of the PDC\_SAP (grey) and SAP (black) light curves for TIC 176516923. Middle: SAP light curve phase folded on the derived rotation period of 8.7821 d. Bottom: amplitude spectrum of the PDC\_SAP light curve showing the pulsational variability. The vertical dotted lines indicate the multiplet components, with the arrows indicating possible further modes.

rotation period of  $4.7788 \pm 0.0005$  d. A phase folded light curve is shown in the top panel of Fig. 5.

There are two pulsation modes seen in this star: a singlet at a frequency of  $2.23870 \pm 0.00004$  mHz ( $193.424 \pm 0.004$  d<sup>-1</sup>), and a triplet with a central frequency of  $2.39524 \pm 0.00004$  mHz ( $206.949 \pm 0.003$  d<sup>-1</sup>). Since the two modes show different multiplet structures, it is logical to assume that the modes are of different degree, however their identification is unclear. Given that the large frequency separation in roAp stars ranges between  $\sim 30 - 100$   $\mu$ Hz, the separation of the two modes,  $\approx 156.5$   $\mu$ Hz, is plausibly 1.5 times the large frequency separation. However, this would require the two modes, and an unobserved mode in the middle, all being the same degree, which the lack of multiplet structure of the lower frequency mode suggests is not the case. If the separation is 2 times the large frequency separation, the two modes seen would be of different degree, as the structure suggests. Further observations of this star are needed to resolve this issue.



**Figure 5.** Top: the light curve of TIC 178575480 phase folded on the derived rotation period of 8.7821 d. Bottom: amplitude spectrum of the light curve showing the pulsational variability. The vertical dotted lines indicate the pulsation modes and the multiplet components of the high-frequency triplet.

#### 4.1.6 TIC 294266638

TIC 294266638 (TYC 6021-415-1) is a star with no mention in the literature, other than a Gaia parallax measurement of  $2.91 \pm 0.05$  mas, implying a distance of  $344.1 \pm 0.6$  pc (Gaia Collaboration et al. 2018).


We obtained a spectrum for this star with the SpUpNIC instrument to confirm its nature as an Ap star (Fig. B2). The spectrum shows enhancements of Sr and Eu leading to a classification of Ap SrEu. There are signs of Cr in the spectrum at 4111 Å, but no other lines of this element are clearly enhanced.

This star was observed in sector 8. There is no evidence of rotational variability in the TESS light curve, however there are clear pulsation signals, as shown in Fig. 6. We extract two pulsation modes from the amplitude spectrum, at  $1.56421 \pm 0.00002$  mHz ( $135.148 \pm 0.002$  d<sup>-1</sup>) and  $1.62882 \pm 0.00002$  mHz ( $140.730 \pm 0.002$  d<sup>-1</sup>). After prewhitening these peaks, there is still evidence of excess power in the residual spectrum. This is indicative of phase/amplitude variability. Furthermore, there is evidence of 2 additional modes at low amplitude around 1.6 mHz which are significant. However, given their broad nature of the peaks, a clear frequency value cannot be determined.


#### 4.1.7 TIC 294769049

TIC 294769049 (HD 161423) was classified as Ap SrEu(Cr) by Houk & Cowley (1975), and listed in the catalogue of Renson & Manfroid (2009) with a spectral type of A2 SrEu. TIC 294769049 has been the target in two searches for rapid variability; both Martinez & Kurtz (1994b) and Joshi et al. (2016) returned null results.

TESS observed this star during sectors 12 and 13, providing a light curve with a time base of 57.4 d. From this we measure a



**Figure 6.** Pulsation modes in TIC 294266638. After removing the obvious pulsations, excess power is remaining which is suggestive of frequency variability.



**Figure 7.** Top: phase folded light curve of TIC 294769049, folded on a period of 10.4641 d. Bottom: amplitude spectrum showing the only pulsation mode in this star, and the lack of a rotationally split multiplet.

456 rotation period of  $10.4641 \pm 0.0007$  d, and show a phase folded  
 457 light curve in Fig. 7. Analysis of the light curve at high frequency  
 458 shows a single pulsation mode at  $2.26044 \pm 0.00001$  mHz ( $195.302 \pm$   
 459  $0.001$  d $^{-1}$ ; Fig. 7). The lack of an obliquely split multiplet suggests  
 460 that either the pulsation axis is aligned with the rotation axis (thus  
 461  $\beta = 0$ , assuming the original oblique pulsator model), despite the  
 462 significant rotational variation, or the mode is an undistorted radial  
 463 mode. The former would imply that the spots causing the mean light  
 464 variation are not close to the pulsation pole (Kochukhov et al. 2004;  
 465 Alecian & Stift 2010), or that the pulsation axis is aligned with the  
 466 rotation axis, rather than the magnetic axis (Bigot & Dziembowski  
 467 2002).

#### 468 4.1.8 TIC 310817678

469 TIC 310817678 (HD 88507) was classified as Ap SrEu(Cr) by Houk  
 470 (1978) and in the catalogue of Renson & Manfroid (2009) it is  
 471 quoted as an Ap SrEu star. There is only one evaluation of the  
 472 atmospheric parameters given by the TIC:  $T_{\text{eff}} = 8230$  K and  $\log g =$   
 473  $4.08$  (cm s $^{-2}$ ).

474 TESS observed this star in sectors 9 and 10, providing a 51.8 d  
 475 time base. We measure a rotation period of  $2.75003 \pm 0.00008$  d  
 476 (Fig. 8). The pulsation spectrum in this star is rich, consisting of  
 477 a dipole triplet centred on  $1.17098 \pm 0.00002$  mHz ( $101.173 \pm$   
 478  $0.002$  d $^{-1}$ ), a quadrupole quintuplet with missing  $\nu_{\text{rot}}$  sidelobes  
 479 centred on  $1.205272 \pm 0.000003$  mHz ( $104.1355 \pm 0.0002$  d $^{-1}$ ), and  
 480 another dipole with a central frequency at  $1.22989 \pm 0.00003$  mHz  
 481 ( $106.263 \pm 0.003$  d $^{-1}$ ). The two dipole modes are separated by  
 482  $58.9$   $\mu$ Hz which is plausibly the large frequency separation for this  
 483 star.

484 The high-frequency dipole sidelobes are unlike those seen in  
 485 the other multiplets: they form doublets. We suspect that this is a  
 486 signature of magnetic distortion since despite the sidelobes having


487 greater amplitude than the low-frequency dipole mode, the central  
 488 component is entirely absent. This star is another example of how  
 489 the multiplet structures vary for the same degree mode.

#### 490 4.1.9 TIC 356088697


491 TIC 356088697 (HD 76460) was classified as an Ap Sr by Houk  
 492 & Cowley (1975) with Elkin et al. (2012) measuring a magnetic  
 493 field strength of 3.6 kG through partial Zeeman of Fe lines. The  
 494 authors also determined the effective temperature to be 7200 K and  
 495  $\nu \sin i = 3$  km s $^{-1}$ , although this value was at the resolution limit of  
 496 their data.

497 Observed in three consecutive TESS sectors (9-11), this star  
 498 shows no signature of rotation. To confirm its Ap nature, we obtained  
 499 a spectrum with SpUpNIC and derived a spectral class of Ap SrEuCr  
 500 (Fig. B3).


501 A single pulsation signal is found in this star at a frequency  
 502 of  $0.646584 \pm 0.000008$  mHz ( $55.8649 \pm 0.0007$  d $^{-1}$ ) with an ampli-  
 503 tude of  $0.053 \pm 0.006$  mmag (Fig. 9), which was also noted by  
 504 Mathys et al. (2020b). This mode does not show any amplitude  
 505 or phase variability over the observing window, which would be  
 506 consistent with a rotation period much longer than the observing  
 507 window (Mathys et al. 2020b). The single mode in this star is at  
 508 a frequency commonly associated with the  $\delta$  Sct stars, and would  
 509 make this star the longest period roAp star, after HD 177765 which  
 510 was discovered by Alentiev et al. (2012) with the mode frequency  
 511 later refined to be  $0.702580 \pm 0.000006$  mHz ( $60.7029 \pm 0.0005$  d $^{-1}$ ;  
 512 Holdsworth 2016). With just a single mode, and an Ap stellar classi-  
 513 fication, this star can be classed as an roAp star, however modelling  
 514 of the pulsation is needed to confirm this.



**Figure 8.** Top: phase folded light curve of TIC 310817678, folded on a rotation period of 2.75003 d. Bottom: amplitude spectrum showing the rich pulsation pattern in this star. The vertical dotted lines denote the frequencies of rotationally split multiplet components.



**Figure 10.** Top: phase folded light curve on the assumed rotation period of 4.189 d. The significant scatter is a result of a second significant low frequency. Bottom: amplitude spectrum of TIC 380651050 showing the two pulsation modes in this star.



**Figure 9.** Amplitude spectrum of TIC 356088697 showing the single pulsation mode in this star.

#### 4.1.10 TIC 380651050

TIC 380651050 (HD 176384) has no in-depth studies in the literature. Reiners & Zechmeister (2020) provide a mass of  $1.25 M_{\odot}$  and  $T_{\text{eff}} = 6531$  K, with Paunzen (2015) giving Strömgren-Crawford photometric indices of  $b - y = 0.285$ ,  $m_1 = 0.162$ ,  $c_1 = 0.446$  and  $H\beta = 2.679$ .

TIC 380651050 (HD 176384) was observed in sector 13. There are two harmonic series at low frequency which both give plausible rotation periods; they are:  $4.19 \pm 0.01$  d and  $3.51 \pm 0.02$  d. We take the highest amplitude signal to be the rotation frequency (Fig. 10), but suggest further observations are obtained to corroborate this. One of the two periods may be the result of binarity or contamination by a background star; spectroscopic observations will provide insight to this.

Two pulsation modes are found in this star:  $1.85817 \pm$


$0.00002$  mHz ( $160.546 \pm 0.002$  d $^{-1}$ ) and  $1.87771 \pm 0.00002$  mHz ( $162.234 \pm 0.002$  d $^{-1}$ ) which are separated by  $19.53 \pm 0.03$   $\mu$ Hz (Fig. 10). This separation is plausible half of the large frequency separation, and would imply the that the star is quite evolved, possibly beyond the main-sequence.

#### 4.1.11 TIC 387115314

TIC 387115314 (TYC 9462-347-1) has a spectral classification of A5 (Jackson & Stoy 1954), and has atmospheric parameters  $T_{\text{eff}} = 7765$  K, surface gravity  $\log g = 4.08$  (cm s $^{-2}$ ), and metallicity  $[M/H] = 0.18$  determined by Kordopatis et al. (2013) on the basis of RAdial Velocity Experiment (RAVE) data.

The star was observed in sector 13, from which we derive a rotation period of  $5.264 \pm 0.002$  d (Fig. 11). We find a rich pulsation spectrum for this star at high frequencies. Although we have a well determined rotation period, and significant mean light variations, the modes do not show significant rotationally split sidelobes which implies we see mostly one pulsation pole over the entire rotation period. The pulsation frequencies that we identify are: a singlet at  $1.31902 \pm 0.00004$  mHz ( $113.963 \pm 0.003$  d $^{-1}$ ), two components of a triplet with the mode at  $1.32838 \pm 0.00002$  mHz ( $114.772 \pm 0.002$  d $^{-1}$ ), a triplet centred on  $1.356225 \pm 0.000006$  mHz ( $117.1778 \pm 0.0005$  d $^{-1}$ ) and another triplet centred on  $1.384083 \pm 0.000007$  mHz ( $119.5848 \pm 0.0006$  d $^{-1}$ ). These modes, with their multiplet components, are shown in Fig. 11.

The separation of the three highest modes is  $\sim 27.85$   $\mu$ Hz, with the lowest frequency two modes separated by  $9.36$   $\mu$ Hz. Since we cannot determine the degree of the modes showing multiplets, due to the low amplitude of the obliquely split sidelobes, we are unable to determine if the separation of  $\sim 27.85$   $\mu$ Hz is the large frequency



**Figure 11.** Top: phase folded light curve of TIC 387115314, folded on the rotation period of 5.264 d. Bottom: amplitude spectrum showing the pulsation modes in this star.


separation, or half of it. The separation of  $9.36 \mu\text{Hz}$  is plausible the small frequency separation for this star.

#### 4.1.12 TIC 466260580

TIC 466260580 (TYC 9087-1516-1) is a relatively obscure star with no previous reports of any variability. The RAVE survey (Kunder et al. 2017) obtained  $T_{\text{eff}} = 6780 \pm 170 \text{ K}$ ,  $\log g = 4.03 \pm 0.27 \text{ (cm s}^{-2}\text{)}$  and  $[\text{M}/\text{H}] = -0.7 \pm 0.2$  which are consistent with the TIC values, although the metallicity is strikingly low for an Ap star. These suggest the star is spectral type F3, which is consistent with the Tycho  $B - V = 0.40 \pm 0.07$ . However, the  $T_{\text{eff}}$  is uncertain; for example, McDonald et al. (2017) obtained  $T_{\text{eff}} = 6420 \pm 170 \text{ K}$  from SED fitting, while Gaia Collaboration et al. (2018) give a value around 7000 K.

We have also obtained a spectrum of this star with the HRS on SALT. From this, we obtain  $T_{\text{eff}} = 6800 \pm 200 \text{ K}$ ,  $\log g = 4.0 \pm 0.2 \text{ (cm s}^{-2}\text{)}$  and  $[\text{Fe}/\text{H}] = -0.6 \pm 0.2$ , which are consistent with the RAVE results. We estimate a spectral type of Ap EuCr, and note that rare earth elements are enhanced in the region around  $6150 \text{ \AA}$ , confirming the star to be an Ap star. A full abundance analysis is beyond the scope of this work, but is required to address the low metallicity in this star.

The TESS data, obtained in sector 13, do not show clear signs of rotational variability, but exhibit many low-frequency variations. The pulsation signal is clear in this star, at a frequency of  $1.34006 \pm 0.00002 \text{ mHz}$  ( $115.781 \pm 0.002 \text{ d}^{-1}$ ; Fig. 12).



**Figure 12.** Amplitude spectrum showing the pulsation mode in TIC 466260580.

## 4.2 roAp stars discovered by TESS in the literature

### 4.2.1 TIC 12968953

TIC 12968953 (HD 217704) star was observed in TESS sector 2 and was analysed by Cunha et al. (2019). We find no significant differences between our analysis here and that previously presented. We refer the reader to the aforementioned publication for the details on TIC 12968953. However, we obtained a new spectrum of this target. TIC 12968953 was classified as Ap Sr by Houk (1982); with our SAAO spectrum (Fig. B4), we classify this star as Ap SrEuCr, and determine a temperature of  $T_{\text{eff}} = 7800 \pm 200 \text{ K}$  through comparison with synthetic spectra.

### 4.2.2 TIC 17676722


TIC 17676722 (HD 63773) was observed in sector 7, and reported as an roAp star by Balona et al. (2019). We present a revised rotation period of  $1.5995 \pm 0.0003 \text{ d}$ , which was derived by fitting a 10-component harmonic series to the data. As in Balona et al. (2019) we find four significant peaks which we interpret to be a single quadrupole pulsation mode ( $1.941545 \pm 0.000002 \text{ mHz}$ ;  $167.7495 \pm 0.0002 \text{ d}^{-1}$ ) and 3 sidelobes split by the rotation frequency of the star.

This star has been classified as A2 (Cannon & Pickering 1993), however our SAAO classification spectrum shows the star to be Ap SrEuCr with an effective temperature of  $T_{\text{eff}} = 7800 \pm 200 \text{ K}$  (Fig. B5).

### 4.2.3 TIC 41259805

TIC 41259805 (HD 43226) was reported by both Cunha et al. (2019) and Balona et al. (2019) to be an roAp star, based on 7 sectors of data. There are now 12 sectors of data available for this star from Cycle 1, with no sector 9 data. With this extended time base, we are able to report a more precise rotation period of  $1.714489 \pm 0.000002 \text{ d}$ ; Fig. 13).

There is clear frequency variability in this star, and seen by the ragged nature of the pulsation mode at  $2.311039 \pm 0.000001 \text{ mHz}$  ( $199.6738 \pm 0.0001 \text{ d}^{-1}$ ) in Fig. 13, but not in the mode at  $2.326368 \pm 0.000003 \text{ mHz}$  ( $200.9982 \pm 0.0003 \text{ d}^{-1}$ ). The transient nature of the modes explains the presence of the  $\nu - \nu_{\text{rot}}$  peak found by Cunha et al. (2019), and the lack of its detection by Balona et al. (2019). We note that here the amplitude is about half that reported in Cunha et al. (2019) as a result of the power being spread over a broad frequency range. Since the frequency variability is only seen



**Figure 13.** Top: Phase folded light curve of TIC 41259805, folded on a period of 1.714489 d. Bottom: the two pulsation modes found in this star, with the lower frequency mode showing evidence of frequency variability.


in one mode, it is likely to be intrinsic to the star, rather than caused by binary motion, for example. This type of frequency variability seems common in the roAp stars studied with high-precision data (Holdsworth 2021), with a possible explanation of stochastic perturbations of classical pulsators (Avelino et al. 2020; Cunha et al. 2020), although this is currently untested.

We have obtained a classification spectrum of this star, since the literature class of A0 SrEu implies a hot star which is uncommon amongst the roAp stars. With our spectrum (Fig. B6) we derive a spectral class of Ap SrEu(Cr), with Balmer and metal lines indicating a A6p star.

#### 4.2.4 TIC 49818005

TIC 49818005 (HD 19687), was given the spectral type A9 IV/V in the Michigan Catalog (Houk & Swift 1999b). Conversely, Balona et al. (2019) list a spectral type of A3, which is inconsistent with the 6829-K temperature they cite (from Stevens et al. 2017). There is no information on its peculiarity, if any, but Balona et al. (2019) found a significant high frequency at 1.6424 mHz ( $141.9 \text{ d}^{-1}$ ), suggesting this could be an roAp star. We obtained a new low-resolution classification spectrum to clarify this point (Fig. B7). We find the star to be an Fp SrEu(Cr), with Balmer and metal lines indicating an F1p star, and confirm the presence of rare earth element lines in a high-resolution SALT spectrum. With the high-resolution spectrum, we determine  $T_{\text{eff}} = 7100 \pm 100 \text{ K}$ , which is consistent with the TIC value and previous determinations.

The TESS observations of this star were collected in sector 4, which were analysed by Balona et al. (2019). We present a slightly different value for the pulsation frequency in this star,  $1.641910 \pm 0.000008 \text{ mHz}$  ( $141.8619 \pm 0.0007 \text{ d}^{-1}$ ), as shown in Fig. 14. We suspect this difference is a result of the data treatment at low-frequency, which is not discussed in Balona et al. (2019).



**Figure 14.** The pulsation mode in TIC 49818005.

#### 4.2.5 TIC 152808505

TIC 152808505 (HD 216641) star was observed in TESS sector 1 and was analysed by Cunha et al. (2019). We find no significant differences between our analysis here and that previously presented, so we refer the reader to the aforementioned publication for the details on this star, but provide the derived rotation period and pulsation frequencies in Table 1.

Since the only classification for this star is F3 IV/V (Houk 1978), we obtained a new spectrum of this target with the SpUpNIC instrument at SAAO (Fig. B8). We classify this star as Fp EuCr, and determine a temperature of  $T_{\text{eff}} = 6900 \pm 300 \text{ K}$  through comparison with synthetic spectra. This confirms this star to be an roAp star.

#### 4.2.6 TIC 156886111

TIC 156886111 (HD 47284) was observed in sectors 6 and 7, and originally reported as an roAp star by Balona et al. (2019). The spectral type of this star is commonly given in the literature as A5 SiEuCr (Houk 1978). However, we have obtained a new spectrum of this star (Fig. B9) which shows this star to be an Ap SrEuCr star (showing Sr rather than Si).


We precisely determine the rotation period of this star to be  $6.8580 \pm 0.0003 \text{ d}$  through analysis of the mean light variations, which is inline with the literature value (Watson et al. 2006). After the rotation signal, the dominant mode in this star is at  $0.063251 \pm 0.000002 \text{ mHz}$  ( $5.4649 \pm 0.0002 \text{ d}^{-1}$ ). This may be another example of a  $\delta$  Sct-roAp hybrid, as discussed for TIC 158637987.

We find no more pulsation modes in the data than were presented by Balona et al. (2019); we find a singlet at  $1.27457 \pm 0.00002 \text{ mHz}$  ( $110.123 \pm 0.002 \text{ d}^{-1}$ ), a doublet which is presumed to be a dipole triplet with a missing central component at  $1.30274 \pm 0.00001 \text{ mHz}$  ( $112.557 \pm 0.001 \text{ d}^{-1}$ ), and a quadrupole quintuplet centred on a frequency of  $1.317816 \pm 0.000008 \text{ mHz}$  ( $113.8593 \pm 0.0007 \text{ d}^{-1}$ ).


#### 4.2.7 TIC 259587315

TIC 259587315 (HD 30849) was classified as an Ap SrEuCr by Houk (1978), and has  $T_{\text{eff}}$  measures of 7720 K in the TIC, 7250 K (Glagolevskij 1994), 8187 K (North et al. 1997) and 8000 K (Netopil et al. 2017).

The TESS data for this star, collected in sectors 4 and 5, have been analysed independently by Balona et al. (2019) and Sikora et al. (2019c) who both announced this star to be an roAp star, but



**Figure 15.** Top: phase folded light curve of TIC 259587315, folded on a period of 15.776 d. Bottom: the pulsation modes in TIC 259587315. The vertical dotted lines show the components of obliquely split multiplets, with the dashed vertical line is an independent mode.



**Figure 16.** Top: phase folded SAP light curve of TIC 349945078, folded on a period of 13.4256 d. The discontinuities, and the dip, are instrumental artefacts. Bottom: the pulsation modes in TIC 349945078. The vertical dotted lines show the components of obliquely split multiplets.

provided conflicting rotational periods. Previous searches for high-frequency variability by (Martinez & Kurtz 1994b) and Hubrig et al. (2000) were unsuccessful.

Here, we use the SAP data from sector 4 and the PDC\_SAP data from section 5 to determine a rotation period of  $15.776 \pm 0.005$  d. This is similar to the first reported period in the literature ( $15.865 \pm 0.005$  d; Hensberge et al. 1981) and that reported by Balona et al. (2019), but is twice the period reported by Sikora et al. (2019c), who have detected the first harmonic of the true period.

The star shows several pulsation modes: a triplet with a missing  $+v_{\text{rot}}$  sidelobe, a closely spaced singlet (denoted by the dashed vertical line in Fig. 15), followed by two quintuplets. There appears to be further low amplitude modes between the multiplet groups, but at low S/N. The separation between the triplet and the singlet is  $1.41 \mu\text{Hz}$ . The spacing between the two quintuplets is  $\approx 19.7 \mu\text{Hz}$  which is plausibly the large frequency separation for this star, indicating it is very evolved.

#### 4.2.8 TIC 349945078

TIC 349945078 (HD 57040) was classified as A2 EuCr by Renson & Manfroid (2009). The TIC reports  $T_{\text{eff}} = 7203$  K, though literature values for this parameter range from as low as 6518 K (Schofield et al. 2019) to as high as 8412 K (Ammons et al. 2006). Elkin et al. (2012) reported the presence of a 7.5 kG magnetic field on the star. Martinez & Kurtz (1994b) were unable to detect oscillations from ground-based observations, so it was not identified as an roAp star until Balona et al. (2019) used TESS data to detect two oscillation frequencies.

This star was observed in sectors 6-9, thus we revisit this star to provide a more detailed analysis of the variability with the availability of more data. To extract the rotation signal, we used the

SAP data since the rotation signature in the PDC\_SAP data has been significantly altered. This was evident upon visual inspection. We fitted a four element harmonic series to derive a rotation period of  $13.4256 \pm 0.0008$  d (Fig. 16). This is different from that presented by Balona et al. (2019) by a factor of 2. We confirm our rotation period through the analysis of the oblique pulsations below.


For the pulsation analysis, we used the PDC\_SAP data and prewhitened the light curve to remove the rotation signal and instrumental artefacts. As a final correction, we removed outlying points. The resulting amplitude spectrum shows two groups of peaks composed of a quintuplet and a triplet (Fig. 16). The two modes in this star are separated by  $\approx 60 \mu\text{Hz}$  which is plausibly the large frequency separation.

#### 4.2.9 TIC 350146296

TIC 350146296 (HD 63087) was classified as A7IV by Houk & Cowley (1975). However, due to the clearly spotted nature of this star, and the pulsation variability, we obtained several spectra with the SpUpNIC instrument to find this star to be an F0p EuCr star (Fig. B10).

TIC 350146296 was first reported as an roAp star by Cunha et al. (2019), with Balona et al. (2019) providing an analysis of extra sectors of data. There are now 13 sectors for this star from Cycle 1 which we analyse here. With the extended time base (357 d) we derive a rotation period of  $2.66387 \pm 0.00002$  d. The phased light curve, shown in Fig. 17, reveals a complex spot structure on this star.

There are five pulsation modes in TIC 350146296, all of which have been split into a multiplet by oblique pulsation. The multiplet structures allow us to conclude that the modes are alternating dipole and quadrupole modes, with three dipole modes and two quadrupole




**Figure 17.** Top: phase folded light curve of TIC 350146296, folded on a period of 2.66387 d. Bottom: the rich pattern of pulsation modes in TIC 350146296. The vertical dotted lines show the components of obliquely split multiplets.

modes. All frequencies in this star show signs of variability, making precise frequency extraction difficult. All the modes are split by  $\approx 40 \mu\text{Hz}$  which, given the multiplet structures, we understand to be half the large frequency separation. Shi et al. (submitted) have conducted an extensive study on this star, to which we refer the reader for further details.


#### 4.2.10 TIC 431380369

TIC 431380369 (HD 20880) was identified as a chemically peculiar star by Houk & Cowley (1975) who gave the classification of Ap Sr(EuCr). It was later reported by Cunha et al. (2019) to be a new roAp star from the analysis of the TESS sector 2 data, after Joshi et al. (2016) presented a null result in the search for pulsations. The star was subsequently observed in sectors 6 and 13, where the significant gaps serve to produce a complicated window function. With the extended data, we are able to derive a more precise rotation period for this star of  $5.19716 \pm 0.00005$  d.

This star shows a complex pulsation spectrum, as can be seen in Fig. 18, which is not helped by the significant gaps in the data. The mode at  $0.805252 \pm 0.000002$  mHz ( $69.5737 \pm 0.0002$  d $^{-1}$ ) is significantly broad, suggesting frequency and/or amplitude variability. The mode at  $0.860412 \pm 0.000001$  mHz ( $74.3396 \pm 0.0001$  d $^{-1}$ ) also shows amplitude variability, but includes a rotationally split sidelobe at  $\nu + \nu_{\text{rot}}$ . The four significant modes are listed in Table 1. We extract two extra modes, and the sidelobe, in addition to those reported by Cunha et al. (2019). This is likely a combination of the longer data set where the noise is slightly lower, and the mode lifetime of the pulsations seen in this star. A detailed study, on a rotation period-by-rotation period basis, of this star is required to extract the full information from the TESS data.



**Figure 18.** Pulsation modes in TIC 431380369. The window function gives fine structure to the peaks. However, there is also significant amplitude and frequency variability in this star, as can be seen in the mode at 0.860412 mHz. The vertical dotted lines denote the extracted modes, apart from the  $\nu - \nu_{\text{rot}}$  sidelobe of the highest amplitude peak which is lost in the noise.



**Figure 19.** Amplitude spectrum of TIC 6118924.

### 4.3 Known roAp stars prior to TESS launch

In this section, we present an analysis of the roAp stars which were known prior to the launch of the TESS mission. As with the new roAp star discoveries, we present only new information on stars which have already been discussed in the literature.

#### 4.3.1 TIC 6118924

TIC 6118924 (HD 116114) was observed in sector 10. The star was classified as F0Vp SrCrEu by Abt et al. (1979) and was identified as an roAp star by Elkin et al. (2005a) who discovered variability in the Eu II lines with a period around 21-min ( $0.790$  mHz;  $68.26$  d $^{-1}$ ). A search for photometric variability after the spectroscopic detection by Lorenz et al. (2005) returned a null result at an amplitude level of about 0.5 mmag in Johnson *B*.

The rotation period of TIC 6118924 was determined to be 27.6 d by Landstreet & Mathys (2000) which is longer than the TESS data span. We see no evidence of this period in the light curve. We detect one significant frequency at  $0.76923 \pm 0.00004$  mHz ( $66.462 \pm 0.003$  d $^{-1}$ ) which is consistent with that found in the spectroscopic study (Fig. 19). This is the first photometric detection of the pulsation in this star.

4.3.2 *TIC 33601621*

TIC 33601621 (HD 42659) was classified as Ap SrCrEu by Houk & Smith-Moore (1988). There are several occurrences of the stellar parameters in the literature with  $T_{\text{eff}}$  ranging between 7450–7940 K (Kochukhov & Bagnulo 2006; Balona et al. 2019). Kochukhov & Bagnulo (2006) also provides  $\log L/L_{\odot} = 1.48 \pm 0.09$  and  $M = 2.1 \pm 0.10 M_{\odot}$ . There are also several measures of  $\langle B_z \rangle$  which provide an average of  $0.40 \pm 0.67$  kG (Kochukhov & Bagnulo 2006; Hubrig et al. 2006; Bagnulo et al. 2015).

This star was discovered to be an roAp star by Martinez et al. (1993b) with later observations providing more detailed results by Martinez & Kurtz (1994a). The TESS sector 6 observations have been previously analysed by Balona et al. (2019), with a more detailed analysis presented by Holdsworth et al. (2019), analysing the dipole triplet. We do not present a re-analysis of the same data here, but refer the reader to the previous literature. It is worthy of note, however, that TIC 33601621 is the only roAp star known to be a member of a spectroscopic binary (SB1) system (Hartmann & Hatzes 2015; Holdsworth et al. 2019).


4.3.3 *TIC 35905913*

TIC 35905913 (HD 132205) was classified as an Ap EuSrCr star by Houk & Cowley (1975) and later discovered by Kochukhov et al. (2013) to be an roAp star through time-resolved spectroscopic analysis. Their data revealed a pulsation at  $201.68 \pm 0.59$  d<sup>-1</sup> with an amplitude less than  $100$  m s<sup>-1</sup>. The star had previously been monitored photometrically for pulsations many times (Martinez & Kurtz 1994b), but no signal was detected which is unsurprising given the small amplitude in the spectroscopic data.

TESS observed TIC 35905913 during sectors 11 and 12. The PDC\_SAP light curves for these two sectors differ significantly. A clear rotation signature can be seen the sector 12 data, whereas the sector 11 data have a greatly reduced amplitude attributed to this variability. On inspection, the contamination factor for this star is high, at 0.28, with the contaminating star about 18 arcsec away. According to Gaia DR2 Gaia Collaboration et al. (2018), the *G*-band magnitude difference is 3.25 mag which suggests, under the assumption of negligible extinction differences, that the contaminant star is a mid G-type star.

By fitting a harmonic series of 5 components to the SAP sector 11 and PDC\_SAP sector 12 data simultaneously, we derive a rotation period of  $7.513 \pm 0.001$  d (derived from  $\nu_1$  in Fig. 20). After the removal of this harmonic series, there are clear signs of variability remaining in the data at low frequency ( $\nu_2$  and  $\nu_3$  in the figure). The source of variability is uncertain given the proximity of the contaminating source. In fitting and removing a harmonic series to  $\nu_2$ , power remains around these frequencies in the amplitude spectrum, indicating an unstable frequency/amplitude which is common with star spots that change in size. This leads us to conclude that the contaminant is the source of the  $\nu_2$  harmonic series. The origin of  $\nu_3$  required further investigation, but could be a g mode in the Ap or contaminant star.

We detect no pulsation in the roAp range to a limit of about  $25$   $\mu$ mag in the combined PDC\_SAP flux data. This is unsurprising given the broad red bandpass of the TESS data, coupled with the low pulsation amplitude in spectroscopy.




**Figure 20.** Top: rotation curve for TIC 35905913. There is clear variability on a shorter time scale. Middle: low-frequency amplitude spectrum showing the rotation frequency of the Ap star ( $\nu_1$ ) and its harmonic.  $\nu_2$  and its harmonic are also indicated. Bottom: low-frequency amplitude spectrum after removing the Ap star rotation signature. Further harmonics of  $\nu_2$  are evident, with an additional mode,  $\nu_3$ , and its harmonic labelled.

4.3.4 *TIC 44827786*


TIC 44827786 (HD 150562) was classified as F5Vp SrCrEu by Buscombe & Foster (1995) and discovered to be an roAp star by Martinez & Kurtz (1992) with a period of 10.75 min.

The TIC gives  $T_{\text{eff}} = 7350$  K, which is consistent with that obtained using *uvbyH $\beta$*  photometry from Joshi et al. (2016) and the Moon & Dworetzky (1985) calibration. However, several literature sources give a significantly lower value: 6390 K (Gaia Collaboration et al. 2018), 6620 K (McDonald et al. 2017) and 6820 K (Ammons et al. 2006). The star appears to be slowly rotating, with Elkin et al. (2008) obtaining a  $\nu \sin i = 1.5 \pm 0.5$  km s<sup>-1</sup> and Mathys et al. (2020b) finding no obvious signature of rotational variation from TESS data. The mean magnetic field modulus was found to be 4.8 kG (Kudryavtsev & Romanyuk 2003), while more recently Bagnulo et al. (2015) obtained a value of  $\sim 2$  kG.

We detect two significant modes of pulsation in the TESS sector 12 data (Fig. 21). The highest amplitude mode at a frequency of  $1.54700 \pm 0.00002$  mHz ( $133.661 \pm 0.002$  d<sup>-1</sup>) is separated from



**Figure 21.** The pulsation signatures seen in TIC 44827786. This is the first detection of the highest amplitude mode, whilst the low amplitude mode is reported in the literature.



**Figure 22.** The pulsation signatures seen in TIC 49332521. Modes marked with dotted lines are separated by multiples of  $\approx 26 \mu\text{Hz}$  while those with dashed lines do not fit this pattern.

the previously known pulsation mode ( $1.55040 \pm 0.00005 \text{ mHz}$ ;  $133.955 \pm 0.004 \text{ d}^{-1}$ ) by  $3.40 \pm 0.05 \mu\text{Hz}$  which is plausibly the small frequency separation.

#### 4.3.5 TIC 49332521

TIC 49332521 (HD 119027) is classified on SIMBAD as an  $\alpha^2 \text{ CVn}$  star, but [Martinez et al. \(1998a\)](#) found no rotational signal, and suggested a period of greater than 6 months if there is any rotational variation. [Mathys et al. \(2020b\)](#) also found no rotational variation. Its spectral type is Ap SrEu(Cr) ([Houk 1982](#)). It has been known to be an roAp star for thirty years ([Martinez & Kurtz 1991b](#)), and has a rich p-mode oscillation spectrum ([Martinez et al. 1998b](#)). Magnetic field properties have been measured for HD 119027 ([Mathys 2017](#)), with measured values for  $B_z$ ,  $B_0$  and  $Q_0$  of 0.5, 3.2, and 3.7 kG, respectively. [Mathys \(2017\)](#) provided a  $\langle B \rangle$  value of 3.17 kG.

TESS observed HD 119027 in sector 11, with multiple oscillation frequencies discussed by [Mathys et al. \(2020b\)](#). Here we detect 7 independent modes in TIC 49332521. Six of these modes are almost the same as presented by [Martinez et al. \(1998b\)](#): their five roughly equally split modes (separated by  $\approx 26 \mu\text{Hz}$ ) plus their  $\nu_{2p}$  mode. Our additional mode is found at a lower frequency ( $1.57070 \pm 0.00005 \text{ mHz}$ ;  $135.709 \pm 0.004 \text{ d}^{-1}$ ) which is separated from the next mode by  $264 \mu\text{Hz}$ . This means that modes spanning about 5 radial orders are not excited to a detectable amplitude in this star, which although is unusual, has been seen in other roAp stars, an excellent example of this being TIC 139191168 (HD 217522; Section 4.3.10).

We show in Fig. 22 the amplitude spectrum and the identified modes. Those marked by vertical dotted lines are separated by multiples of  $\approx 26 \mu\text{Hz}$  while those with dashed lines do not fit this pattern. We suspect, given the stellar parameters, that this separation is half of the large frequency separation, such that  $\Delta\nu \approx 52 \mu\text{Hz}$  for TIC 49332521. The separation between the two highest amplitude modes ( $1.78 \mu\text{Hz}$ ) could represent the small frequency separation in this star, although this value is different from the value of  $1.95 \mu\text{Hz}$  proposed by [Martinez et al. \(1998b\)](#).

#### 4.3.6 TIC 69855370

TIC 69855370 (HD 213637) was observed in sector 2. The star is classified as A (pEuSrCr) by [Houk & Smith-Moore \(1988\)](#) and was identified as an roAp star by [Martinez et al. \(1998c\)](#) who reported the discovery of two pulsation modes with periods around 11–12 min,

while acknowledging that the frequency values identified needed to be confirmed due to the presence of  $1 \text{ d}^{-1}$  aliases in the data. Based on the TESS data, [Cunha et al. \(2019\)](#) confirmed the presence of two pulsation modes in the same period range, the first corresponding to one of the modes identified by [Martinez et al. \(1998c\)](#) and the other differing from their second mode by  $1 \text{ d}^{-1}$ .

No sign of rotation modulation was found by [Cunha et al. \(2019\)](#) in their analysis of the TESS data for this star. Nevertheless, based on the measurement of a projected rotational velocity of  $\nu \sin i = 3.5 \pm 0.5 \text{ km s}^{-1}$ , [Kochukhov \(2003\)](#) suggested a rotation period of 25 d. The lack of a rotational modulation in the TESS data could result from an unfavourable alignment ( $i$  or  $\beta$  close to zero), as discussed by [Cunha et al. \(2019\)](#), or be a result of a long rotation period ([Mathys et al. 2020b](#)).

Since the TESS sector 2 data have been analysed in detail, we refer the reader to [Cunha et al. \(2019\)](#) for further discussion, but include the pulsation frequencies in Table 1 for completeness.


#### 4.3.7 TIC 93522454

TIC 93522454 (HD 143487) was observed in sector 12. The star is classified as A3 SrEuCr in [Renson & Manfroid \(2009\)](#). It was identified as an roAp star by [Elkin et al. \(2010\)](#), following the analysis of high time resolution spectra obtained with UVES from which a low amplitude pulsation with a period around 10 min was inferred. The roAp nature of this star was later confirmed by [Kochukhov et al. \(2013\)](#) based on the analysis of a larger set of UVES spectra which allowed the detection of the pulsations in the lines of rare-earth elements as well as in the core of  $\text{H}\alpha$  and the determination of a pulsation period of 9.63 min ( $1.730 \text{ mHz}$ ;  $149.47 \text{ d}^{-1}$ ) with amplitudes up to  $110 \text{ m s}^{-1}$ . No rotation period is reported in the literature for this star.


The TESS data do not show a clear pulsation signal. We show, in Fig. 23, the amplitude spectrum in the frequency region where pulsations have been previously found. We indicate the possible location of the pulsation in this star with an arrow, but caution its presence. The lack of an obvious detection is not surprising given the low amplitude radial velocity variations reported.

#### 4.3.8 TIC 125297016

TIC 125297016 (HD 69013) was observed in sector 7. The star is classified as Ap EuSr by [Houk & Smith-Moore \(1988\)](#). It was identified as an roAp star with a pulsation period around 11 min by [Elkin](#)



**Figure 23.** The amplitude spectrum of TIC 93522454 in the frequency region of interest. We see no clear indication of pulsation, but mark with an arrow a possible signature. The smooth solid curve represents the shape of the peak used by Kochukhov et al. (2013) to provide a pulsation frequency of 1.73 mHz ( $149.47 \text{ d}^{-1}$ ), demonstrating the broad frequency range where the mode may be found.




**Figure 24.** The amplitude spectrum of TIC 125297016 in the frequency region of interest. The highest amplitude peak is at a frequency of 1.47519 mHz which is similar to that reported in the literature, however we are not confident in the detection.

et al. (2011), following the analysis of high time resolution spectra obtained with UVES. Pulsational variability with a similar period was also detected in photometric data published in the same work. The roAp nature of this star was confirmed by Kochukhov et al. (2013) based on the analysis of a larger set of UVES spectra which allowed the detection of the pulsations in the lines of rare-earth elements and the determination of a pulsation period of 11.22 min ( $1.4854 \text{ mHz}$ ;  $128.34 \text{ d}^{-1}$ ). No rotation period is reported in the literature and no sign of rotation variability was found by Mathys et al. (2020b) in their analysis of the TESS data for this star.

The TESS data of this star do not clearly show the known pulsation frequency. A significant peak is detected at a frequency of  $1.47519 \pm 0.00005 \text{ mHz}$  ( $127.456 \pm 0.004 \text{ d}^{-1}$ ; Fig. 24) which is similar to the spectroscopic literature value, but further TESS observations are required to reduce the noise in the data to allow a confident photometric detection of the pulsation.

#### 4.3.9 TIC 136842396

TIC 136842396 (HD 9289) is classified as A(p) SrEuCr by Houk & Swift (1999b) and its mean magnetic field modulus was determined to be  $\langle B \rangle = 2.0 \text{ kG}$  (Ryabchikova et al. 2007). The star was discovered to be an roAp star by Kurtz et al. (1994b) who identified 3 pul-



**Figure 25.** The pulsation signatures seen in TIC 136842396. The vertical dotted lines represent the extracted peaks, all of which are, in their groups, separated by integer values of the rotation frequency.


sation modes in the amplitude spectrum of photometric data. Later, Gruberbauer et al. (2011) observed TIC 136842396 with the MOST satellite and found a greater number of frequencies, but with few in agreement with the first study. TESS observed TIC 136842396 during sector 3, with Balona et al. (2019) presenting the first analysis of those data.

Here, we derive a rotation period of the star from the SAP data since the pipeline has clearly altered the astrophysical signal. We measure a rotation period of  $8.660 \pm 0.006 \text{ d}$ , which is different from that provided by Balona et al. (2019). The pulsation spectrum for this star poses questions due to the presence, and lack thereof, of many rotationally split sidelobes. We extract 9 significant peaks from the prewhitened light curve of this star, as shown in Fig. 25. Given the lack of complete multiplets, we investigated the frequency separation between the assumed mode frequencies and found a recurrence of a separation of  $\approx 31 \mu\text{Hz}$ . Under the assumption that the highest amplitude mode is a quadrupole mode with missing  $\pm \nu_{\text{rot}}$  sidelobes, the pulsation spectrum consists of: a quadrupole mode showing only the  $\nu_{\text{rot}}$  sidelobes, a dipole doublet split by the rotation frequency (with the pulsation mode being the highest amplitude peak), the quadrupole mode that is missing the  $\pm \nu_{\text{rot}}$  sidelobes, and a final quadrupole mode showing only the  $\nu_{\text{rot}}$  sidelobes. These tentative mode identifications imply a large frequency separation of  $\approx 62 \mu\text{Hz}$ .


#### 4.3.10 TIC 139191168

TIC 139191168 (HD 217522) was observed in sector 1. The star is classified as A5 SrEuCr in Renson & Manfroid (2009) and has an upper limit on  $\langle B \rangle$  of  $1.5 \text{ kG}$  (Ryabchikova et al. 2008). It was first identified as an roAp star with pulsations periods around 13.72 min ( $1.2151 \text{ mHz}$ ;  $104.98 \text{ d}^{-1}$ ) by Kurtz (1983), based on photometric data. The authors noted a significant night-to-night pulsation amplitude variability which was confirmed by a multi-site photometric campaign conducted by Kreidl et al. (1991) in 1989. In addition to the pulsational variability around the period detected earlier, the latter data revealed a new pulsation with a period around 8.26 min ( $2.0174 \text{ mHz}$ ;  $174.23 \text{ d}^{-1}$ ). In 2008, further photometric data, as well as high time resolution UVES/ESO spectra were collected. The analysis of these data, published by Medupe et al. (2015), showed that pulsational variability around both periods was still present.

The Cycle 1 TESS data for this star were analysed by Cunha et al. (2019) who detected four modes with frequencies in the interval  $1.2008 - 1.2152 \text{ mHz}$ , thus in the frequency region where all



**Figure 26.** The pulsation signatures seen in TIC 146715928. The vertical dotted lines represent the extracted peaks, which are separated by  $40 \mu\text{Hz}$ .



**Figure 27.** The pulsation signatures seen in TIC 167695608. There is significant frequency variability in most modes.

1021 earlier data sets exhibited pulsations. However, the higher frequency  
 1022 mode, detected in the 1989 and 2008 data sets, was not detected in  
 1023 the TESS data. Similar to the earlier ground-based data, the TESS  
 1024 data showed evidence for amplitude and/or frequency modulation.  
 1025 No rotation period is reported in the literature for this star with  
 1026 Mathys et al. (2020b) classifying the star as a super slowly rotat-  
 1027 ing Ap (ssrAp) star, although the longitudinal field measurements  
 1028 reported in the catalogue by Bagnulo et al. (2015) do show some  
 1029 indication of variability on the time scale of a few years. No sign  
 1030 of rotation variability was found in any of the published analyses of  
 1031 the TESS data (Cunha et al. 2019; David-Uraz et al. 2019; Mathys  
 1032 et al. 2020b).

1033 Since the TESS data for this star have already been analysed in  
 1034 detail, we refer the reader to those works, but include the pulsation  
 1035 frequencies in Table 1 for completeness.

#### 1036 4.3.11 TIC 146715928

1037 TIC 146715928 (HD 92499) was classified as A2p SrEuCr (Houk  
 1038 1978) with  $T_{\text{eff}}$  values ranging from 7200 K to 7810 K (Hubrig &  
 1039 Nesvacil 2007; Freyhammer et al. 2008b; Elkin et al. 2010), and  
 1040  $\log g = 4.1 \pm 0.2$  ( $\text{cm s}^{-2}$ ). Elkin et al. (2010) also provided a full  
 1041 chemical abundance analysis of this star. A mean magnetic field  
 1042 modulus of 8.5 kG was derived from FEROS spectra by Hubrig &  
 1043 Nesvacil (2007) which was corroborated by Mathys (2017) who also  
 1044 suggested the rotation period of TIC 146715928 must be greater than  
 1045 5 yr. TIC 146715928 has a  $v \sin i \approx 3.5 \text{ km s}^{-1}$  (Hubrig & Nesvacil  
 1046 2007; Elkin et al. 2010).

1047 Rapid oscillations were found in the spectra of TIC 146715928  
 1048 by Elkin et al. (2010) with a period of  $10.4 \pm 0.3$  min. TESS observed  
 1049 the star during sectors 9 and 10. Mathys et al. (2020b) used the data  
 1050 to classify the star as an ssrAp star from the lack of rotation signal.  
 1051 They also provided a brief analysis of the two pulsation modes in  
 1052 this star that are separated by  $40 \mu\text{Hz}$ , which could be the large  
 1053 frequency separation, or half of it. We present the extracted mode  
 1054 frequencies in Table 1.

#### 1055 4.3.12 TIC 167695608

1056 TIC 167695608 (TYC 8912-1407-1) was observed in 12 of the 13  
 1057 sectors of Cycle 1 (all but sector 5). The star was classified as  
 1058 F0p SrEu(Cr) by Holdsworth et al. (2014a), who collected spectra  
 1059 to confirm its peculiar nature following the detection of rapid vari-  
 1060 ability, with a period around 11 min ( $1.532 \text{ mHz}$ ;  $132.38 \text{ d}^{-1}$ ) in

a survey conducted using the SuperWASP archive Pollacco et al. (2006).

The first two sectors of Cycle 1 TESS data were analysed by Cunha et al. (2019) who confirmed the pulsation detected previously and identified two more modes at slightly higher frequencies. A follow-up analysis of TESS data collected up to sector 7 revealed yet another pulsation mode at slightly lower frequency than the modes detected previously (Balona et al. 2019). Altogether, the TESS data up to sector 7 had revealed four pulsation frequencies. No sign of rotation variability was found for this star in the SuperWASP (Holdsworth et al. 2014a) or in the year-long TESS Cycle 1 data (Mathys et al. 2020b).


With the longer data set, we are now able to detect 6 pulsation frequencies in this star (Fig. 27), most of which show significant frequency variability, evidenced by broad ragged peaks in the amplitude spectrum that are not cleanly extracted when prewhitened. There is no clear pattern to the splitting of the pulsation modes, thus inhibiting the determination of the large or small frequency separation in this star.

#### 1080 4.3.13 TIC 168383678

1081 TIC 168383678 (HD 96237) is an roAp star discovered by  
 1082 Kochukhov et al. (2013) who determined the following stel-  
 1083 lar parameters:  $T_{\text{eff}} = 7800 \text{ K}$ ,  $\log g = 4.3$  ( $\text{cm s}^{-2}$ ),  $v \sin i =$   
 1084  $6 \text{ km s}^{-1}$  and a mean magnetic field modulus of  $\langle B \rangle = 2.9 \text{ kG}$ .  
 1085 TIC 168383678 has a spectral classification of A4 SrEuCr (Renson  
 1086 & Manfroid 2009) and a rotation period of 20.91 d (Freyhammer  
 1087 et al. 2008b). The dominant pulsation mode was reported at 1.2 mHz  
 1088 (Elkin et al. 2011; Kochukhov et al. 2013).

1089 TESS observed the star during sector 9. The SPOC pipeline  
 1090 has removed the astrophysical signal associated with the rotation  
 1091 period in this star, so we use the SAP data. Upon inspection of the  
 1092 light curve, it is not clear if we have observed one rotation cycle or  
 1093 half of it. Inspection of the two observed light minima show subtly  
 1094 different shapes (Fig. 28). The data for this star span just 24.2 d  
 1095 which is just a little longer than the rotation period reported in the  
 1096 literature. From the light curve, we estimate a rotation period of  
 1097 either  $\sim 21$  d or  $\sim 42$  d.

1098 We detect a significant peak in the amplitude of the SAP  
 1099 light curve at a frequency of  $1.19610 \pm 0.00003 \text{ mHz}$  ( $103.343 \pm$   
 1100  $0.003 \text{ d}^{-1}$ ) which has a  $S/N$  of 7.6. There are, perhaps, further  
 1101 modes in this star at slightly higher frequencies than that detected,  
 1102 but more data are required to confirm their presence.



**Figure 28.** Top: the SAP light curve of TIC 168383678. It appears that one cycle is completed, but inspection of the unequal light minima suggests we may be seeing only half of a rotation cycle. Bottom: the amplitude spectrum of TIC 168383678 with the known pulsation mode indicated. There may be further modes at higher frequency, but with low significance in this data set.


#### 4.3.14 TIC 170419024

TIC 170419024 (HD 151860) was classified as Ap SrEu(Cr) (Houk & Cowley 1975). Kochukhov et al. (2013) measured a mean magnetic field modulus of 2.5 kG, and detected rapid pulsations through the analysis of high-resolution spectra collected with UVES, after the null detection in photometry (Martinez & Kurtz 1994b). An average pulsation period of  $12.30 \pm 0.09$  min was established from the radial velocity of lines of rare earth elements. Some lower amplitude peaks at other frequencies were detected indicating this star to be a multiperiodic roAp star.


The TESS sector 12 data show this star to be an ssrAp star (Mathys et al. 2020b) since the data show no rotation signature. Mathys et al. (2020b) also commented on the pulsation modes in this star, as detected in the TESS data. There is a significant peak at  $0.85014 \pm 0.00002$  mHz ( $73.452 \pm 0.002$  d<sup>-1</sup>) which was not reported by Kochukhov et al. (2013), and a second peak at  $1.36330 \pm 0.00004$  mHz ( $117.789 \pm 0.003$  d<sup>-1</sup>; Fig. 29) which is consistent with the spectroscopic analysis.

#### 4.3.15 TIC 173372645

TIC 173372645 (HD 154708), with a rotation period of 5.37 d, an effective temperature of 7200 K (Joshi et al. 2016),  $\log g = 4.11$  (cm s<sup>-2</sup>), and  $v \sin i = 4.0$  km s<sup>-1</sup>, is classified as an ApEuSrCr star (Renson & Manfroid 2009). The dominant pulsation mode has a frequency of 2.088 mHz (180.4 d<sup>-1</sup>), which was discovered by Kurtz et al. (2006b) through the analysis of high time resolution UVES spectra at an amplitude of just  $\approx 60$  m s<sup>-1</sup>. TIC 173372645 has a magnetic field of 24.5 kG, the strongest ever



**Figure 29.** The pulsation signatures seen in TIC 170419024. The vertical dotted lines represent the extracted peaks.



**Figure 30.** The amplitude spectrum of TIC 173372645. The arrow indicated a potential peak, whereas the broad smooth curve represents the shape of the peak used by Kurtz et al. (2006b) to identify the pulsation mode in this star.


measured for an roAp star (Hubrig et al. 2005), and one of the strongest for an Ap star.

TESS observed TIC 173372645 in sector 12. The light curve is rotationally modulated with a period of  $5.363 \pm 0.001$  d. There is no clear indication of pulsation in this star in the TESS data, only a few peaks in the amplitude spectrum in the frequency region where the pulsation was previously found (Fig. 30; Kurtz et al. 2006b). The low amplitude in the spectroscopic data seems to have translated to an undetectable peak in the TESS photometry.


#### 4.3.16 TIC 189996908

TIC 189996908 (HD 75445) was classified as an Ap SrEu(Cr) star by Houk (1982) and has a magnetic field strength of  $\langle B \rangle = 2985 \pm 42$  G (Mathys et al. 1997). The stellar parameters were measured to be  $T_{\text{eff}} = 7700$  K,  $\log g = 4.3$  (cm s<sup>-2</sup>) (Ryabchikova et al. 2004),  $v \sin i \leq 2$  km s<sup>-1</sup> (Kochukhov et al. 2009a) and the Hipparcos parallax resulted in  $\log L = 1.17 \pm 0.06$  L<sub>⊙</sub> (Kochukhov & Bagnulo 2006). Based on radial velocity measurements, Kochukhov et al. (2009a) discovered rapid oscillations in this star, with a main pulsation frequency at 1.85 mHz (159.8 d<sup>-1</sup>).

TESS observed TIC 189996908 in sectors 8 and 9. The data show no rotational modulation which implies a long rotation period, or an unfavourable view. The magnetic field measurements presented by Mathys (2017) show changes over short time periods, implying a low inclination angle, rather than slow rotation. The pulsation signal is weak in this star, and would perhaps be classed



**Figure 31.** The amplitude spectrum of TIC 189996908. The low S/N peak detection agrees with that found in spectroscopic observations.



**Figure 32.** The single pulsation mode seen in TIC 268751602.

as noise if it was not previously known. We find a peak in the amplitude spectrum at  $1.84163 \pm 0.00002$  mHz ( $159.117 \pm 0.002$  d<sup>-1</sup>) with a S/N of 5.3 (Fig. 31). This is another example of how low amplitude modes in spectroscopic observations are difficult to observe photometrically.

#### 4.3.17 TIC 211404370

TIC 211404370 (HD 203932) is classified as A5 SrEu in the catalogue of [Renson & Manfroid \(2009\)](#). [Gelbmann et al. \(1997\)](#) derived  $T_{\text{eff}} = 7540 \pm 100$  K and  $\log g = 4.1$  (cm s<sup>-2</sup>) for this star, while the average magnetic field,  $\langle B \rangle$ , was determined to be 0.25 kG ([Bychkov et al. 2009](#)) and the  $v \sin i$  was measured to be 4.7 km s<sup>-1</sup> ([Smalley et al. 2015](#)). Pulsations were first detected in TIC 211404370 by [Kurtz \(1984\)](#) and was later shown to have 4 significant peaks in the amplitude spectrum between 1.280 and 2.737 mHz ([Martinez et al. 1990](#)).

This star was observed in TESS's sector 1, with the data analysed in [Cunha et al. \(2019\)](#). We find no significant difference in our analysis and that previously presented, so we refer the reader to that work. We include the rotation frequency, and the frequency of the two pulsation modes, in Table 1.

#### 4.3.18 TIC 237336864

TIC 237336864 (HD 218495), with a spectral type of ApEuSr ([Renson & Manfroid 2009](#)), was shown to have a  $v \sin i$  of  $16$  km s<sup>-1</sup> ([Smalley et al. 2015](#)) with an average magnetic field,  $\langle B \rangle$ , of 0.77 kG ([Bychkov et al. 2009](#)). [Cunha et al. \(2019\)](#) derived  $T_{\text{eff}} = 7950 \pm 160$  K and  $\log g = 4.1$  (cm s<sup>-2</sup>). Rapid oscillations were first detected by [Martinez & Kurtz \(1990\)](#) in this star.

This star was observed in sector 1, with the data analysed in [Cunha et al. \(2019\)](#). The data show a complex pulsation pattern with, in increasing frequency, a singlet, a doublet which is split by twice the rotation frequency, a quintuplet with components split by the rotation frequency, and a triplet of frequencies with each component split by twice the rotation frequency. Given the variety of different multiplet structures, a simple understanding if this star is not possible, but to develop a complete understanding is beyond the scope of this work. We provide the inferred pulsation frequencies and amplitudes in Table 1.

#### 4.3.19 TIC 268751602

[Houk & Smith-Moore \(1988\)](#) classified TIC 268751602

(HD 12932) as ApSrEuCr, with [Kochukhov et al. \(2007\)](#) providing parameters of  $T_{\text{eff}} = 7620$  K,  $\log g = 4.15$  (cm s<sup>-2</sup>), and  $v \sin i = 3.5$  km s<sup>-1</sup>, and [Ryabchikova et al. \(2007\)](#) deriving  $\langle B \rangle = 1.7$  kG.


[Schneider & Weiss \(1990\)](#) and [Kreidl \(1990\)](#) independently collected high time resolution photometric data of TIC 268751602 and discovered variability with a period of about 11.6 min (1.4368 mHz; 124.14 d<sup>-1</sup>), thus announcing TIC 268751602 as an roAp star. This period was later refined by [Schneider et al. \(1992\)](#) to be 11.7 min, with the authors suggesting a rotation period of 3.5295 d. Analysis of photometric data by [Martinez et al. \(1994\)](#) showed that TIC 268751602 was a single-mode pulsator with a period of 11.6 min. The earlier identification the rotation period was concluded to be likely incorrect. [Kochukhov et al. \(2007\)](#) identified variability in spectral lines of some elements in the atmosphere of TIC 268751602.

TESS observed this star during sector 3, with an analysis of those data presented by [Balona et al. \(2019\)](#). We confirm the lack of detection of rotation in this star, and agree the with the pulsation frequency presented in that work, namely  $1.436302 \pm 0.000006$  mHz ( $124.0965 \pm 0.0005$  d<sup>-1</sup>). There is only one pulsation mode seen in these data of this star (Fig 32).

#### 4.3.20 TIC 279485093

TIC 279485093 (HD 24712, HR 1217) was classified as A9VpSrEuCr by [Abt & Morrell \(1995\)](#). [Perraut et al. \(2016\)](#) derived a temperature from bolometric flux and optical interferometry of  $7235 \pm 280$  K, while [Sikora et al. \(2019a\)](#) derived  $7150 \pm 170$  K from spectroscopy. The mean magnetic field modulus for TIC 279485093 was determined to be 2.3 kG ([Ryabchikova et al. 2008](#); [Rusomarov et al. 2013](#)). [Kurtz & Marang \(1987b\)](#) obtained a rotation period from the mean light variation of 12.45733 d, and [Kobzar et al. \(2020\)](#) found a signal in the Fourier spectrum of the TESS data that may correspond to a rotation period of  $12.44 \pm 0.02$  d. [Kurtz \(1981\)](#) first discovered 6.15 min (2.71 mHz;  $234.1$  d<sup>-1</sup>) oscillations in this star, and [Balona et al. \(2019\)](#) list 10 independent frequencies. Many of these modes show the expected rotation-modulated amplitude variations, but unexplained variations in this modulation also occur on shorter timescales of days ([White et al. 2011](#)). This star is one of the first discovered and best studied roAp stars, appearing extensively in the literature.

As mentioned, TESS observed this star in sector 5. From the SAP data, we derive a rotation period of  $12.578 \pm 0.008$  d which is different from previous literature values (Fig. 33). The sector 5 SAP data, after removing points obviously affected by instrumental




**Figure 33.** Top: the phase folded SAP light curve of TIC 279485093, folded on a period of 12.578 d. Middle: the amplitude spectrum of the PDC\_SAP data. The vertical dotted lines represent the frequencies extracted. Bottom: amplitude spectrum of the residuals after removing 24 frequencies (9 pulsation modes and significant sidelobes). There is power remaining around the dominant modes which suggests significant frequency variability in this star. It seems likely there are further modes close to the noise level, too.

artefacts, span 25.63 d – a little over twice the derived rotation period, thus this value should be treated with caution.

To inspect the pulsations in this star, we used the PDC\_SAP light curve. The pulsation spectrum is known to be rich in this star, as evidenced by the middle panel in Fig. 33. We are able to extract 9 pulsation modes (as listed in Table 1) and 15 sidelobes that are split from the pulsation frequency by oblique pulsation. In comparison to the best ground based photometric data set for this star (Kurtz et al. 2005), we detect all but their  $\nu_6$  frequency although we see a power excess in the bottom panel of Fig. 33 around this frequency. We detect two further modes (our lowest frequencies) which likely correspond to two modes detected by Mkrtichian & Hatzes (2005) via high-resolution spectroscopy.

#### 4.3.21 TIC 280198016

TIC 280198016 (HD 83368, HR 3831) is a close visual binary, with the roAp star having spectral type A8V SrCrEu and the companion



**Figure 34.** Amplitude spectrum of PDC\_SAP light curve of TIC 280198016. We detect 6 of the 7 septuplet components found by Kurtz et al. (1997a).


having type F9V (Corbally 1984). Sikora et al. (2019a) derive  $T_{\text{eff}} = 7660 \pm 170$  K which is consistent with the TIC value. TIC 280198016 shows a polarity reversing magnetic field strength that varies about zero mean with amplitude  $737 \pm 68$  G (Mathys 1991), and has a mean quadratic magnetic field strength of 11 kG (Mathys 1995). The rotation period of TIC 280198016 was precisely determined to be  $2.851976 \pm 0.000003$  d (Kurtz et al. 1997a), with the star showing a single dipole pulsation mode at  $123.38 \text{ d}^{-1}$  that was exactly split into a septuplet by pulsation amplitude modulation with the rotation frequency. Baldry et al. (1998b) found that the radial velocity variations were amplitude and phase modulated in the same manner as the photometric variations, first confirming the oblique pulsator model and ruling out the spotted pulsator model for roAp stars.

The TESS sector 9 data of TIC 280198016 allow us to derive a rotation frequency (Table 1) which agrees with that of Kurtz et al. (1997a). We detect 6 significant peaks (Fig. 34) of the reported septuplet in TIC 280198016 and find that the mode shows frequency variability, as reported by (Kurtz et al. 1997a). We also detect the second harmonic of the pulsation mode which forms a triplet centred on twice the pulsation frequency, although Kurtz et al. (1997a) show four harmonics.


#### 4.3.22 TIC 315098995

TIC 315098995 (HD 84041) was classified as Ap SrEuCr by Houk (1982), and has effective temperature measurements of 8100 K (Mathys et al. 1996), 7800 K and 7500 K from Elkin et al. (2008) obtained through the photometric calibrations of Moon & Dworetzky (1985) and fitting synthetic spectra to the  $H_{\alpha}$  profile. TIC 315098995 is a magnetic star with Kochukhov & Bagnulo (2006) reporting the first detection of a longitudinal magnetic field of  $497 \pm 91$  G while Bychkov et al. (2009) gave, for the averaged quadratic magnetic field, a value of  $479 \pm 72$  G. The latest evaluation given in the FORS1 catalogue of stellar magnetic field measurements was  $640 \pm 52$  G measured from Balmer and metal lines (Bagnulo et al. 2015).

Martinez (1991) first reported the star as an roAp star with a pulsation period of about 14.6 min ( $1.14 \text{ mHz}$ ;  $98.6 \text{ d}^{-1}$ ). Later Martinez et al. (1993a) clarified that the star pulsated in at least three modes with periods near 15 min and with frequency separations of about  $30 \mu\text{Hz}$ . However, it was noted that these frequencies were not a complete description of the pulsation behaviour of TIC 315098995. It exhibited mean light variations of a double-wave nature, indicating that both magnetic poles were vis-



**Figure 35.** Amplitude spectrum of the two-sector light curve of TIC 315098995. There are at least two very variable peaks in this star. The vertical dashed lines represent the positions of the frequencies found by [Martinez et al. \(1993a\)](#).



**Figure 36.** Amplitude spectrum of the light curve of TIC 322732889. The vertical dotted lines represent the septuplet components split by oblique pulsation.

ible. The rotation period determined from these observations was  $3.69 \pm 0.01$  d. [Elkin et al. \(2008\)](#) reported maximum pulsation amplitude of  $0.5 \text{ km s}^{-1}$  and a pulsation amplitude from the  $H_{\alpha}$  core of  $0.20 \text{ km s}^{-1}$ .

TESS observed TIC 315098995 during sectors 8 and 9. We combined the PDC\_SAP data from sector 8 and the SAP sector 9 data to determine the rotation period since the SPOC pipeline has distorted the astrophysical signal in the latter. We measure a rotation period of  $3.6884 \pm 0.0004$  d, which is consistent with the literature value.

The TESS data show this star to have a very variable pulsation signal, as reported by previous studies. We find two peaks (Fig. 35) that are split by twice the rotation frequency. We assume then, that we are seeing two sidelobes of a dipole triplet where the peak corresponding to the actual pulsation mode frequency is missing, implying that either  $i$  or  $\beta$  are close to  $90^{\circ}$ . This is corroborated when considering the results of [Martinez et al. \(1993a\)](#). We do not detect, in the combined light curve, the other modes reported to be present in this star. A careful inspection of the data, in shorter sections, is required to extract the full information on this star.

#### 4.3.23 TIC 322732889

TIC 322732889 (HD 99563) was classified as A9 III star by [Houk \(1978\)](#) and with a Sr peculiarity added later ([Bidelman 1981](#)), although the star appears in the catalogue of [Renson & Manfroid \(2009\)](#) as an F0 Sr. From high-resolution spectra, [Elkin et al. \(2008\)](#) found  $T_{\text{eff}} = 7700$  K and  $\log g = 4.2$  ( $\text{cm s}^{-2}$ ). The first measure of the longitudinal magnetic field gave  $688 \pm 145$  G ([Hubrig et al. 2004](#)), with a further two measurements at different phases of  $-235 \pm 73$  G and  $670 \pm 37$  G ([Hubrig et al. 2006](#)). These detections were confirmed by [Kochukhov & Bagnulo \(2006\)](#) who measured a field strength of  $-392$  G.

[Dorokhova & Dorokhov \(1998\)](#) first announced the detection of pulsation in TIC 322732889 with a period of about 11.17 min ( $1.4921$  mHz;  $128.9 \text{ d}^{-1}$ ). Later, [Handler & Paunzen \(1999\)](#) confirmed the roAp nature of the star with a period of 10.7 min. [Elkin et al. \(2005b\)](#) discovered remarkably large amplitude pulsations in the star for some spectral lines with amplitudes up to  $5 \text{ km s}^{-1}$  making this star the largest radial velocity amplitude roAp star known. A more detailed study by [Handler et al. \(2006\)](#) using multi-site photometry found a single rotationally modulated mode with a frequency  $1.558$  mHz ( $134.6 \text{ d}^{-1}$ ) connected with the rotation de-

termined to be 2.912 d. Further spectroscopic observations found little variation in the pulsation amplitude in TIC 322732889 ([Kurtz et al. 2006a](#)), while [Freyhammer et al. \(2009\)](#) claimed the presence of  $H_{\alpha}$  spots caused by the settling of He in the presence of the magnetic field. Based on MOST observations, [Gruberbauer et al. \(2011\)](#) found a new candidate frequency independent of the primary multiplet.


TESS observed TIC 322732889 during sector 9. With those data, we derive a rotation period of  $2.9114 \pm 0.0002$  d which is consistent with that in the literature. As reported in previous works, we find a septuplet of peaks in the amplitude spectrum separated by the rotation frequency. The central peak, representing the pulsation mode, is at  $1.557657 \pm 0.000002$  mHz ( $134.5816 \pm 0.0002 \text{ d}^{-1}$ ; Fig. 36), and has a smaller amplitude than the first rotational sidelobes. We detect none of the candidate modes of [Gruberbauer et al. \(2011\)](#).

#### 4.3.24 TIC 326185137


TIC 326185137 (HD 6532) was classified as Ap SrCrEu by [Houk \(1982\)](#), although there are a variety of different classifications ([Skiff 2014](#)). Effective temperature measurements for this star range from 7995 K by [Bai et al. \(2019\)](#) to 8382 K by [McDonald et al. \(2017\)](#). The  $\log g$  value for this star is also poorly constrained in the literature, with an average value of 4.25 ( $\text{cm s}^{-2}$ ). Several measurements of the magnetic field have been made, with the  $\langle B \rangle_{\text{rms}}$  being 0.411 kG ([Mathys & Hubrig 1997](#); [Bychkov et al. 2009](#); [Bagnulo et al. 2015](#)).

TIC 326185137 was discovered to be an roAp star by [Kurtz & Kreidl \(1985\)](#) with a period of 6.9 min ( $2.4155$  mHz;  $208.7 \text{ d}^{-1}$ ). They found a frequency triplet split by the rotation frequency of the star. Subsequently, the star was the subject of many ground-based photometric campaigns, with [Kurtz et al. \(1996b\)](#) concluding the star pulsates with a distorted dipole mode. The rotation period of this star has been refined many times in the literature, with an average period of 1.94 d ([Kurtz & Marang 1987a](#); [Kurtz et al. 1996a](#); [Renson & Manfroid 2009](#); [Balona et al. 2019](#)).

TESS observed this star in sector 3, with an initial analysis of the data being presented by [Balona et al. \(2019\)](#) where it was noted that the multiplet structure was vastly different from previous  $B$  photometric data. [Kurtz & Holdsworth \(2020\)](#) explored this further and postulated why there is a striking difference between the two data sets. They suggest two scenarios to explain the difference: either the star has changed its pulsation axis or, more likely, we are



**Figure 37.** Amplitude spectrum of the light curve of TIC 326185137, showing the single pulsation mode split into a multiplet. Note the different structure seen here compared with the photometric  $B$  data of Kurtz et al. (1996b).



**Figure 38.** Amplitude spectrum of the light curve of TIC 340006157, showing the many pulsation modes. The broad peaks indicate significant amplitude variability in this star, as reported in the literature.

1379 seeing a depth dependence on the mode geometry revealed by the 1421  
1380 two wavelengths of observation. 1422

1381 From the data, we determine a rotation period of  $1.9447 \pm$  1423  
1382  $0.0001$  d, and a pulsation frequency of  $2.402163 \pm 0.000007$  mHz 1424  
1383 ( $207.5469 \pm 0.0006$  d $^{-1}$ ). We identify 4 of the 5 peaks of the dis-  
1384 torted dipole multiplet (Fig. 37) in the TESS Cycle 1 data.

#### 1385 4.3.25 TIC 340006157

1386 TIC 340006157 (HD 60435) was classified as A3 SrEu in Renson 1426  
1387 et al. (1991). Ghazaryan et al. (2019) report abundances for the star 1427  
1388 and classify it as neither He-poor or rich. Kochukhov & Bagnulo 1428  
1389 (2006) detected a surface magnetic field with  $B_z = -315 \pm 87$  G. 1429  
1390 The TIC gives  $T_{\text{eff}} = 8427$  K, though values in the literature range 1430  
1391 as low as 7694 K (McDonald et al. 2012). The rotation period was 1431  
1392 reported by Kurtz et al. (1990) to be 7.68 d, with the light curve 1432  
1393 displaying a double wave of unequal depth. 1433

1394 The star was first identified as an roAp star by Kurtz (1984). 1436  
1395 Matthews (1987) conducted a detailed study and identified 17 fre- 1437  
1396 quencies, which were identified as consecutive radial overtones of 1438  
1397  $\ell = 1$  and  $\ell = 2$  modes, with amplitudes which were highly variable 1439  
1398 on timescales of days. Vauclair & Théado (2004) attempted mod- 1440  
1399 elling based on these frequencies, with results that suggested the 1441  
1400 presence of a He gradient inside the star. Balona et al. (2019) per- 1442  
1401 formed a preliminary analysis using TESS and detected 6 independ- 1443  
1402 ent frequencies between 1.3079 and 1.4352 mHz (113 – 124 d $^{-1}$ ). 1444  
1403 All peaks were broad and poorly resolved, and many displayed 1445  
1404 variable amplitudes which both confirmed the results of Matthews 1446  
1405 (1987) and perhaps explained why not all the frequencies reported 1447  
1406 by the earlier author were not detected in TESS data. 1448

1407 This star was observed in 7 sectors of Cycle 1, but with two 1449  
1408 2-sector gaps. To derive the rotation period here, we used a combi- 1450  
1409 nation of the raw and pipeline corrected data since the pipeline has 1451  
1410 altered the astrophysical signal in sectors 7 and 10 for this star, which 1452  
1411 in total span 296 d. We derive a rotation period of  $7.6797 \pm 0.0001$  d. 1453

1412 The TESS amplitude spectrum is no different for this star as 1454  
1413 reported in the literature: it shows significant frequency/amplitude 1455  
1414 variability, making precise frequency extraction difficult. We are, 1456  
1415 however, able to identify 8 pulsation frequencies in the data, which 1457  
1416 we have listed in Table 1. Some of the extracted frequencies are the 1458  
1417 same as those reported by Matthews (1987), but we do not detect 1459  
1418 the lowest frequency modes reported in that work. Only the second 1460  
1419 highest frequency mode shows rotationally split sidelobes in the 1460  
1420 amplitude spectrum. Due to the broad nature of the signal in

the amplitude spectrum (Fig. 38) we have inflated our uncertain-  
ties to reflect the uncertainty in the frequency determination. This  
star requires a detailed individual study in the light of significant  
amplitude variations and short mode lifetimes.

#### 1425 4.3.26 TIC 348717688


1426 TIC 348717688 (HD 19918) was observed in sectors 1,12 and 13, 1427  
1428 and has a spectral type of A5 SrEuCr (Renson & Manfroid 2009). 1429  
1429 Its pulsations were first detected by Martinez & Kurtz (1991a) and 1430  
1430 revisited later by Martinez et al. (1995) using ground-based photom- 1431  
1431 etry from SAAO. In addition to the dominant pulsation frequency 1432  
1432 at 1.510 mHz (130.5 d $^{-1}$ ) and its harmonics, a second frequency at 1433  
1433 1.4806 mHz (127.9 d $^{-1}$ ) was detected. TIC 348717688 has a known 1434  
1434 mean longitudinal magnetic field strength of  $-625 \pm 87$  G (Hubrig 1435  
1435 et al. 2006), although a rotation period is not currently known. 1436  
1436 More recently, Cunha et al. (2019) analysed TESS sector 1 data of 1437  
1437 TIC 348717688 and extracted eight significant pulsation frequen- 1438  
1438 cies, including the dominant frequency originally reported by Mar- 1439  
1439 tinez & Kurtz (1991a). Mathys et al. (2020b), using the complete 1440  
1440 TESS Cycle 1 data set, classified this star as an ssrAp star through 1441  
1441 the lack of detection of a rotation signal, and very sharp spectral 1442  
1442 lines. 1443

1444 Despite the large gap in the Cycle 1 data set, the window func- 1445  
1445 tion of the data does not significantly interfere with the frequency 1446  
1446 analysis of this star. However, we identify frequency variability in 1447  
1447 this star. We are able to extract seven pulsation frequencies from the 1448  
1448 Cycle 1 light curve (Fig. 39), and note the presence of the first har- 1449  
1449 monic of the highest amplitude mode. The frequency not detected 1450  
1450 here, but quoted in Cunha et al. (2019), is likely the result of the 1451  
1451 frequency variability of a nearby mode as this mode is now split 1452  
1452 into several peaks in the amplitude spectrum. 1453


#### 1451 4.3.27 TIC 363716787

1452 TIC 363716787 (HD 161459) was observed in sector 13, and has a 1453  
1453 spectral type of Ap EuSrCr (Houk 1978). It has a dominant pulsation 1454  
1454 frequency of 1.39 mHz (120.1 d $^{-1}$ ; Martinez & Kauffmann 1990; 1455  
1455 Martinez et al. 1991), but no magnetic field or rotation period has 1456  
1456 been measured yet. 1457

1458 The TESS data allow us to provide a rotation period measure- 1459  
1459 ment of  $5.9660 \pm 0.0014$  d. The folded light curve, Fig. 40, shows a 1460  
1460 double wave, implying that we see both magnetic poles, under the 1460  
1460 assumption that the chemical spots form at or near the magnetic



**Figure 39.** Amplitude spectrum of the light curve of TIC 348717688, showing the multiple pulsation modes. All modes show signs of frequency variability.




**Figure 40.** Top: light curve of TIC 363716787 phased on its rotation period of 5.9660 d. Bottom: amplitude spectrum of the light curve of TIC 363716787. The dashed lines represents the two extracted modes, while the dotted lines show the multiplet components.

poles. The pulsation spectrum of this star, as noted by [Martinez et al. \(1991\)](#), shows amplitude variability of its modes. We detect two independent modes in this star, with one showing rotationally split sidelobes. It is unclear which of the multiplet peaks is the pulsation mode, but we follow [Martinez et al. \(1991\)](#), taking the mode to be at 1.3909 mHz (120.1725 d<sup>-1</sup>). This mode is separated from the next by  $\approx 31 \mu\text{Hz}$  which is plausibly half the large frequency separation in this star.

#### 4.3.28 TIC 368866492

TIC 368866492 (HD 166473) was classified as A5p SrCrEu in [Renson & Manfroid \(2009\)](#). The TESS Asteroseismic Target List (ATL; [Schofield et al. 2019](#)) reports  $T_{\text{eff}} = 6811 \text{ K}$ , but this is the low



**Figure 41.** Amplitude spectrum of the light curve of TIC 368866492 showing the three equally split modes.

end of the range of  $T_{\text{eff}}$  from the literature, which range as high as 8500 K ([Ammons et al. 2006](#)). [Ryabchikova \(2005\)](#) and [Ghazaryan et al. \(2019\)](#) conducted chemical analyses and found nothing peculiar. [Mathys et al. \(2020b\)](#) used magnetic field modulus measurements to determine the rotation period to be  $3836 \pm 30 \text{ d}$ , using high-resolution spectroscopy spanning the period 1992 – 2019, and confirming an earlier report of 3514 d by [Shavrina et al. \(2014\)](#). Recently, [Mathys et al. \(2020a\)](#) reported a mean magnetic field modulus of 7.13 kG for this star.


[Kurtz & Martinez \(1987\)](#) first detected photometric oscillations in TIC 368866492, finding periods between 8.8 and 9.1 min (1.8315–1.8939 mHz; 158.2–163.6 d<sup>-1</sup>), while [Kurtz et al. \(2003\)](#) and [Mathys et al. \(2007\)](#) confirmed the presence of oscillations using radial velocity measurements. [Mathys et al. \(2007\)](#) reported three frequencies, 1.833, 1.886, and 1.928 mHz, with amplitudes up to  $110 \text{ ms}^{-1}$ , but all with low signal-to-noise. They also suggested that their data hinted at periodic variability of the mean magnetic field modulus, with an amplitude of  $21 \pm 5 \text{ G}$ .

TESS observed TIC 368866492 in sector 13. As discussed by [Mathys et al. \(2020b\)](#), there is no sign of rotational variability in the light curve, only low-frequency noise which is suspected of arising from contamination. We detect three main pulsation frequencies in this star, but note there is power excess in the amplitude spectrum after they have been removed. This leads us to speculate there are further low-amplitude modes that are not fully resolved in the TESS data.


The three peaks that we do see ([Fig. 41](#)) have equal phases and are equally split by  $3.69 \mu\text{Hz}$  which would correspond to a rotation frequency of 3.14 d if the multiplet is caused by oblique pulsation. However, from the literature, we know this star to have an extremely long rotation period. A different interpretation of this splitting is either the large or small frequency separation. With a value of  $3.69 \mu\text{Hz}$  this is too small to be the large frequency separation, but is plausibly the small separation. However, in that situation, one would require modes of  $\ell = 1, 3, 5$  or  $\ell = 0, 2, 4$  to be visible and would need an explanation for why they would be equally split. This star needs careful consideration and modelling to solve these conundrums, which we leave to a future study.

#### 4.3.29 TIC 369845536

TIC 369845536 (2MASS J19400781-4420093) is a well-studied, high-amplitude roAp star with a distorted quadrupole mode ([Holdsworth et al. 2018a](#)). It was originally discovered to have high-frequency variability by [Holdsworth et al. \(2014a\)](#) using Su-



**Figure 42.** Amplitude spectrum of the light curve of TIC 369845536 showing the quadrupole mode with its equally split quintuplet of peaks.



**Figure 43.** Amplitude spectrum of the light curve of TIC 394272819 showing the multiple modes in this star. We mark the presumed modes with dashed lines, with rotationally split sidelobes marked by dotted lines.

perWASP photometry, and a dominant pulsation frequency of 2.042 mHz ( $176.4 \text{ d}^{-1}$ ). Spectroscopy obtained by Holdsworth et al. (2018a) led them to classify its spectral type as A7Vp Eu(Cr). The ground-based photometry also allowed the rotation period to be determined as  $9.5344 \pm 0.0012 \text{ d}$ , and modelling of the amplitude and phase modulation of its quadrupole pulsation mode confirmed its distorted nature.

TESS observed TIC 369845536 during sector 13. The data span 26 d which makes the determination of the rotation period less certain in the TESS data, at  $9.50 \pm 0.02 \text{ d}$  over the previous measurement. The pulsation signature is strikingly large in this star; it is the roAp star with the highest amplitude mode seen in  $B$  photometric observations, which is also the case here with the TESS observations (Fig. 42). Beyond the known pulsation mode in this star, we detect no further modes. The multiplet structure of the  $\ell = 2$  mode is similar to that seen in  $B$  data (Holdsworth et al. 2018a)

#### 4.3.30 TIC 394124612

TIC 394124612 (HD 218994) was observed in sector 1, and previously analysed by Cunha et al. (2019). It has a spectral type of A3 Sr (Renson & Manfroid 2009), but is also known to be a visual binary with a separation of 1.2 arcsec (Renson et al. 1991). Pulsations consistent with a  $\delta$  Sct star and a longitudinal magnetic field strength of  $440 \pm 23 \text{ G}$  were detected by Kurtz et al. (2008). Despite attempts to detect roAp pulsations, none have so far been detected in photometry (see e.g. Martínez & Kurtz 1994b). However, analysis of spectroscopic time series data has discovered variability in its Nd III and Pr III lines (González et al. 2008), which showed a frequency of 1.17 mHz ( $101.1 \text{ d}^{-1}$ ).

Since an analysis of the TESS data has already been presented by Cunha et al. (2019), we refer the reader to that work while providing the rotation period and pulsation frequency in Table 1.

#### 4.3.31 TIC 394272819

TIC 394272819 (HD 115226) is a well-known  $\alpha^2$  CVn variable star. It was classified as Ap Sr(Eu) by Houk & Cowley (1975). The  $T_{\text{eff}}$  values in the literature for this star are mostly in agreement, with a value around 7640 K (Kochukhov & Bagnulo 2006; Smalley et al. 2015). Using this value and the Hipparcos parallax, Kochukhov & Bagnulo (2006) determine  $L = 7.2 \pm 1.9 L_{\odot}$ . Later, Kochukhov et al. (2008) determined the projected rotational velocity  $v_e \sin i = 25\text{--}30 \text{ km s}^{-1}$  and conducted an abundance analysis, finding nearly

solar abundances of Mg, Si, Ti, and Fe, an overabundance of Cr, Co, and rare earth elements, and an under-abundance of Ba. Bagnulo et al. (2015) reported a  $\langle B_z \rangle_{\text{rms}}$  value of 0.828 kG.

The roAp nature of TIC 394272819 was discovered by Kochukhov et al. (2008). Using time-series high-resolution HARPS spectroscopy, they measured the high-amplitude 10.87-min ( $1.5333 \text{ mHz}$ ;  $132.5 \text{ d}^{-1}$ ) oscillation in this star from radial velocity variations. Pulsational variability was discovered in the cores of hydrogen lines and in doubly-ionized Pr, Nd, and Dy lines. Recently, the same 102 HARPS spectra were analysed by Binnenfeld et al. (2020) by using the unit-sphere representation periodogram (USuRPER), finding the same period.

Photometric ASAS-3 data of TIC 394272819 were analysed by Hümmerich et al. (2016). They detected a clear signal with a period of 2.9882 d. In addition they checked the Hipparcos data and found marginal variability with  $P = 3.61 \text{ d}$ , which is not present in the ASAS-3 data. Joshi et al. (2016) confirmed the presence of 10.86-min oscillations analysing high-speed photometry from the Nainital-Cape Survey.

TIC 394272819 was observed during sectors 11 and 12. The combined PDC\_SAP data allow us to derive a rotation period of  $2.98827 \pm 0.00008 \text{ d}$  which is consistent with the ASAS-3 data. There is no evidence of the 3.61-d signal found in the Hipparcos data. We detect four pulsation modes in this star. In increasing frequency we see a singlet, three components of a quintuplet, a triplet and another singlet. Fig. 43 shows the amplitude spectrum of this star. We interpret this pattern as a series of alternating  $\ell = 1$  and  $\ell = 2$  modes with some components lost in the noise. This interpretation is reinforced when considering the spacing between the modes; under our assumption they are all separated by  $29.58 \mu\text{Hz}$  which we understand to be the half of the large frequency separation for this star.

#### 4.3.32 TIC 402546736

TIC 402546736 (HD 128898;  $\alpha$  Cir) has many similar classifications in the literature. Here we quote that of Gray et al. (2006): A7Vp SrCrEu. This star is a well studied, naked eye object, and as such has a plethora of information in the literature. We provide here a very limited introduction to the star, and encourage the interested reader to explore the references provided.

Effective temperature measurements for TIC 402546736 range from 7420 K (e.g., Bruntt et al. 2008; Desidera et al. 2015) to around 8000 K (e.g., Kurtz & Martínez 1993; Matthews et al. 1999)


1599 derived through a variety of different methods.  $\log g$  determina-  
 1600 tions also vary, but are consistently around  $4.1 \text{ (cm s}^{-2}\text{)}$  within the  
 1601 quoted errors (e.g., Kochukhov et al. 2009b; Sikora et al. 2019a),  
 1602 with  $v \sin i$  values consistently around  $13 \text{ km s}^{-1}$  (e.g., Kupka et al.  
 1603 1996; Ammler-von Eiff & Reiners 2012). The abundance pattern in  
 1604 TIC 402546736 is typical for Ap stars (Kupka et al. 1996; Bruntt  
 1605 et al. 2008), with under-abundances of C, N, and O, and overabun-  
 1606 dances of rare-earth and some other heavy elements. Kochukhov  
 1607 et al. (2009b) built a self-consistent empirical model atmosphere of  
 1608 TIC 402546736 and conducted stratification modelling of some el-  
 1609 ements, showing that chemical stratification has an important effect  
 1610 on the model structure of TIC 402546736. A  $\langle B \rangle$  value of  $0.9 \text{ kG}$   
 1611 have been reported for this star (Bruntt et al. 2008; Sikora et al.  
 1612 2019b).

1613 TIC 402546736 was classified as roAp star showing an 6.8-min  
 1614 ( $2.4510 \text{ mHz}$ ;  $211.8 \text{ d}^{-1}$ ) oscillation by Kurtz & Cropper (1981) and  
 1615 subsequently, 47 hours of high-speed, ground-based photometry  
 1616 was analysed by Kurtz et al. (1981). Since its discovery as an roAp  
 1617 star, TIC 402546736 has been frequently analysed. Notable are the  
 1618 results of Kurtz et al. (1994a) who found the star to pulsate in  
 1619 a dipole mode with a frequency of  $2.442 \text{ mHz}$  ( $211.0 \text{ d}^{-1}$ ), and  
 1620 subsequent lower amplitude modes. They also provided a precise  
 1621 rotation period of  $4.4790 \text{ d}$ . Later, Bruntt et al. (2009) used 84 d of  
 1622 WIRE observations to refine the separation of the modes in this star  
 1623 to be  $30.2 \mu\text{Hz}$ . Weiss et al. (2016) and Weiss et al. (2020) used  
 1624 4 BRITE satellites to observe TIC 402546736 for a total of 146 d,  
 1625 providing them data to model the spot configuration of the star, and  
 1626 confirm the previously tentative detection of some low amplitude  
 1627 modes.

1628 Beside the photometric observations of TIC 402546736, there  
 1629 have been substantial spectroscopic studies using high time res-  
 1630 olution spectra. These studies (e.g., Baldry et al. 1998a, 1999;  
 1631 Kochukhov & Ryabchikova 2001; Balona & Laney 2003) have  
 1632 shown that the principal pulsation mode can have an amplitude  
 1633 of several  $\text{km s}^{-1}$ , with some spectral lines pulsating in anti-phase  
 1634 with others, suggesting a node, or false node, in the atmosphere of  
 1635 the star (Sousa & Cunha 2011; Quiral-Manosalva et al. 2018).

1636 TESS observed TIC 402546736 during sectors 11 and 12. We  
 1637 use the SAP data to determine a rotation period of  $4.4812 \pm 0.0005 \text{ d}$   
 1638 which is different from that of Kurtz et al. (1994a) by  $4.5\sigma$ . We also  
 1639 use the SAP data for the pulsation analysis that follows, since the  
 1640 SPOC pipeline has introduced significant noise to this bright star.  
 1641 As expected from the literature, the amplitude spectrum is rich in  
 1642 pulsation modes and rotationally split sidelobes (Fig. 44). We  
 1643 extract 23 significant peaks from the amplitude spectrum, but note  
 1644 there is evidence of further modes seemingly present in the data.  
 1645 Of those 23 peaks, we identify 10 as pulsation modes (see Table 1),  
 1646 one as a harmonic and 2 as peaks due to non-linear interactions.  
 1647 The latter 3 peaks are all found above the Nyquist frequency and  
 1648 are reminiscent of those seen in the K2 data of 33 Lib (HD 137949;  
 1649 Holdsworth et al. 2018c), making TIC 402546736 only the second  
 1650 roAp star to show such non-linear interactions.

1651 Although not all modes are separated by the same frequency,  
 1652 many are separated by  $30.18 \mu\text{Hz}$  which we take to be half of the  
 1653 large frequency separation in this star, as did Bruntt et al. (2009).  
 1654 For those modes where the frequency spacing is not a multiple of  
 1655  $30.18 \mu\text{Hz}$ , we postulate that these mode frequencies have been  
 1656 affected by the magnetic field, as was seen in TIC 279485093  
 1657 (HR 1217; Kurtz et al. 2005).




**Figure 44.** Amplitude spectrum of the SAP light curve of TIC 402546736 ( $\alpha$  Cir) showing the multiple modes in this star. The top panel shows the region where the pulsation modes are detected. We mark the modes with dashed lines, with rotationally split sidelobes marked by dotted lines. The bottom panel shows the harmonic of the dominant mode, and two additional peaks that are the result on non-linear interactions in this star. Note the change in scale between the two panels.

#### 4.3.33 TIC 434449811

1658 TIC 434449811 (HD 80316) was classified as Ap Sr(Eu?) with ‘extre-  
 1659 mly strong Sr II’ by Bonsack (1974). There is no detailed spec-  
 1660 troscopic analysis of the fundamental properties in the literature,  
 1661 so  $T_{\text{eff}}$  values are based on photometry with values ranging from  
 1662  $8110 \text{ K}$  (Gaia Collaboration et al. 2018) to  $9120 \text{ K}$  (Ammons et al.  
 1663 2006). Using  $uvbyH\beta$  from Paunzen (2015), and the calibrations of  
 1664 Moon & Dworetzky (1985), gives  $T_{\text{eff}} = 8420 \text{ K}$ , which is consistent  
 1665 with  $8280 \text{ K}$  obtained from the spectral energy distribution (McDon-  
 1666 ald et al. 2012). The mean longitudinal magnetic field strength has  
 1667 been determined to be  $-183 \pm 38 \text{ G}$  by Hubrig et al. (2006) and  
 1668  $-249 \pm 89 \text{ G}$  by Bagnulo et al. (2015).

1669 The rotation period was determined to be  $2.09 \text{ d}$  (Manfroid  
 1670 & Mathys 1986; Kurtz et al. 1997b), although a recent value of  
 1671  $1.91 \text{ d}$  was given by Oelkers et al. (2018). With  $v \sin i = 32 \pm$   
 1672  $5 \text{ km s}^{-1}$  (Wade 1997) estimated a rotational axis inclination of  
 1673  $i = 60 \pm 15^\circ$ . The radial velocity is given as  $9.5 \pm 1.1 \text{ km s}^{-1}$  by  
 1674 Gontcharov (2006). However, 5 measurements with HARPS show a  
 1675 wide variation of a 2-yr time-span, ranging from  $10.6$  to  $46.7 \text{ km s}^{-1}$   
 1676 (Trifonov et al. 2020), indicating this could be a spectroscopic binary  
 1677 star. The star is part of a very wide ( $> 70 \text{ arcsec}$ ) optical pair with  
 1678 HD 80298 (BD-19 2673; WDS J09184-2022A; Mason et al. 2001),  
 1679 but there is no physical association.

1680 TIC 434449811 is a known roAp star, with a  $7.4\text{-min}$   
 1681 ( $2.2523 \text{ mHz}$ ;  $194.6 \text{ d}^{-1}$ ) oscillation discovered by Kurtz (1990)  
 1682 in photometric observations. Elkin et al. (2008) found a pulsation  
 1683 signal of the same period with an amplitude of  $0.32 \text{ km s}^{-1}$  in spec-  
 1684 tral lines. A later, more in depth photometric study was presented



**Figure 45.** Amplitude spectrum of the light curve of TIC 434449811. The top panel shows the low-frequency modes found in this star, with the bottom panel showing the multiplet around the known pulsation mode.

by Kurtz et al. (1997b) who found a frequency triplet with equal amplitudes in a *B* filter.

TESS observed the star during sector 8. The PDC\_SAP data, after the removal of some outlying points at the start of the second orbit, provide a rotation period for this star of  $2.08862 \pm 0.00004$  d, which is consistent with the literature value. After removal of the rotation signature and its harmonics, there are two peaks remaining in the  $\delta$  Sct frequency range (Fig. 45). It is unclear whether these are mode frequencies in TIC 434449811 or in a companion star. Given the relatively weak magnetic field in this star, it is possible that the  $\delta$  Sct frequencies are in the Ap star (Murphy et al. 2020), but detailing modelling and the exclusion of other possibilities is required.

The roAp pulsation signature in this star has a different multiplet structure than previously seen (Kurtz et al. 1997b); there are, in the TESS data, two peaks with similar amplitudes, and a further 3 peaks with low signal to noise. This change in multiplet structure is similar to that seen in TIC 326185137 (HD 6532; Kurtz & Holdsworth 2020) and may be a result of the different filters probing different depths in the stellar atmosphere. A detailed analysis of this star is required to fully understand its pulsation properties.

#### 4.3.34 TIC 469246567

TIC 469246567 (HD 86181) is listed as an  $\alpha^2$  CVn star on SIMBAD and has a spectral type Ap Sr (Houk & Cowley 1975). Kurtz & Martinez (1994) discovered 6.2-min (2.6882 mHz;  $232.3$  d $^{-1}$ ) oscillations in TIC 469246567, making it one of the most rapidly oscillating Ap stars. A magnetic field strength of  $536 \pm 58$  kG was measured by Bagnulo et al. (2015). Despite the  $\alpha^2$  CVn classification, there is only one recent measure of the rotation period of this star (Shi et al. submitted).

TIC 469246567 was observed by TESS during sectors 9 and

10. These data have been extensively analysed by Shi et al. (submitted) so we refer the reader there to a detailed discussion. In short, the pulsation spectrum consists of alternating dipole, quadrupole, dipole modes, all rotationally split into multiplets. The TESS data also provide the first measure of the rotation period of this star to be  $2.05115 \pm 0.00006$  d. The pulsation frequencies, and rotation period, are included in Table 1 for completeness.

## 5 CANDIDATE ROAP STARS

In this section, we provide information on a number of stars which were flagged by the search teams as possible roAp stars. These stars are classed as candidate roAp stars since either their spectral classification does not indicate chemical peculiarity, their pulsation signature does not show signs of multiplets split by the rotation frequency, or their presumed pulsation signal is close to a signal to noise limit of 4.0.

### 5.1 TIC 1727745

TIC 1727745 (HD 113414) was classified as an F7/8V type star by Houk & Smith-Moore (1988). There have been numerous studies that attempt to calculate physical parameters of the star. Among these, Haakonsen & Rutledge (2009) identified the star as an x-ray source and Suchkov et al. (2003) found its x-ray luminosity to be  $\log L_x = 29.45$  erg s $^{-1}$  in the investigation of x-ray properties of the ROSAT F stars. Suchkov et al. (2003) also estimated the temperature and metallicity to be  $T_{\text{eff}} = 6200$  K and Fe/H = 0.04, respectively.

Using precision radial velocities from HIRES on the Keck I telescope, Butler et al. (2017) stated that the star showed chromospheric emission variations, which led to radial velocity variations, due to either stellar rotation or long-term activity cycles. Additionally,  $(b - y)$ ,  $m_1$ ,  $c_1$  colour indexes and H $\beta$  index of the star were provided to be 0.357, 0.182, 0.386 and 2.624, respectively (Paunzen 2015; Hauck & Mermilliod 1998), which do not imply this star is chemically peculiar.


Observed in sector 10, the TESS data show a rotationally modulated light curve, which has a period of  $3.172 \pm 0.001$  d (Fig. 46). The well defined phased light curve suggests that the surface inhomogeneities are stable over the observations period. The pulsation signal in the star occurs at  $2.29116 \pm 0.00004$  mHz ( $197.956 \pm 0.004$  d $^{-1}$ ) which is in the frequency range of known roAp pulsations.

The contamination factor, in the TIC, for this star is very small (0.0001) which implies all of the light in the photometric aperture is originating from the target star.


### 5.2 TIC 3814749

TIC 3814749 (HD 3748) was also listed in Balona et al. (2019) as an roAp candidate. The star is a known double system (with a contamination factor of 1.431) which have spectral types of A0/1 IV/V and A9/F0V (Houk & Swift 1999b), but with also A5 quoted in the literature. We obtained a classification spectrum from SAAO and confirm the lack of strong chemical peculiarities in this star, suggesting it is not an Ap star (Fig. B11). However, the Ca II K line is weak in the star, as is the Mg II 4481 Å line.

TESS observed TIC 3814749 during sector 3. With those data, we detect a low frequency signal corresponding to a period of  $1.689 \pm 0.002$  d. We also find two peaks at higher frequency which we assume are a result of pulsational variability (Fig. 47). These



**Figure 46.** Top: phase folded light curve of TIC 1727745, phased on a period of 3.172 d. Bottom: the high frequency pulsation mode found in this star.



**Figure 47.** Top: phase folded light curve of TIC 3814749, phased on a period of 1.689 d. Bottom: the two high frequency pulsation modes found in this star, as marked by dotted lines.

1772 two modes are well separated in frequency, by 1.12 mHz, which  
1773 requires a significant range of unexcited overtones as has been seen  
1774 in TIC 139191168 (HD 217522) and now also in TIC 49332521  
1775 (HD 119027).

1776 However, the star is listed as an Algol type binary (Samus'  
1777 et al. 2003). It is possible that the low frequency signal is the orbital  
1778 frequency of the A star and a compact companion. In this case, a  
1779 large gap between the observed frequencies is less uncommon (e.g.,  
1780 Sahoo et al. 2020). High-resolution spectroscopic monitoring of  
1781 this system is needed to solve this problem.

### 1782 5.3 TIC 158637987

1783 There is little information in the literature about TIC 158637987  
1784 (HD 10330). Houk & Cowley (1975) gave the spectral type as  
1785 A9V, indicating a chemically normal star. Paunzen (2015) pro-  
1786 vided Strömgren-Crawford photometric indices of  $b - y = 0.207$ ,  
1787  $m_1 = 0.143$ ,  $c_1 = 0.702$ , and  $H\beta = 2.732$ . Using the calibrations  
1788 of Crawford (1979), these indices suggest the star is significantly  
1789 metal weak.

1790 The TESS observations, collected during sectors 2 and 3, show  
1791 significant variability in the low-frequency range which we take to be  
1792 g mode pulsations (Fig. 48). The high-frequency variability in this  
1793 star occurs at about 2.71 mHz ( $236 \text{ d}^{-1}$ ; Fig. 48), where we identify  
1794 two peaks of marginal significance. Spectroscopic observations are  
1795 required of this star to test the existence of chemical peculiarities  
1796 and/or a magnetic field. If either exist, this star provides another  
1797 example of a  $\gamma$  Dor-roAp hybrid star. If this is a roAp star, it is  
1798 unique in having apparently low metallicity.


1799 The TIC provides a contamination factor of 0.048 for  
1800 TIC 158637987, suggesting there is very little contamination from  
1801 a nearby (on the sky) star.

### 1802 5.4 TIC 324048193


1803 TIC 324048193 (HD 85892) was classified as Ap Si by Houk &  
1804 Cowley (1975). The literature provides  $T_{\text{eff}}$  values of between  
1805 7770 K (Bai et al. 2019) to 10110 K (Anders et al. 2019), but with  
1806 most estimates around 9300 K (McDonald et al. 2012; Chandler  
1807 et al. 2016; Ammons et al. 2006). The TIC estimate is by far the  
1808 highest temperature estimate at 11950 K, and we treat that with cau-  
1809 tion. The presence of Si in the stellar spectrum suggests a late B early  
1810 A star which is consistent with the higher temperature range. The  
1811 mass and radius estimates for the star,  $5.2 M_{\odot}$  and  $4.5 R_{\odot}$  (Kervella  
1812 et al. 2019; McDonald et al. 2017) suggest it has evolved away  
1813 from the zero-age main-sequence. A  $v \sin i$  value of  $45 \text{ km s}^{-1}$  was  
1814 quoted by Renson & Manfroid (2009), with Watson et al. (2006)  
1815 measuring a rotation period of 4.295 d.

1816 TESS observed TIC 324048193 during sectors 11 and 12. The  
1817 light curve shows clear modulation due to spots, as expected for a  
1818 magnetic CP star. From the variations, we derive a rotation period  
1819 of  $4.2953 \pm 0.0001 \text{ d}$  which is consistent with that in the litera-  
1820 ture. The pulsation signal in this star is at a high frequency of  
1821  $3.15347 \pm 0.00002 \text{ mHz}$  ( $272.460 \pm 0.002 \text{ d}^{-1}$ ; Fig. 49). This is  
1822 among the highest frequencies known in the roAp stars. This star is  
1823 also at the upper temperature range of the known roAp stars, after  
1824 TIC 123231021 (KIC 7582608; Holdsworth et al. 2014b), but it is  
1825 well within the theoretical instability strip for this class of pulsator  
1826 (Cunha 2002).

1827 We classify this as a candidate roAp star since the signal is  
1828 detected at marginal significance ( $S/N = 5.4$ ), and lacks signs of ro-  
1829 tationally split sidelobes (which would be lost in the noise). Further  
1830 TESS observations of this star are due, which will hopefully allow  
1831 this star to be confirmed as a new roAp star. Since the contamination  
1832 factor is small (0.0032) we are confident that the signal is from this  
1833 star.



**Figure 48.** Top: g-mode frequencies seen in TIC 158637987. Bottom: the two pulsation modes found in this star, as marked by dotted lines.




**Figure 49.** Suspected pulsation mode in TIC 324048193.

### 5.5 TIC 410163387

TIC 410163387 (HD 76279) was classified as an ApSrEuCr star by (Houk & Smith-Moore 1988), although they note it as a ‘weak case’. There are differing  $T_{\text{eff}}$  values in the literature, with most in the region of 7100 K (e.g., Ammons et al. 2006; Bai et al. 2019). There has been one previous published attempt to find pulsations in this star, however the authors returned a null result at a magnitude limit of 0.8 mmag in 2-hrs of photometric  $B$  observations Nelson & Kreidl (1993).

TESS observed TIC 410163387 in sector 8. An amplitude spectrum of the light curve at low frequency (Fig. 50) reveals the presence of probable g-mode pulsations in this star. At high frequency, we detect a single peak at  $1.43155 \pm 0.00003$  mHz ( $123.686 \pm 0.003$  d $^{-1}$ ) which has a S/N of 7.0. We cannot conclude, with confidence, that this peak is a true pulsation mode, and, coupled with the presence of the g-modes, we rate this star as a candidate roAp star until further observations are made to confirm the existence of the mode. We also note that the TIC contamination



**Figure 50.** Top: suspected g-mode pulsations in TIC 410163387. Bottom: suspected roAp pulsation mode in TIC 410163387.

factor for this star is non-negligible at 0.01, implying that some of the signals may be from a contaminant source.

## 6 CONCLUSIONS

We have presented the results of an unbiased systematic survey of hot ( $> 6000$  K) stars observed during the first Cycle of TESS observations, covering the Southern Ecliptic hemisphere, in 2-min cadence. We selected over 50 700 stars in this temperature range, and searched for pulsational variability with frequencies greater than 0.52 mHz, resulting in over 6713 detections. After the removal of obvious contaminant stars, the final sample consisted of 163 stars for detailed frequency analysis. The main discoveries from this work can be summarised as:

(i) We have reported the discovery of 12 new roAp stars. Six of these stars show multiple pulsation frequencies in their light curves, making them of utmost importance for asteroseismic modelling. In addition to these stars, another (TIC 119327278) shows the presence of a  $\delta$  Sct mode, suggesting this star belongs to the rare  $\delta$  Sct-roAp hybrid stars. This star also shows the shortest rotation period among the class of roAp stars, at 1.085 d. We have identified the roAp star with the lowest frequency pulsation, TIC 356088697, which shows a single pulsation mode at 0.646584 mHz ( $P = 25.8$  min), allowing us to test the lowest end of the radial orders excited in the roAp stars. Of the 12 new roAp stars, we provided the first measures of the rotation period for 5, and find three to show no rotational variability.

(ii) We have presented the results of TESS observations of a further 44 roAp stars that were known in the literature. Of these 44, we find three (TIC 35905913 – HD 132205, TIC 93522454 – HD 143487, and TIC 173372645 – HD 154708) show no clear pulsation signal in the high-precision TESS data and note that these three stars were discovered to have low amplitude pulsations through

the analysis of time-resolved high resolution spectroscopic observations. Furthermore, we present the tentative detection of modes in 2 other roAp stars which were discovered through spectroscopic observations.

(iii) Of the entire number of 56 roAp stars observed, 68 per cent (38) show rotationally modulated light curves, with rotation periods in the range 1.085 – 21 (or 42) d. It is not yet clear if there is a causal link between rotation and pulsation, but these observations will provide new results to test this relationship.

(iv) There are many instances where the rotationally split multiplets show different structures, in the same star, for the same degree mode (e.g., TIC 176516923 – HD 38823, TIC 310817678 – HD 88507, TIC 259587315 – HD 30849). This provides a problem for the application of the oblique pulsator model where the stellar inclination angle,  $i$ , and the magnetic obliquity,  $\beta$ , can be derived or constrained. This, coupled with the difference in structure between observations made in different filters (e.g., TIC 326185137 – HD 6532; Kurtz & Holdsworth 2020), provides a significant challenge in understanding the results of the oblique pulsator model.

(v) We have reported the presence of non-linear interactions in the pulsation spectrum of TIC 402546736 (HD 128898) around the harmonic of the principal mode. This is only the second star (after 33 Lib; Holdsworth et al. 2018c) where this phenomenon is seen.

The results presented here have provided new insights into the roAp class of pulsationally variable stars. We have identified new pulsation frequencies in many stars, and shown frequency variability is a common feature of the roAp stars. These observational results will drive forwards the development of state-of-the-art theoretical models of these stars, providing insight into the physical processes at play in these magnetic, chemically peculiar stars. Puzzling frequency separations, high-frequency modes, and a lack of roAp stars probing the blue edge of the instability strip, mean these results will have a significant impact on the theoretical understanding of the rapidly oscillating Ap stars. We will use these results to conduct a homogeneous study of the properties of roAp stars and explore any correlations between stellar and pulsation properties, and investigate any differences between the observed frequencies and those predicted by models.

## ACKNOWLEDGEMENTS

DLH acknowledges financial support from the Science and Technology Facilities Council (STFC) via grant ST/M000877/1. MSC acknowledges the support by FCT/MCTES through the research grants UIDB/04434/2020, UIDP/04434/2020 and PTDC/FIS-AST/30389/2017, and by FEDER - Fundo Europeu de Desenvolvimento Regional through COMPETE2020 - Programa Operacional Competitividade e Internacionalização (grant: POCI-01-0145-FEDER-030389). MSC is supported by national funds through FCT in the form of a work contract. VA was supported by a research grant (00028173) from VILLUM FONDEN. Funding for the Stellar Astrophysics Centre is provided by The Danish National Research Foundation (Grant agreement no.: DNR106).

This paper includes data collected by the TESS mission, which are publicly available from the Mikulski Archive for Space Telescopes (MAST). Funding for the TESS mission is provided by NASA's Science Mission directorate. Funding for the TESS Asteroseismic Science Operations Centre is provided by the Danish National Research Foundation (Grant agreement no.: DNR106), ESA PRODEX (PEA 4000119301) and Stellar Astrophysics Centre (SAC) at Aarhus University. We thank the TESS team and staff

and TASC/TASOC for their support of the present work. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France.


## REFERENCES

- Abt H. A., Morrell N. I., 1995, *ApJS*, **99**, 135  
 Abt H. A., Brodzik D., Schaefer B., 1979, *PASP*, **91**, 176  
 Alecian G., Stift M. J., 2010, *A&A*, **516**, A53  
 Alentiev D., Kochukhov O., Ryabchikova T., Cunha M., Tsymbal V., Weiss W., 2012, *MNRAS*, **421**, L82  
 Ammler-von Eiff M., Reiners A., 2012, *A&A*, **542**, A116  
 Ammons S. M., Robinson S. E., Strader J., Laughlin G., Fischer D., Wolf A., 2006, *ApJ*, **638**, 1004  
 Anders F., et al., 2019, *A&A*, **628**, A94  
 Avelino P. P., Cunha M. S., Chaplin W. J., 2020, *MNRAS*, **492**, 4477  
 Babcock H. W., 1960, *ApJ*, **132**, 521  
 Bagnulo S., Fossati L., Landstreet J. D., Izzo C., 2015, *A&A*, **583**, A115  
 Bai Y., Liu J., Bai Z., Wang S., Fan D., 2019, *AJ*, **158**, 93  
 Baldry I. K., Bedding T. R., Viskum M., Kjeldsen H., Frandsen S., 1998a, *MNRAS*, **295**, 33  
 Baldry I. K., Kurtz D. W., Bedding T. R., 1998b, *MNRAS*, **300**, L39  
 Baldry I. K., Viskum M., Bedding T. R., Kjeldsen H., Frandsen S., 1999, *MNRAS*, **302**, 381  
 Balmforth N. J., Cunha M. S., Dolez N., Gough D. O., Vauclair S., 2001, *MNRAS*, **323**, 362  
 Balona L. A., Laney C. D., 2003, *MNRAS*, **344**, 242  
 Balona L. A., et al., 2011, *MNRAS*, **413**, 2651  
 Balona L. A., Holdsworth D. L., Cunha M. S., 2019, *MNRAS*, **487**, 2117  
 Bernhard K., Hümmerich S., Paunzen E., 2020, *MNRAS*, **493**, 3293  
 Bidelman W. P., 1981, *AJ*, **86**, 553  
 Bidelman W. P., MacConnell D. J., 1973, *AJ*, **78**, 687  
 Bigot L., Dziembowski W. A., 2002, *A&A*, **391**, 235  
 Bigot L., Kurtz D. W., 2011, *A&A*, **536**, A73  
 Binnenfeld A., Shahaf S., Zucker S., 2020, *A&A*, **642**, A146  
 Blanco-Cuaresma S., 2019, *MNRAS*, **486**, 2075  
 Blanco-Cuaresma S., Soubiran C., Heiter U., Jofré P., 2014, *A&A*, **569**, A111  
 Bonsack W. K., 1974, *PASP*, **86**, 408  
 Bramall D. G., et al., 2010, in *Ground-based and Airborne Instrumentation for Astronomy III*. p. 77354F, doi:10.1117/12.856382  
 Bruntt H., et al., 2008, *MNRAS*, **386**, 2039  
 Bruntt H., et al., 2009, *MNRAS*, **396**, 1189  
 Buckley D. A. H., Swart G. P., Meiring J. G., 2006, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*. p. 62670Z, doi:10.1117/12.673750  
 Buscombe W., Foster B. E., 1995, MK spectral classifications. Twelfth general catalogue, Epoch 2000 including UVB photometry  
 Butler R. P., et al., 2017, *AJ*, **153**, 208  
 Bychkov V. D., Bychkova L. V., Madej J., 2009, *MNRAS*, **394**, 1338  
 Cannon A. J., Pickering E. C., 1993, *VizieR Online Data Catalog*, p. III/135A  
 Chandler C. O., McDonald I., Kane S. R., 2016, *AJ*, **151**, 59  
 Chen P. S., Liu J. Y., Shan H. G., 2017, *AJ*, **153**, 218  
 Chojnowski S. D., et al., 2019, *ApJ*, **873**, L5  
 Corbally C. J., 1984, *ApJS*, **55**, 657  
 Crause L. A., et al., 2014, in *Ground-based and Airborne Instrumentation for Astronomy V*. p. 91476T, doi:10.1117/12.2055635  
 Crause L. A., et al., 2016, in *Proc. SPIE*. p. 990827, doi:10.1117/12.2230818  
 Crawford D. L., 1979, *AJ*, **84**, 1858  
 Cunha M. S., 2001, *MNRAS*, **325**, 373  
 Cunha M. S., 2002, *MNRAS*, **333**, 47  
 Cunha M. S., 2006, *MNRAS*, **365**, 153  
 Cunha M. S., Gough D., 2000, *MNRAS*, **319**, 1020  
 Cunha M. S., Fernandes J. M. M. B., Monteiro M. J. P. F. G., 2003, *MNRAS*, **343**, 831  
 Cunha M. S., Alentiev D., Brandão I. M., Perraut K., 2013, *MNRAS*, **436**, 1639

- 2007 Cunha M. S., et al., 2019, *MNRAS*, **487**, 3523
- 2008 Cunha M. S., Avelino P. P., Chaplin W. J., 2020, *MNRAS*, **499**, 4687
- 2009 David-Uraz A., et al., 2019, *MNRAS*, **487**, 304
- 2010 Deal M., Cunha M. S., Keszthelyi Z., Perraut K., Holdsworth D. L., 2021, arXiv e-prints, p. arXiv:2104.08097
- 2011 Desidera S., et al., 2015, *A&A*, **573**, A126
- 2012 Dorokhova T. N., Dorokhov N. I., 1998, Contributions of the Astronomical Observatory Skalnaté Pleso, **27**, 338
- 2013 Elkin V. G., Riley J. D., Cunha M. S., Kurtz D. W., Mathys G., 2005a, *MNRAS*, **358**, 665
- 2014 Elkin V. G., Kurtz D. W., Mathys G., 2005b, *MNRAS*, **364**, 864
- 2015 Elkin V., Kurtz D. W., Mathys G., 2008, Contributions of the Astronomical Observatory Skalnaté Pleso, **38**, 317
- 2016 Elkin V. G., Kurtz D. W., Mathys G., Freyhammer L. M., 2010, *MNRAS*, **404**, L104
- 2017 Elkin V. G., Kurtz D. W., Worters H. L., Mathys G., Smalley B., van Wyk F., Smith A. M. S., 2011, *MNRAS*, **411**, 978
- 2018 Elkin V. G., Kurtz D. W., Nitschelm C., 2012, *MNRAS*, **420**, 2727
- 2019 Freyhammer L. M., Kurtz D. W., Cunha M. S., Mathys G., Elkin V. G., Riley J. D., 2008a, *MNRAS*, **385**, 1402
- 2020 Freyhammer L. M., Elkin V. G., Kurtz D. W., Mathys G., Martinez P., 2008b, *MNRAS*, **389**, 441
- 2021 Freyhammer L. M., Kurtz D. W., Elkin V. G., Mathys G., Savanov I., Zima W., Shibahashi H., Sekiguchi K., 2009, *MNRAS*, **396**, 325
- 2022 Gaia Collaboration et al., 2018, *A&A*, **616**, A1
- 2023 Gelbmann M., Kupka F., Weiss W. W., Mathys G., 1997, *A&A*, **319**, 630
- 2024 Ghazaryan S., Alecian G., Hakobyan A. A., 2019, *MNRAS*, **487**, 5922
- 2025 Glagolevskij Y. V., 1994, Bulletin of the Special Astrophysics Observatory, **38**, 152
- 2026 Gontcharov G. A., 2006, *Astronomy Letters*, **32**, 759
- 2027 González J. F., Hubrig S., Kurtz D. W., Elkin V., Savanov I., 2008, *MNRAS*, **384**, 1140
- 2028 Gray R. O., Corbally C. J., 1994, *AJ*, **107**, 742
- 2029 Gray R. O., Corbally Christopher J., 2009, Stellar Spectral Classification
- 2030 Gray R. O., Corbally C. J., Garrison R. F., McFadden M. T., Bubar E. J., McGahee C. E., O'Donoghue A. A., Knox E. R., 2006, *AJ*, **132**, 161
- 2031 Grenier S., et al., 1999, *A&AS*, **137**, 451
- 2032 Gruberbauer M., et al., 2008, *A&A*, **480**, 223
- 2033 Gruberbauer M., et al., 2011, *A&A*, **530**, A135
- 2034 Gustafsson B., Edvardsson B., Eriksson K., Jørgensen U. G., Nordlund Å., Plez B., 2008, *A&A*, **486**, 951
- 2035 Haakonsen C. B., Rutledge R. E., 2009, *ApJS*, **184**, 138
- 2036 Handler G., Paunzen E., 1999, *A&AS*, **135**, 57
- 2037 Handler G., et al., 2006, *MNRAS*, **366**, 257
- 2038 Hartmann M., Hatzes A. P., 2015, *A&A*, **582**, A84
- 2039 Hauck B., Mermilliod M., 1998, *A&AS*, **129**, 431
- 2040 Hensberge H., et al., 1981, *A&AS*, **46**, 151
- 2041 Hey D. R., et al., 2019, *MNRAS*, **488**, 18
- 2042 Holdsworth D. L., 2016, *IBVS*, **6185**, 1
- 2043 Holdsworth D. L., 2021, *Frontiers in Astronomy and Space Sciences*, **8**, 31
- 2044 Holdsworth D. L., et al., 2014a, *MNRAS*, **439**, 2078
- 2045 Holdsworth D. L., Smalley B., Kurtz D. W., Southworth J., Cunha M. S., Clubb K. I., 2014b, *MNRAS*, **443**, 2049
- 2046 Holdsworth D. L., Kurtz D. W., Smalley B., Saio H., Handler G., Murphy S. J., Lehmann H., 2016, *MNRAS*, **462**, 876
- 2047 Holdsworth D. L., et al., 2018a, *MNRAS*, **473**, 91
- 2048 Holdsworth D. L., Saio H., Bowman D. M., Kurtz D. W., Sefako R. R., Joyce M., Lambert T., Smalley B., 2018b, *MNRAS*, **476**, 601
- 2049 Holdsworth D. L., Cunha M. S., Shibahashi H., Kurtz D. W., Bowman D. M., 2018c, *MNRAS*, **480**, 2976
- 2050 Holdsworth D. L., Saio H., Kurtz D. W., 2019, *MNRAS*, **489**, 4063
- 2051 Houk N., 1978, Michigan catalogue of two-dimensional spectral types for the HD stars
- 2052 Houk N., 1982, Michigan Catalogue of Two-dimensional Spectral Types for the HD stars. Volume\_3. Declinations -40\_f0 to -26\_f0.
- 2053 Houk N., Cowley A. P., 1975, University of Michigan Catalogue of two-dimensional spectral types for the HD stars. Volume I. Declinations -90\_to -53\_f0.
- 2054 Houk N., Smith-Moore M., 1988, Michigan Catalogue of Two-dimensional Spectral Types for the HD Stars. Volume 4, Declinations -26°.0 to -12°.0. Vol. 4
- 2055 Houk N., Swift C., 1999b, Michigan catalogue of two-dimensional spectral types for the HD Stars ; vol. 5. Vol. 5
- 2056 Houk N., Swift C., 1999a, Michigan Spectral Survey, **5**, 0
- 2057 Hubrig S., Nesvacil N., 2007, *MNRAS*, **378**, L16
- 2058 Hubrig S., Kharchenko N., Mathys G., North P., 2000, *A&A*, **355**, 1031
- 2059 Hubrig S., Szeifert T., Schöller M., Mathys G., Kurtz D. W., 2004, *A&A*, **415**, 685
- 2060 Hubrig S., et al., 2005, *A&A*, **440**, L37
- 2061 Hubrig S., North P., Schöller M., Mathys G., 2006, *Astronomische Nachrichten*, **327**, 289
- 2062 Hubrig S., Järvinen S. P., Madej J., Bychkov V. D., Ilyin I., Schöller M., Bychkova L. V., 2018, *MNRAS*, **477**, 3791
- 2063 Hümmerich S., Paunzen E., Bernhard K., 2016, *AJ*, **152**, 104
- 2064 Jackson J., Stoy R. H., 1954, Annals of the Cape Observatory, **17**
- 2065 Joshi S., et al., 2016, *A&A*, **590**, A116
- 2066 Kervella P., Arenou F., Mignard F., Thévenin F., 2019, *A&A*, **623**, A72
- 2067 Kniazev A. Y., Gvaramadze V. V., Berdnikov L. N., 2016, *MNRAS*, **459**, 3068
- 2068 Kniazev A. Y., Gvaramadze V. V., Berdnikov L. N., 2017, in Balega Y. Y., Kudryavtsev D. O., Romanyuk I. I., Yakunin I. A., eds, Astronomical Society of the Pacific Conference Series Vol. 510, Stars: From Collapse to Collapse. p. 480 (arXiv:1612.00292)
- 2069 Kobzar O., et al., 2020, in Wade G., Alecian E., Bohlender D., Sigut A., eds, Vol. 11, Stellar Magnetism: A Workshop in Honour of the Career and Contributions of John D. Landstreet. pp 214–218 (arXiv:2003.05337)
- 2070 Kochukhov O., 2003, *A&A*, **404**, 669
- 2071 Kochukhov O., 2011, in Prasad Choudhary D., Strassmeier K. G., eds, Vol. 273, Physics of Sun and Star Spots. pp 249–255 (arXiv:1010.0264), doi:10.1017/S1743921311015328
- 2072 Kochukhov O., Bagnulo S., 2006, *A&A*, **450**, 763
- 2073 Kochukhov O., Ryabchikova T., 2001, *A&A*, **377**, L22
- 2074 Kochukhov O., Drake N. A., Piskunov N., de la Reza R., 2004, *A&A*, **424**, 935
- 2075 Kochukhov O., Ryabchikova T., Weiss W. W., Landstreet J. D., Lyashko D., 2007, *MNRAS*, **376**, 651
- 2076 Kochukhov O., Ryabchikova T., Bagnulo S., Lo Curto G., 2008, *A&A*, **479**, L29
- 2077 Kochukhov O., Bagnulo S., Lo Curto G., Ryabchikova T., 2009a, *A&A*, **493**, L45
- 2078 Kochukhov O., Shulyak D., Ryabchikova T., 2009b, *A&A*, **499**, 851
- 2079 Kochukhov O., Alentiev D., Ryabchikova T., Boyko S., Cunha M., Tsymbal V., Weiss W., 2013, *MNRAS*, **431**, 2808
- 2080 Kordopatis G., et al., 2013, *AJ*, **146**, 134
- 2081 Kreidl T. J., 1990, *IBVS*, **3539**, 1
- 2082 Kreidl T. J., Kurtz D. W., Bus S. J., Kuschnig R., Birch P. B., Candy M. P., Weiss W. W., 1991, *MNRAS*, **250**, 477
- 2083 Kudryavtsev D. O., Romanyuk I. I., 2003, *Astrophysics*, **46**, 234
- 2084 Kudryavtsev D. O., Romanyuk I. I., 2012, *Astronomische Nachrichten*, **333**, 41
- 2085 Kudryavtsev D. O., Romanyuk I. I., Elkin V. G., Paunzen E., 2006, *MNRAS*, **372**, 1804
- 2086 Kudryavtsev D. O., Romanyuk I. I., Semenko E. A., Solov'ev G. A., 2007, *Astrophysical Bulletin*, **62**, 147
- 2087 Kunder A., et al., 2017, *AJ*, **153**, 75
- 2088 Kupka F., Ryabchikova T. A., Weiss W. W., Kuschnig R., Rogl J., Mathys G., 1996, *A&A*, **308**, 886
- 2089 Kurtz D. W., 1978, *IBVS*, **1436**, 1
- 2090 Kurtz D. W., 1981, *IBVS*, **1915**, 1
- 2091 Kurtz D. W., 1982, *MNRAS*, **200**, 807
- 2092 Kurtz D. W., 1983, *MNRAS*, **205**, 3
- 2093 Kurtz D. W., 1984, *MNRAS*, **209**, 841
- 2094 Kurtz D. W., 1985, *MNRAS*, **213**, 773
- 2095 Kurtz D. W., 1990, *MNRAS*, **242**, 489
- 2096 Kurtz D. W., Cropper M. S., 1981, *IBVS*, **1987**, 1

- 2142 Kurtz D. W., Holdsworth D. L., 2020, *Astrophysics and Space Science* 2210  
 2143 *Proceedings*, 57, 313
- 2144 Kurtz D. W., Kreidl T. J., 1985, *MNRAS*, 216, 987
- 2145 Kurtz D. W., Marang F., 1987a, *MNRAS*, 228, 141
- 2146 Kurtz D. W., Marang F., 1987b, *MNRAS*, 229, 285
- 2147 Kurtz D. W., Martinez P., 1987, *MNRAS*, 226, 187
- 2148 Kurtz D. W., Martinez P., 1993, in Dworetzky M. M., Castelli F., Faraggiana 2216  
 2149 R., eds, *Astronomical Society of the Pacific Conference Series Vol. 44,* 2217  
 2150 *IAU Colloq. 138: Peculiar versus Normal Phenomena in A-type and* 2218  
 2151 *Related Stars.* p. 561
- 2152 Kurtz D. W., Martinez P., 1994, *IBVS*, 4013, 1
- 2153 Kurtz D. W., Allen S., Cropper M. S., 1981, *IBVS*, 2033, 1
- 2154 Kurtz D. W., van Wyk F., Marang F., 1990, *MNRAS*, 243, 289
- 2155 Kurtz D. W., Sullivan D. J., Martinez P., Tripe P., 1994a, *MNRAS*, 270, 674
- 2156 Kurtz D. W., Martinez P., Tripe P., 1994b, *MNRAS*, 271, 421
- 2157 Kurtz D. W., Marang F., van Wyk F., Roberts G., 1996a, *MNRAS*, 280, 1
- 2158 Kurtz D. W., Martinez P., Koen C., Sullivan D. J., 1996b, *MNRAS*, 281, 2226  
 2159 883
- 2160 Kurtz D. W., van Wyk F., Roberts G., Marang F., Handler G., Medupe R., 2227  
 2161 Kilkenny D., 1997a, *MNRAS*, 287, 69
- 2162 Kurtz D. W., Martinez P., Tripe P., Hanbury A. G., 1997b, *MNRAS*, 289, 2228  
 2163 645
- 2164 Kurtz D. W., Elkin V. G., Mathys G., 2003, *MNRAS*, 343, L5
- 2165 Kurtz D. W., et al., 2005, *MNRAS*, 358, 651
- 2166 Kurtz D. W., Elkin V. G., Mathys G., 2006a, *MNRAS*, 370, 1274
- 2167 Kurtz D. W., Elkin V. G., Cunha M. S., Mathys G., Hubrig S., Wolff B., 2233  
 2168 Savanov I., 2006b, *MNRAS*, 372, 286
- 2169 Kurtz D. W., Hubrig S., González J. F., van Wyk F., Martinez P., 2008, 2234  
 2170 *MNRAS*, 386, 1750
- 2171 Kurtz D. W., et al., 2011, *MNRAS*, 414, 2550
- 2172 Landstreet J. D., Mathys G., 2000, *A&A*, 359, 213
- 2173 Lorenz D., Handler G., Kurtz D. W., 2005, *IBVS*, 5651, 1
- 2174 Lüftinger T., Kochukhov O., Ryabchikova T., Piskunov N., Weiss W. W., 2242  
 2175 Ilyin I., 2010, *A&A*, 509, A71
- 2176 Manfroid J., Mathys G., 1986, *A&AS*, 64, 9
- 2177 Martinez P., 1991, *IBVS*, 3621, 1
- 2178 Martinez P., Kauffmann G., 1990, *IBVS*, 3507, 1
- 2179 Martinez P., Kurtz D. W., 1990, *IBVS*, 3509, 1
- 2180 Martinez P., Kurtz D. W., 1991a, *IBVS*, 3553, 1
- 2181 Martinez P., Kurtz D. W., 1991b, *IBVS*, 3611, 1
- 2182 Martinez P., Kurtz D. W., 1992, *IBVS*, 3750, 1
- 2183 Martinez P., Kurtz D. W., 1994a, *MNRAS*, 271, 118
- 2184 Martinez P., Kurtz D. W., 1994b, *MNRAS*, 271, 129
- 2185 Martinez P., Kurtz D. W., Heller C. H., 1990, *MNRAS*, 246, 699
- 2186 Martinez P., Kurtz D. W., Kauffmann G. M., 1991, *MNRAS*, 250, 666
- 2187 Martinez P., Kurtz D. W., Kreidl T. J., Koen C., van Wyk F., Marang F., 2255  
 2188 Roberts G., 1993a, *MNRAS*, 263, 273
- 2189 Martinez P., Kurtz D. W., Ashley R., 1993b, *IBVS*, 3844, 1
- 2190 Martinez P., Kurtz D. W., van Wyk F., 1994, *MNRAS*, 271, 305
- 2191 Martinez P., Kurtz D. W., Hoffman M. J. H., van Wyk F., 1995, *MNRAS*, 2259  
 2192 276, 1435
- 2193 Martinez P., Marang F., van Wyk F., Roberts G. R., 1998a, *The Observatory*, 2261  
 2194 118, 153
- 2195 Martinez P., Koen C., Sullivan D. J., 1998b, *MNRAS*, 300, 188
- 2196 Martinez P., Meintjes P., Ratcliff S. J., Engelbrecht C., 1998c, *A&A*, 334, 2264  
 2197 606
- 2198 Mason B. D., Wycoff G. L., Hartkopf W. I., Douglass G. G., Worley C. E., 2266  
 2199 2001, *AJ*, 122, 3466
- 2200 Mathys G., 1991, *A&AS*, 89, 121
- 2201 Mathys G., 1995, *A&A*, 293, 746
- 2202 Mathys G., 2017, *A&A*, 601, A14
- 2203 Mathys G., Hubrig S., 1997, *A&AS*, 124, 475
- 2204 Mathys G., Kharchenko N., Hubrig S., 1996, *A&A*, 311, 901
- 2205 Mathys G., Hubrig S., Landstreet J. D., Lanz T., Manfroid J., 1997, *A&AS*, 2273  
 2206 123, 353
- 2207 Mathys G., Kurtz D. W., Elkin V. G., 2007, *MNRAS*, 380, 181
- 2208 Mathys G., Khalack V., Landstreet J. D., 2020a, *A&A*, 636, A6
- 2209 Mathys G., Kurtz D. W., Holdsworth D. L., 2020b, *A&A*, 639, A31
- 2210 Matthews J. M., 1987, PhD thesis, THE UNIVERSITY OF WESTERN 2211  
 2212 ONTARIO (CANADA).
- 2213 Matthews J. M., Kurtz D. W., Martinez P., 1999, *ApJ*, 511, 422
- 2214 McDonald I., Zijlstra A. A., Boyer M. L., 2012, *MNRAS*, 427, 343
- 2215 McDonald I., Zijlstra A. A., Watson R. A., 2017, *MNRAS*, 471, 770
- 2216 Medupe R., Kurtz D. W., Elkin V. G., Mguda Z., Mathys G., 2015, *MNRAS*, 446, 1347
- 2217 Michaud G., Megessier C., Charland Y., 1981, *A&A*, 103, 244
- 2218 Mkrichian D. E., Hatzes A. P., 2005, *A&A*, 430, 263
- 2219 Moon T. T., Dworetzky M. M., 1985, *MNRAS*, 217, 305
- 2220 Murphy S. J., Saio H., Takada-Hidai M., Kurtz D. W., Shibahashi H., Takata 2221  
 2222 M., Hey D. R., 2020, *MNRAS*, 498, 4272
- 2223 Nelson M. J., Kreidl T. J., 1993, *AJ*, 105, 1903
- 2224 Netopil M., Paunzen E., Hümmerich S., Bernhard K., 2017, *MNRAS*, 468, 2745
- 2225 North P., 1984, *A&A*, 141, 328
- 2226 North P., Jaschek C., Hauck B., Figueras F., Torra J., Kunzli M., 1997, in 2227  
 2228 Bonnet R. M., et al., eds, *ESA Special Publication Vol. 402, Hipparcos* 2229  
 2230 *- Venice '97.* pp 239–244
- 2231 Oelkers R. J., et al., 2018, *AJ*, 155, 39
- 2232 Paunzen E., 2015, *A&A*, 580, A23
- 2233 Paunzen E., Netopil M., Rode-Paunzen M., Handler G., Božić H., Ruždjak 2234  
 2235 D., Sudar D., 2012, *A&A*, 542, A89
- 2236 Paunzen E., et al., 2018, *Research in Astronomy and Astrophysics*, 18, 135
- 2237 Perraut K., Brandão I., Cunha M., Shulyak D., Mourard D., Nardetto N., ten 2238  
 2239 Brummelaar T. A., 2016, *A&A*, 590, A117
- 2240 Perry C. L., 1991, *PASP*, 103, 494
- 2241 Perry C. L., Christodoulou D. M., 1996, *PASP*, 108, 772
- 2242 Pollacco D. L., et al., 2006, *PASP*, 118, 1407
- 2243 Quiral-Manosalva P., Cunha M. S., Kochukhov O., 2018, *MNRAS*, 480, 1676
- 2244 Reiners A., Zechmeister M., 2020, *ApJS*, 247, 11
- 2245 Renson P., Manfroid J., 2009, *A&A*, 498, 961
- 2246 Renson P., Gerbaldi M., Catalano F. A., 1991, *A&AS*, 89, 429
- 2247 Richer J., Michaud G., Turcotte S., 2000, *ApJ*, 529, 338
- 2248 Ricker G. R., et al., 2015, *Journal of Astronomical Telescopes, Instruments,* 2249  
 2250 *and Systems*, 1, 014003
- 2251 Romanyuk I. I., Semenko E. A., Kudryavtsev D. O., Moiseeva A. V., 2016, 2252  
 2253 *Astrophysical Bulletin*, 71, 302
- 2254 Rusomarov N., et al., 2013, *A&A*, 558, A8
- 2255 Ryabchikova T. A., 2005, *Astronomy Letters*, 31, 388
- 2256 Ryabchikova T., Nesvacil N., Weiss W. W., Kochukhov O., Stütz C., 2004, 2257  
 2258 *A&A*, 423, 705
- 2259 Ryabchikova T., Sachkov M., Kochukhov O., Lyashko D., 2007, *A&A*, 473, 907
- 2260 Ryabchikova T., Kochukhov O., Bagnulo S., 2008, *A&A*, 480, 811
- 2261 Ryabchikova T., Piskunov N., Kurucz R. L., Stempels H. C., Heiter U., 2262  
 2263 Pakhomov Y., Barklem P. S., 2015, *Phys. Scr.*, 90, 054005
- 2264 Sahoo S. K., Baran A. S., Sanjayan S., Ostrowski J., 2020, *MNRAS*, 499, 5508
- 2265 Saio H., 2005, *MNRAS*, 360, 1022
- 2266 Saio H., Gautschi A., 2004, *MNRAS*, 350, 485
- 2267 Samus' N. N., et al., 2003, *Astronomy Letters*, 29, 468
- 2268 Schneider H., Weiss W. W., 1990, *IBVS*, 3520, 1
- 2269 Schneider H., Kreidl T. J., Weiss W. W., 1992, *A&A*, 257, 130
- 2270 Schofield M., et al., 2019, *ApJS*, 241, 12
- 2271 Shavrina A. V., Khalack V., Glagolevskij Y., Lyashko D., Landstreet J., 2272  
 2273 Leone F., Giarrusso M., 2014, in Petit P., Jardine M., Spruit H. C., eds, 2274  
 2275 Vol. 302, *Magnetic Fields throughout Stellar Evolution.* pp 274–275, 2276  
 2277 doi:10.1017/S1743921314002269
- 2278 Shibahashi H., 1979, *PASJ*, 31, 87
- 2279 Shibahashi H., Saio H., 1985, *PASJ*, 37, 601
- 2280 Sikora J., Wade G. A., Power J., Neiner C., 2019a, *MNRAS*, 483, 2300
- 2281 Sikora J., Wade G. A., Power J., Neiner C., 2019b, *MNRAS*, 483, 3127
- 2282 Sikora J., et al., 2019c, *MNRAS*, 487, 4695
- 2283 Skiff B. A., 2014, *VizieR Online Data Catalog*, p. B/mk
- 2284 Smalley B., et al., 2015, *MNRAS*, 452, 3334
- 2285 Sousa J. C., Cunha M. S., 2011, *MNRAS*, 414, 2576

- 2278 Stepień K., 2000, *A&A*, **353**, 227  
 2279 Stevens D. J., Stassun K. G., Gaudi B. S., 2017, *AJ*, **154**, 259  
 2280 Stibbs D. W. N., 1950, *MNRAS*, **110**, 395  
 2281 Suchkov A. A., Makarov V. V., Voges W., 2003, *ApJ*, **595**, 1206  
 2282 Tassoul M., 1980, *ApJS*, **43**, 469  
 2283 Trifonov T., Tal-Or L., Zechmeister M., Kaminski A., Zucker S., Mazeh T.,  
 2284 2020, *A&A*, **636**, A74  
 2285 Vauclair S., Théado S., 2004, *A&A*, **425**, 179  
 2286 Vogt N., Faundez A. M., 1979, *A&AS*, **36**, 477  
 2287 Wade G. A., 1997, *A&A*, **325**, 1063  
 2288 Watson C. L., Henden A. A., Price A., 2006, Society for Astronomical  
 2289 Sciences Annual Symposium, **25**, 47  
 2290 Weiss W. W., et al., 2016, *A&A*, **588**, A54  
 2291 Weiss W. W., et al., 2020, *A&A*, **642**, A64  
 2292 White T. R., Bedding T. R., Stello D., Kurtz D. W., Cunha M. S., Gough  
 2293 D. O., 2011, *MNRAS*, **415**, 1638



**Figure B1.** SAAO spectrum of TIC 170586794 (centre), with stars of spectral class F5V (above) and F0V (below) for comparison. Lines of Eu and Cr are enhanced in this star, leading to the classification of F5VpEuCr.

## 2294 APPENDIX A: AUTHOR AFFILIATIONS


- 2295 <sup>1</sup>Jeremiah Horrocks Institute, University of Central Lancashire, Pre-  
 2296 ston PR1 2HE, UK  
 2297 <sup>2</sup>Instituto de Astrofísica e Ciências do Espaço, Universidade do  
 2298 Porto CAUP, Rua das Estrelas, PT4150-762 Porto, Portugal  
 2299 <sup>3</sup>Centre for Space Research, Physics Department, North West Uni-  
 2300 versity, Mahikeng, 2745, South Africa  
 2301 <sup>4</sup>DTU Space, National Space Institute, Technical University of Den-  
 2302 mark, Elektrovej 328, 2800 Kgs., Lyngby, Denmark  
 2303 <sup>5</sup>Stellar Astrophysics Centre, Aarhus University, DK-8000 Aarhus  
 2304 C, Denmark  
 2305 <sup>6</sup>School of Physics, Sydney Institute for Astronomy (SfA), The  
 2306 University of Sydney, NSW 2006, Australia  
 2307 <sup>7</sup>Institute of Astronomy, KU Leuven, Celestijnenlaan 200D, 3001  
 2308 Leuven, Belgium  
 2309 <sup>8</sup>Département de Physique et d’Astronomie, Université de Mon-  
 2310 cton, 18 avenue Antonine-Maillet, Moncton, NB E1A 3E9, Canada  
 2311 <sup>9</sup>Department of Chemistry and Physics, Florida Gulf Coast Univer-  
 2312 sity, 10501 FGCU Blvd. S., Fort Myers, FL 33965 USA  
 2313 <sup>10</sup>Astronomical Institute of University of Wrocław, ul. Kopernika  
 2314 11, 51-622 Wrocław, Poland  
 2315 <sup>11</sup>Ankara University, Faculty of Science, Dept. of Astronomy and  
 2316 Space Sciences, 06100, Tandogan - Ankara / Turkey  
 2317 <sup>12</sup>Department of Astronomy, School of Physics, Peking University,  
 2318 Beijing 100871, People’s Republic of China  
 2319 <sup>13</sup>Kavli Institute for Astronomy and Astrophysics, Peking Univer-  
 2320 sity, Beijing 100871, People’s Republic of China  
 2321 <sup>14</sup>Konkoly Observatory, MTA CSFK, H-1121, Konkoly Thege Mik-  
 2322 lós út 15-17, Budapest, Hungary  
 2323 <sup>15</sup>MTA CSFK Lendület Near-Field Cosmology Research Group  
 2324 <sup>16</sup>Unknown  
 2325 <sup>17</sup>Instituto de Astronomía–Universidad Nacional Autónoma de  
 2326 México, Ap. P. 877, Ensenada, BC 22860, Mexico  
 2327 <sup>18</sup>Department of Physics and Astronomy, Uppsala University, Box  
 2328 516, 75120 Uppsala, Sweden  
 2329 <sup>19</sup>Instituto de Astrofísica de Andalucía (CSIC), Glorieta de la As-  
 2330 tronomía s/n, E-18008 Granada, Spain  
 2331 <sup>20</sup>Physics Department, Mount Allison University, Sackville, NB,  
 2332 Canada, E4L1E6  
 2333 <sup>21</sup>National Astronomical Research Institute of Thailand, 260 Moo  
 2334 4, T. Donkaew, A. Maerim, Chiangmai, 50180 Thailand  
 2335 <sup>22</sup>Department of Theoretical Physics and Astrophysics, Masaryk  
 2336 University, Kotlářská 2, 611 37 Brno, Czech Republic  
 2337 <sup>23</sup>Unknown

- 2338 <sup>24</sup>Physics and Astronomy Department, Bishop’s University, 2600  
 2339 College Street, Sherbrooke, QC J1M 1Z7, Canada  
 2340 <sup>25</sup>School of Physics, Xi’an Jiaotong University, 710049 Xi’an, Peo-  
 2341 ple’s Republic of China  
 2342 <sup>26</sup>New?  
 2343 <sup>27</sup>Unknown  
 2344 <sup>28</sup>Dept. Física Teórica y del Cosmos, Universidad de Granada, Cam-  
 2345 pus de Fuentenueva s/n, E-18071 Granada, Spain  
 2346 <sup>29</sup>Los Alamos National Laboratory, MS T-082, Los Alamos, NM  
 2347 87545 USA  
 2348 <sup>30</sup>Astrophysics Group, Keele University, Staffordshire ST5 5BG,  
 2349 UK  
 2350 <sup>31</sup>Institute of Astronomy and NAO, Bulgarian Academy of Sci-  
 2351 ences, blvd. Tsarigradsko chaussee 72, Sofia 1784, Bulgaria


## 2352 APPENDIX B: SPECTRAL CLASSIFICATION PLOTS

2353 Here we provide the spectral classification plots used to provide  
 2354 new or updated spectral classifications. We have used the SpUp-  
 2355 NIC long slit spectrograph mounted on the SAAO 1.9-m telescope  
 2356 to record these spectra. We used grating 4 with a grating angle of  
 2357  $4.2^\circ$ , with observations typically made with a 1 arcsec slit to reach  
 2358 a resolving power of  $R \sim 3000$ , but was widened in poor seeing or  
 2359 low sky transparency situations. The data were flat field and bias  
 2360 corrected, cleaned of cosmic rays, and had a wavelength solution  
 2361 applied using a CuAr reference spectrum obtained directly after the  
 2362 science observation. The spectra were normalised using an auto-  
 2363 mated spline fitting procedure in *ISPEC* spectral analysis software  
 2364 (Blanco-Cuaresma et al. 2014; Blanco-Cuaresma 2019).


2365 This paper has been typeset from a  $\text{\TeX}/\text{\LaTeX}$  file prepared by the author.




**Figure B2.** SAAO spectrum of TIC 294266638. Lines of Sr and Eu are noted as being pronounced in this star, leading to the classification of Ap SrEu.




**Figure B5.** The SAAO spectrum of TIC 17676722 with lines of Sr, Eu and Cr identified. We refine the spectral classification to Ap SrEuCr.




**Figure B3.** The SAAO spectrum of TIC 356088697 with lines of Sr, Eu and Cr identified. We refine the spectral classification to Ap SrEuCr.




**Figure B6.** The SAAO spectrum of TIC 41259805 with lines of Sr, Eu and Cr identified. We refine the spectral classification to Ap SrEu(Cr).




**Figure B4.** The SAAO spectrum of TIC 12968953 with lines of Sr, Eu and Cr identified. We refine the spectral classification to Ap SrEuCr.




**Figure B7.** The SAAO spectrum of TIC 49818005 with lines of Sr, Eu and Cr identified. We provide a spectral classification of Fp SrEu(Cr).




**Figure B8.** The SAAO spectrum of TIC 152808505. We provide a spectral classification of Fp EuCr.



**Figure B11.** The SAAO spectrum of TIC 3814749 showing the lack of strong chemical peculiarities in this star.



**Figure B9.** The SAAO spectrum of TIC 156886111 with which we deduce a classification of Ap SrEuCr.



**Figure B10.** The SAAO spectrum of TIC 350146296 with which we deduce a classification of F0p EuCr.