

# TESS Cycle 2 observations of roAp stars with 2-min cadence data

D. L. Holdsworth,<sup>\*</sup> M. S. Cunha, M. Lares-Martiz, D. W. Kurtz, V. Antoci,  
 S. Barceló Forteza, P. De Cat, A. Derekas, C. Kayhan, D. Ozuyar, M. Skarka, D. R. Hey,  
 F. Shi, D. M. Bowman, O. Kobzar, A. Ayala, Zs. Bognár, D. L. Buzasi,  
 M. Ebadi, L. Fox-Machado, A. García Hernández, H. Ghasemi, J. A. Guzik, G. Handler,  
 A. Hasanzadeh, R. Jayaraman, V. Khalack, O. Kochukhov, C. C. Lovekin, P. Mikołajczyk,  
 D. Mkrtichian, S. J. Murphy, E. Niemczura, B. G. Olafsson, J. Pascual-Granado, E. Paunzen,  
 N. Posiłek, A. Ramón-Ballesta, H. Safari, A. Samadi-Ghadim, B. Smalley, Á. Sódor,  
 I. Stateva, J. C. Suárez, R. Szabó, T. Wu, E. Ziaali, and W. Zong

*Author affiliations shown in Appendix A.*

27 October 2023

## ABSTRACT

We present the results of a systematic search of the *Transiting Exoplanet Survey Satellite* (*TESS*) 2-min cadence data for new rapidly oscillating Ap (roAp) stars observed during the Cycle 2 phase of its mission. We found seven new roAp stars previously unreported as such and present the analysis of a further 25 roAp stars that are already known. Three of the new stars show multiperiodic pulsations, while all new members are rotationally variable stars, leading to almost 70 per cent (22) of the roAp stars presented being  $\alpha^2$  CVn stars. We show that targeted observations of known chemically peculiar stars is likely to overlook many new roAp stars, and demonstrate that multi-epoch observations are necessary to see pulsational behaviour in some cases. We found a lack of roAp stars close to the blue edge of the theoretical roAp instability strip, and reaffirmed that mode instability is observed more frequently with precise, space-based observations. In addition to the Cycle 2 observations, we analysed any available *TESS* data for all known roAp stars. This amounts to 18 further roAp stars observed by *TESS*. Finally, we list six known roAp stars that *TESS* is yet to observe. With this information, we deduce that the incidence of roAp stars amongst the Ap star population is just 5.5 per cent, raising fundamental questions about the conditions required to excite pulsations in Ap stars. This work, coupled with Paper I (Holdsworth et al. 2021), presents the most comprehensive, homogeneous study of the roAp stars in the *TESS* nominal mission, with a collection of 112 confirmed roAp stars in total.

**Key words:** asteroseismology – stars: chemically peculiar – stars: oscillations – techniques: photometric – stars: variables – stars: individual

## 1 INTRODUCTION

The *Transiting Exoplanet Survey Satellite* (*TESS*; Ricker et al. 2015) is providing a rich photometric dataset for millions of stars in most regions of the sky. While the primary mission focus is on the detection of exoplanet transits, the acquired data are well suited for studies of stellar variability. *TESS* observes the sky in strips of  $24^\circ \times 96^\circ$  for about 27 d each, referred to as *sectors*. *TESS* is observing in 1-year *cycles*, each composed of 13 sectors. For the nominal mission comprising Cycles 1 & 2, *TESS* recorded data in two cadences: 30 min for the Full Frame Images (FFIs) and 2 min for a selection of  $\sim 20\,000$  stars per sector. This 2-min cadence data provides a dataset with which it is possible to perform a homogeneous search for high-frequency stellar variability. We have previously exploited this capability to search for high-frequency pulsations in the chemically

peculiar Ap stars (Holdsworth et al. 2021, hereafter Paper I) in the Cycle 1 data. Here we extend this study to encompass the 2-min Cycle 2 observations.

The rapidly oscillating Ap (roAp) stars are a rare subset of the chemically peculiar, magnetic, Ap stars (Kurtz 1982). To date there are of the order 100 known in the literature (see e.g., Smalley et al. 2015; Joshi et al. 2016; Cunha et al. 2019; Holdsworth et al. 2021, for catalogues and lists), discovered through targeted high-speed ground-based photometry and spectroscopy (e.g., Martinez et al. 1991; Paunzen et al. 2018; Freyhammer et al. 2008; Kochukhov et al. 2013) and space-based survey data (e.g., Gruberbauer et al. 2011; Balona et al. 2011b; Hey et al. 2019; Cunha et al. 2019).

The broader population of Ap stars are characterised by strong, global, magnetic fields (Mathys 2017) that suppress near-surface convection in the stellar atmosphere (Michaud et al. 1981; LeBlanc et al. 2009; Alecian & Stift 2010), leading to gravitational settling of some

<sup>\*</sup> e-mail: dlholdsworth@uclan.ac.uk

light elements and radiative levitation of other heavy elements such as Pr, Nd, Eu and Y, which can reach atmospheric abundances of a million times that of the Sun (e.g., Lüftinger et al. 2010). These levitated chemical elements create persistent atmospheric anomalies that may be stable for at least a century (e.g., Bychkov et al. 2016). These anomalies, i.e. chemical spots, modulate the observed flux, enabling the accurate determination of the stellar rotation period. Rotationally variable Ap stars are known as  $\alpha^2$  CVn stars.

The roAp stars are characterised by high overtone ( $n \gtrsim 15$ ), low degree ( $\ell \lesssim 3$ ) pressure (p) mode pulsations with frequencies in the range 0.7 – 3.6 mHz (60 – 310 d<sup>-1</sup>;  $P = 4.7$  – 23.6 min). In many non-magnetic stars, the pulsation axis is aligned with the rotation axis as this is the axis of maximum deformation from spherical symmetry. In the roAp stars, however, the pulsation axis is closely aligned with the magnetic axis which in turn is inclined to the rotation axis. Such a geometry means the pulsations are viewed from varying aspects as the star rotates, leading to the oblique pulsator model (e.g., Kurtz 1982; Bigot & Dziembowski 2002), where the mode amplitude and phase appear modulated over the stellar rotation period. In a Fourier spectrum of a light curve, this results in a multiplet with  $2\ell + 1$  components for a non-distorted mode, with the components split from the central mode frequency by exactly the stellar rotation frequency. With a distorted mode, the multiplet shows more components (e.g., Holdsworth et al. 2016) and the pulsation amplitude never reaches zero as a node crosses the line of sight. The distortion arises from the magnetic field’s interaction with the pulsation mode’s eigenfunction, which then requires higher order spherical harmonics to describe the mode (see Holdsworth et al. 2018a; Shi et al. 2021, and references therein for discussions).

The driving mechanism for the pulsations in these stars is still uncertain. In many cases, theoretical models can recreate the observations by employing the opacity ( $\kappa$ ) mechanism acting in the hydrogen ionisation layers in regions of the star where the magnetic field suppresses convection (Balmforth et al. 2001). However, this method cannot explain all of the observed frequencies, especially those at very high frequency, where instead it is suggested that turbulent pressure plays a role in the mode excitation (Cunha et al. 2013).

The most promising stars for asteroseismic insight are those that show multiperiodic variability. In a non-magnetic star, the high frequency pulsations occur in the asymptotic regime, forming a series of alternating odd and even degree modes (Shibahashi 1979; Tassoul 1980) that allow the use of asteroseismic techniques to constrain the stellar properties, e.g., Deal et al. (2021). However, the situation is complicated in some roAp stars as the pulsations become magneto-acoustic in the outer layers of the star, and their frequencies suffer from magnetic perturbation (Cunha & Gough 2000; Cunha 2006; Saio & Gautschy 2004). Detecting such phenomena in the roAp stars helps to drive forward the theoretic modelling of the class, and provide the opportunity to fully understand roAp stars.

Recently, the validity of the roAp class was questioned by Balona (2022), who suggested that all roAp stars are just  $\delta$  Sct stars. However, this is clearly not the case. While there is overlap in the frequency regime where both  $\delta$  Sct pulsations and roAp pulsations are found (in different stars), the roAp stars are distinct from the  $\delta$  Sct stars given their Ap chemical peculiarities, oblique pulsations, and magnetic fields. There are some cases where roAp stars do show low-frequency  $\delta$  Sct pulsations (e.g., Murphy et al. 2020), and new results where a magnetic field has been confirmed in an Ap star potentially showing  $\delta$  Sct pulsations (Hubrig et al. 2023). While the former work rules out a companion to the roAp star, they place limits on the magnetic field strength that can suppress low-overtone pulsations. The latter work cautions that an undetected companion to the Ap star may provide

the  $\delta$  Sct pulsations and that their derived field strength is close to the limit provided by Murphy et al. (2020), thus no certain conclusions can be drawn on the presence of  $\delta$  Sct modes in a magnetic Ap star. The suggestion, therefore, that the roAp classification should be abandoned is not useful and is rejected. It is only with careful study and interpretation of valid evidence that stars should be admitted to the roAp class of pulsating variable.

## 2 DATA SAMPLE AND SEARCH STRATEGY

To present a homogeneous sample of roAp stars observed by *TESS* during its nominal mission, we followed the same procedure as in Paper I. Team 1 took the 2-min target list for each sector of observations from *TESS*<sup>1</sup> which consisted of 20 000 stars per sector. These lists were then crossmatched with version 8 of the *TESS* Input Catalogue (TIC; Stassun et al. 2019) with an 8 arcsec radius, giving us access to the secondary information in the TIC to refine our targets. Using only the  $T_{\text{eff}}$  parameter, we selected stars with temperatures of 6000 K or greater (including stars with a ‘null’ value). This amounted to about 7500 stars per sector, with 93 163 light curves in total, corresponding to 36 209 unique stars (with 15 675 stars observed in more than one sector). The light curves for each star in all available sectors were downloaded from the Mikulski Archive for Space Telescopes (MAST) server. These data have been processed with the Science Processing Operations Center (SPOC) pipeline (Jenkins et al. 2016), which are the most suitable data for the investigation of stellar variability (Skarka et al. 2022). In the following we used the Pre-search Data Conditioning Simple Aperture Photometry (PDC\_SAP) data unless otherwise stated.

Initially, two independent analysis techniques were implemented for the search of pulsational variability. For one team, Team 1, each individual light curve was automatically prewhitened to a frequency of 0.23 mHz (20 d<sup>-1</sup>) to an amplitude limit of the approximate noise level between 2.3 – 3.3 mHz (200 – 300 d<sup>-1</sup>) to remove instrumental artefacts and any low-frequency signal whose window function affects the noise at high frequency. If multiple sectors for a target were available, the prewhitened sectors were then combined. An amplitude spectrum of the light curve was calculated to the Nyquist frequency of 4.2 mHz (360 d<sup>-1</sup>), which also included a calculation of the false alarm probability (FAP).

From these amplitude spectra, we selected stars that showed peaks with frequencies > 0.52 mHz (> 45 d<sup>-1</sup>) and a corresponding FAP of < 0.1. The amplitude spectra of these stars were then plotted for visual inspection. This totalled 6713 stars, of which 2125 stars have multi-sector observations.

A second team, Team 2, used a complementary method which calculated the skewness of the amplitude spectrum (Murphy et al. 2019) of the MAST PDC\_SAP data above 0.46 mHz (40 d<sup>-1</sup>). Where multiple sectors of data were available, they were combined to a single light curve. If the skewness was greater than 5, then all peaks in the amplitude spectrum with FAP values below 0.05 were extracted and the star flagged as variable for later human inspection. This method produced a total of 189 variable star candidates.

These two lists were then combined to produce a master list. An initial pass was made with the *ECHO* code (described in Antoci et al. 2019) to flag possible false positive detections with human inspection of all produced outputs (light curves, Fourier transform plots and detected peaks). A detection was determined to be false positive if

<sup>1</sup> <https://tess.mit.edu/observations/target-lists/>

the S/N of the high-frequency signal was below 4.0; if there was obvious contamination from a low-frequency harmonic series; or if the amplitude spectrum displayed obvious characteristics of either  $\delta$  Sct stars (with many modes in the  $\approx 0.2 - 1.0$  mHz range, e.g., Bowman & Kurtz 2018), sdBV stars (stars with pulsations in low and high-frequency ranges) or pulsating white dwarfs (modes with amplitudes significantly greater than those known in roAp stars). The final sample consisted of 163 stars.

Further, we solicited input from the TESS Asteroseismic Science Consortium (TASC) Working Group 4 (WG4; AF-stars). Contributions from the WG4 are as follows:

- *Team A* used the TESS Input Catalog (Stassun et al. 2018) and Gaia DR2 Catalog (Andrae et al. 2018) to look for all stars with SIMBAD spectral types between B8 and F2 (stars with no classification were retained) in the Northern ecliptic hemisphere with effective temperatures between 6000 K and 10000 K, luminosity between 5 and  $100 L_{\odot}$  and a contamination ratio lower than 0.01. Using the interpolation and frequency analysis routine of Barceló Forteza et al. (2015), the team found 121 candidates where the highest amplitude peak was in the frequency range 0.7 – 4.0 mHz and without lower frequency modes in order to avoid the  $\delta$  Sct star pulsation regime (e.g., Aerts et al. 2010; Barceló Forteza et al. 2020, and references therein).

- *Team B* used the Cycle 2 data to search for targets with a magnitude brighter than  $T = 10$  mag and a temperature of  $>6000$  K. The resultant SAP light curves were used to derive rotation periods which were then removed from the data before pulsation frequencies were searched for using the Lomb-Scargle method. This procedure resulted in 9 candidates.

- *Team C* considered bright stars located close to the northern ecliptic pole (within  $15^{\circ}$ ), which were brighter than  $T = 11$  mag, with temperatures in the range  $6000 < T_{\text{eff}} < 10000$  K (Skarka et al. 2022). Frequency spectra, in the range 0 – 4.2 mHz ( $0 - 360 \text{ d}^{-1}$ ) were created from the PDC\_SAP data obtained from the MAST and searched for significant peaks. In total, 1600 stars showed peaks in the frequency range 0.9 – 4.2 mHz ( $80 - 360 \text{ d}^{-1}$ ) with a SNR  $> 4$ . However, only stars with peaks with SNR  $> 5$  were selected as good candidates (8 stars).

- *Team D* provided 55 roAp candidates using the TESS-Ap procedure (Khalack et al. 2021). A search for pulsations in the frequency range 1.2–4.2 mHz ( $100 - 360 \text{ d}^{-1}$ ) was conducted to limit the inclusion of  $\delta$  Sct stars. Only stars in the temperature range 6000–10000 K were considered. Peaks were considered real if the SNR  $> 4$ . After the automatic selection, manual inspection was conducted to reject false positives.

- *Team E* used the TIC to select all stars in the temperature range 6000 – 10000 K for which 2-min cadence data were available. The light curves were downloaded from the MAST server, from which the PDC\_SAP flux data were extracted. Only the first  $\sim 12.5$  d of data were selected to avoid the central downlink gap. Any other gaps are much smaller in size and a piecewise cubic Hermite polynomial was used to fill them with interpolated data at 2-min cadence. No detrending of low-frequency variability was performed. The sampling intervals were then regularised in order to obtain an evenly sampled time series that was analysed using an FFT to calculate the amplitude spectrum to the Nyquist frequency. A star is selected to be variable when a local maximum is found in the amplitude spectrum to have a SNR  $> 5$  and a frequency greater than 0.5 mHz ( $\sim 43 \text{ d}^{-1}$ ). This procedure returned 1241 stars which were vetted by eye to produce a final list of 1134 variable candidates.

- Finally, 8 additional targets were submitted from members of WG4 where individuals had made private note of stars of interest.

All of these inputs from WG4 members were cross-checked with the results from Team 1 and Team 2 to avoid duplication. Team 1 then re-vetted all unique submissions to reject obvious non-roAp variable stars to produce a final sample list of 116 stars. This final sample was redistributed amongst the members of the TASC WG4 for detailed analysis to confirm the presence of a positive roAp detection, and to extract rotation periods from the light curves.

This process identified seven new roAp stars previously unreported as such in the literature (Sec. 4.1) and nine roAp stars discovered through TESS observations (Sec. 4.2). There were also positive detections of pulsations in 12 roAp stars known prior to the launch of TESS (Sec. 4.3), with four roAp stars where TESS did not detect pulsational variability. These four stars were originally identified in Kepler data which has a lower noise threshold than the TESS data analysed here. Finally, we present seven roAp candidate stars where the observational and archival data do not allow us to conclude with certainty that the star is a true roAp star.

### 3 SPECTROSCOPIC OBSERVATIONS

#### 3.1 LAMOST data

The Large Sky Area Multi-Object Fiber Spectroscopic Telescope (LAMOST) is a 4-m class telescope located at the Xinglong observatory (China) that is operated and managed by the National Astronomical Observatories, Chinese Academy of Sciences (Cui et al. 2012). Its scientific observations started in October 2011. With its quasi-meridian transit configuration, it is possible to observe northern stars with a declination above  $\sim 10^{\circ}$  from about two hours before to about two hours after meridian passage. This unique instrument combines a large aperture (primary mirror of  $5.72 \text{ m} \times 4.40 \text{ m}$ ; secondary mirror of  $6.67 \text{ m} \times 6.05 \text{ m}$ ) with a large field of view ( $\sim 20 \text{ deg}^2$ ). This circular field with a diameter of  $5^{\circ}$  is homogeneously covered with 4000 fibres, connected to 16 spectrographs with 250 fibres each. Currently, observations can be done either in the full optical range ( $3700 - 9000 \text{ \AA}$ ) in the low-resolution mode (LAMOST-LRS;  $R \sim 1800$ ; Zhao et al. 2012) or in a blue ( $4950 - 5350 \text{ \AA}$ ) and red ( $6300 - 6800 \text{ \AA}$ ) wavelength range in the medium-resolution mode (LAMOST-MRS;  $R \sim 7500$ ; Liu et al. 2020). The LAMOST-LRS observations are single shot observations while the LAMOST-MRS observations are done in groups of observations in concatenation as long as the observed field remains visible. We checked the LAMOST database and found data for six of our targets. Four targets had one spectrum each (TIC 21024812, TIC 101624823, TIC 118247716, and TIC 467074220) in LAMOST-LRS and two targets had groups of observations in LAMOST-MRS (TIC 26833276; 1 group of spectra, TIC 259017938; 6 groups of spectra). The pipeline-reduced spectra (Luo et al. 2015) from data release 9 (DR9, v1.0)<sup>2</sup> were used to check if typical chemical peculiarities expected for Ap stars can be identified in the data.

#### 3.2 HERMES data

The High Efficiency and Resolution Mercator Echelle Spectrograph (HERMES; Raskin et al. 2011) is mounted to the 1.2-m Mercator telescope at the Observatorio del Roque de los Muchachos on La

<sup>2</sup> <http://www.lamost.org/dr9/>

Palma in the Canary Islands, Spain, and is operated by KU Leuven<sup>3</sup>. HERMES is a fibre-fed prism-cross-dispersed echelle spectrograph with an effective wavelength coverage between 3770 and 9000 Å, and in its standard high-resolution mode has a spectral resolving power of  $\sim 85000$ . Reduced spectra are automatically produced by an efficient PYTHON pipeline, HERMESDRS (v7.0), which includes bias subtraction and flat fielding, as well as wavelength calibration using a ThArNe spectrum, echelle order merging, and removal of cosmic rays. With its Northern hemisphere observing window, as of semester 2018b, we embarked upon a large programme (program 85; PI Bowman) to assemble high signal-to-noise ( $S/N > 100$ ), multi-epoch (minimally two epochs), high-resolution HERMES spectra of all bright ( $V \leq 10$  mag) Ap stars in the catalogue of [Renson & Manfroid \(2009\)](#). The programme ran for two years and included more than 450 Ap stars in total with  $\delta > -30^\circ$ . Thus the majority of observed Ap stars are located in the Northern hemisphere ensuring complementarity with *TESS* Cycle 2 observations.

### 3.3 Analysis

Data from both LAMOST and HERMES were normalised using the SUPPNET software ([Rózański et al. 2022](#)) with the default settings. Where we provide spectral classifications, comparison is made between the observed spectrum and a set of MK Standard stars. We aimed to find a match between the Balmer line profiles and the non-peculiar metals to determine the broad spectral type, then note those peculiar lines that are particularly abnormal in the observed spectrum.

The high resolution HERMES spectra were also analysed to provide estimates of the effective temperature ( $T_{\text{eff}}$ ), surface gravity ( $\log g$ ) and the projected rotational velocity ( $v \sin i$ ). Two separate teams carried out this task, with their procedures below. We combine their results in the relevant star section.

The first team used the software packages `iSPEC` ([Blanco-Cuaresma et al. 2014](#); [Blanco-Cuaresma 2019](#)) and the extended version of `HANDY`<sup>4</sup> ([Rózański](#), private communication). Firstly, with `iSPEC`, Balmer line profiles were fitted and an estimate of the effective temperature with spectral synthesis was made using the `SYNTHE` ([Kurucz 1993](#); [Sbordone et al. 2004](#)) programme with the Castelli and Kurucz model atmospheres ([Castelli & Kurucz 2003](#)) and the VALD line list ([Kupka et al. 2011](#)). This initial result was then used in `HANDY-ext` where synthetic Balmer lines were calculated using the same underlying codes/inputs to determine  $\log g$ ,  $v_{\text{mic}}$ ,  $T_{\text{eff}}$  and  $v \sin i$  via manually adjusting the parameters until a good (visual) match with the Balmer lines was obtained. Those parameters were then used to calculate a second, full, synthetic spectrum in `iSPEC` with which non-blended lines of Fe I and II were found so that  $\log g$ ,  $v \sin i$  and  $T_{\text{eff}}$  were re-calculated, via spectral synthesis, with error estimates.

The second team used the ‘Balmer’ code, written in PYTHON, based on the spectrum synthesis method. The atmospheric models and synthetic spectra were calculated with the `ATLAS9/SYNTHE` software ([Kurucz 2005](#)). The determination of  $T_{\text{eff}}$  and  $\log g$  was performed using four Balmer lines ( $H\alpha$ ,  $H\beta$ ,  $H\gamma$ ,  $H\delta$ ). In the case of temperatures lower than about 8500 K, the Balmer lines become insensitive to  $\log g$ , and only the temperature was determined, with  $\log g$  assumed as  $4.0 \text{ (cm s}^{-2}\text{)}$ . Microturbulence was assumed to be  $\xi = 1 \text{ km s}^{-1}$ . An approximate value of  $v \sin i$  was determined from

the Mg 4481 Å line and metal lines. Errors of estimated parameters were calculated as a weighted median standard deviation of the fit to each line considered.

## 4 RESULTS

Here we present the results of our search and analysis of new and known roAp stars in *TESS* Cycle 2 data. We provide an overview of the targets to be discussed in Table 1 and divide this section, and the table, into subsections addressing new discoveries reported in this work, previous discoveries reliant on the *TESS* data, and *TESS* observations of known roAp stars. Finally, we provide a list of candidate roAp stars where the currently available data do not allow us to confirm the nature of these stars.

As with [Paper I](#), we present the numerical results of the reference team to provide a homogeneous dataset both within this paper and across the two papers. The results have been verified with the WG4 members, and discrepancies were resolved with secondary checks.

Furthermore, we use the same approach as in [Paper I](#) which briefly comprises of: for each star, we combined available sectors to produce a single light curve for analysis. We calculated a Discrete Fourier Transform, following [Kurtz \(1985\)](#), using the `PDC_SAP` data obtained from the MAST servers. Initially we searched in the low-frequency range (up to 0.11 mHz;  $10 \text{ d}^{-1}$ ) to identify signs of rotational variability. Where present, we fitted a harmonic series by non-linear least squares to determine the rotation frequency, and used the highest amplitude peak to determine the rotation frequency. Where no rotational variability was found, we also checked the `SAP` light curve for signs of rotation as the `SPOC` pipeline can remove low-frequency astrophysical signal.

Subsequently, we removed low-frequency peaks (rotational or instrumental) via iterative pre-whitening of the light curve to a noise level determined in the range 2.3 – 3.3 mHz ( $200 - 300 \text{ d}^{-1}$ ). A new amplitude spectrum was calculated to the Nyquist frequency, and any signals found were extracted through non-linear least squares fitting. Both the derived rotation period and the pulsation frequency(ies) for each star are listed in Table 1. We provide only the *pulsation mode* frequencies and not rotationally split sidelobes; any sidelobes detected are mentioned in the text, and indicated on the appropriate figure. Where multiplets due to oblique pulsation exist, we apply the relations of [Kurtz et al. \(1990\)](#) to investigate the geometry of the pulsation mode as derived by the multiplet amplitudes. For a dipole mode, the multiplet amplitudes give a relation between the inclination angle,  $i$ , and the angle of obliquity,  $\beta$ , that is the angle between the rotation and pulsation axes. When conducting these tests we set a zero point in time for the linear least squares fit such that the phases of the  $\pm v_{\text{rot}}$  sidelobes are equal. It is expected that all peaks in a multiplet have equal phase for a non-distorted mode, thus applying the oblique pulsator model provides insight to the mode structure.

For stars that have previously been discussed at length in the literature, with use of the *TESS* data, we do not present an in-depth analysis unless significant differences in the results were found. We do, however, include these stars in Table 1 to provide a complete inventory of *TESS* observations of roAp stars in Cycle 2. For all stars newly classified roAp stars in this work, and roAp stars that have been discovered in *TESS* data, we provide plots of their pulsation frequencies. For the known roAp stars, we provide plots in Appendix C unless there is new information in the *TESS* data.

Where available, we provide the Strömgren-Crawford indices for a given object. We used these values, in conjunction with the cal-

<sup>3</sup> <http://www.mercator.iac.es/instruments/hermes/>

<sup>4</sup> <https://github.com/RozanskiT/HANDY>

**Table 1.** Details of the stars analysed in this paper. The columns provide the TIC identifier and star name, the TIC v8.1 *TESS* magnitude, the spectral type as referenced in the stars discussion section, where ‘\*’ denotes a classification derived in this work, the effective temperature as provided in the TIC, the sectors in which *TESS* observed the target in Cycle 2. The final three columns provide the stellar rotation period derived in this work, with a † indicating a more precise period is presented in the literature and a ‡ indicating this could be an orbital period, the pulsation frequency(ies) found in this work (note that sidelobes that arise from oblique pulsation are not listed), and the pulsation amplitude(s) seen in the *TESS* data. A ‘\*\*\*’ denotes rotation periods derived from rotationally split sidelobes. A ‘\*\*\*\*’ denotes a tentative detection.

TIC	HD/TYC name	<i>TESS</i> mag	Spectral type	$T_{\text{eff}}^{\text{TIC}}$ (K)	Sectors	$P_{\text{rot}}$ (d)	Pulsation frequency (mHz)	Pulsation amplitude (mmag)
<i>New TESS roAp stars found in this work</i>								
101624823	100598	8.18	Ap Cr(SrEu)*	6780	22	$4.050 \pm 0.001$	$1.44176 \pm 0.00001$	$0.124 \pm 0.006$
165052884	51561	8.81	A5	6900	20	~ 22	$2.176721 \pm 0.000007$	$0.243 \pm 0.008$
233200244	161846	8.81	A3	8220	14-19, 21-26	$7.25002 \pm 0.00006$	$1.2614617 \pm 0.0000016$ $1.4390374 \pm 0.0000005$	$0.030 \pm 0.003$ $0.0905 \pm 0.003$
259017938	210684	7.15	F0	7570	15,16	$5.0226 \pm 0.0005$	$1.352116 \pm 0.000006$	$0.064 \pm 0.003$
335457083	48409	8.20	A3	7510	20	$2.9139 \pm 0.0001$	$2.08819 \pm 0.00002$ $2.20575 \pm 0.00003$ $2.24835 \pm 0.00003$ $2.29691 \pm 0.00005$	< 0.015 $0.049 \pm 0.006$ $0.043 \pm 0.006$ $0.029 \pm 0.006$
445493624	11948	7.7	A5 SrEu(Cr)*	8060	18	$7.288 \pm 0.002$	$1.22522 \pm 0.00003$	$0.06 \pm 0.01$
467074220	40142	7.55	A8 SrCr(Eu)*	7410	19	$5.4195 \pm 0.0004$	$0.70462 \pm 0.00006$ $0.72476 \pm 0.00003$	$0.030 \pm 0.007$ $0.059 \pm 0.007$
<i>roAp stars previously discovered by TESS</i>								
21024812	14522	8.59	A8 SrEu(Cr)*	8070	18	~ 24	$1.10259 \pm 0.00002$	$0.0963 \pm 0.0074$
25676603	2194-2347-1	8.50	F0 SrEu	6680	15	>26	$1.787321 \pm 0.000027$ $1.819500 \pm 0.000002$ $1.846724 \pm 0.000008$	$0.072 \pm 0.008$ $0.864 \pm 0.008$ $0.249 \pm 0.008$
26833276	10682	7.51	A3	6970	17	>25	$1.58317 \pm 0.00002$ $1.60714 \pm 0.00003$ $2.74058 \pm 0.00004$ $2.80741 \pm 0.00004$	$0.081 \pm 0.005$ $0.046 \pm 0.005$ $0.033 \pm 0.005$ $0.032 \pm 0.005$
72392575	225578	8.83	A Sr:	7570	14	$3.9016 \pm 0.0008$	$1.57343 \pm 0.00003$	$0.09 \pm 0.01$
118247716	12519	8.13	A7 SrEuCr*	7060	17	No signature	$1.94247 \pm 0.00004$ $1.97093 \pm 0.00002$ $2.03280 \pm 0.00002$	$0.040 \pm 0.006$ $0.098 \pm 0.006$ $0.070 \pm 0.006$
120532285	213258	7.40	Fp	7210	16	No signature	$2.171017 \pm 0.000011$ $2.198746 \pm 0.000008$ $2.226270 \pm 0.000006$	$0.095 \pm 0.004$ $0.128 \pm 0.004$ $0.159 \pm 0.004$
129820552	2322-1440-1	9.90	Ap	7400	18	$6.230 \pm 0.002$	$2.01305 \pm 0.00002$	$0.23 \pm 0.01$
394860395	2666-85-1	9.23		7170	14	$22 \pm 2^{**}$	$1.18050 \pm 0.00005$	$0.09 \pm 0.01$
435263600	218439	7.59	A5 SrSi		17, 24	$3.0739 \pm 0.0002$	$1.097186 \pm 0.000003$ $1.097049 \pm 0.000003$ $1.117976 \pm 0.000003$ $1.124019 \pm 0.000003$	$0.063 \pm 0.010$ $0.059 \pm 0.010$ $0.039 \pm 0.005$ $0.037 \pm 0.005$

ibrations of Crawford (1975, 1979) to determine if the values are indicative of a chemically peculiar star.

#### 4.1 New *TESS* roAp stars

We classified the following stars as new roAp stars based on the presence of high frequency pulsation in the target and either an Ap spectral classification, the presence of multiplets in the amplitude spectrum split by the stellar rotation frequency, or Strömgren-Crawford indices that indicate a chemically peculiar star.

##### 4.1.1 TIC 101624823

TIC 101624823 (HD 100598) has a spectral spectral classification in the literature of A2 (Cannon & Pickering 1919). Masana et al. (2006) estimate the effective temperature to 6800 K, based on *V* and 2MASS IR photometry, and Stassun et al. (2019) estimate  $\log g = 3.8$  ( $\text{cm s}^{-2}$ ). The surface gravity was not obtained via spec-

troscopy, but calculated from the reported mass and radius. In addition, the value for the effective temperature of 7000 K and for the surface gravity  $\log g = 3.85$  ( $\text{cm s}^{-2}$ ) are extracted from the Gaia Collaboration (2022) *Gaia* DR3 catalogue. These values are extracted from GSP-Phot Aeneas best library using BP/RP spectra, apparent G magnitude and parallax. The final result is the median of the MCMC samples. The literature values of radial velocity for this star are  $-15.45 \pm 0.33$   $\text{km s}^{-1}$  measured by Gaia Collaboration (2018), with a more recent value from the Gaia Collaboration (2022) *Gaia* DR3 catalogue of  $-15.64 \pm 0.19$   $\text{km s}^{-1}$ , thus both in agreement. This star appears in the Strömgren-Crawford measurements of Paunzen (2015) with values of  $b - y = 0.223$ ,  $m_1 = 0.157$ ,  $c_1 = 0.672$ ,  $H\beta = 2.714$ , which give  $\delta m_1 = 0.018$  and  $\delta c_1 = 0.105$ . Both of these values indicate a chemically normal star, given the  $H\beta$  value.

*TESS* data for TIC 101624823 used in this work were collected during sector 22. The data show a clear rotational signature of  $4.050 \pm 0.001$  d (Fig. 1). This is in agreement with the value reported by Balona (2022) using the same data, although the previous author did

Table 1 – *continued*

TIC	HD/KIC/TYC name	TESS mag	Spectral type	$T_{\text{eff}}^{\text{TIC}}$ (K)	Sectors	$P_{\text{rot}}$ (d)	Pulsation frequency (mHz)	Pulsation amplitude (mmag)
<i>Known roAp stars prior to TESS launch</i>								
26418690	K11296437	11.38	A9 EuCr	7010	14,15	$7.11 \pm 0.02$	$1.409717 \pm 0.000007$	$0.32 \pm 0.02$
26749633	K11031749	12.39	F1 SrCrEu	7000	14,15	No signature	No signal in TESS data	<0.18
27395746	K11409673	12.58	A9 SrEu	7500	14,15	No signature	No signal in TESS data	<0.15
77128654	97127	9.06	F3 SrEu(Cr)	6710	22	No signature	$1.233855 \pm 0.000004$ $1.248661 \pm 0.000031$	$0.489 \pm 0.009$ $0.061 \pm 0.009$
123231021	K7582608	11.43	A7 EuCr	8400	14,26	$19.805 \pm 0.005^{\ddagger}$	$2.1034011 \pm 0.0000002$	$1.32 \pm 0.02$
158216369	K7018170	12.89	F2 (SrCr)Eu	6950	14,26	No signature	No signal in TESS data	<0.28
158271090	K10195926	10.31	F0 Sr	7390	14,15,26	$5.6845 \pm 0.0002$	$0.974579 \pm 0.000002$	$0.019 \pm 0.009$
158275114	K8677585	10.00	A5 p	7460	14,26	No signature	$1.5853813 \pm 0.0000009$ $1.6217146 \pm 0.0000019$	$0.107 \pm 0.009$ $0.053 \pm 0.009$
159392323	3547-2692-1	11.92	F3 SrEuCr	6390	15	No signature	$1.47658 \pm 0.00005$ $1.49021 \pm 0.00002$ $1.49610 \pm 0.00003$	$0.21 \pm 0.05$ $0.66 \pm 0.05$ $0.34 \pm 0.05$
169078762	225914	9.00	A5 SrCrEu	8100	14	$5.181 \pm 0.003^{\ddagger}$	$0.71130 \pm 0.00005$	$0.05 \pm 0.01$
171988782	258048	10.02	F4 EuCr(Sr)	6370	20	No signature	$1.962238 \pm 0.000005$ $1.975095 \pm 0.000023$	$0.55 \pm 0.01$ $0.13 \pm 0.01$
264509538	K10685175	11.79	A4 Eu	7970	14,15	$3.1029 \pm 0.0005^{\ddagger}$	$2.216610 \pm 0.000006$	$0.57 \pm 0.03$
272598185	K10483436	11.16	A7 SrEuCr	7980	14,15	$4.2857 \pm 0.0006$	No signal in TESS data	<0.05
273777265	K6631188	13.38	F0 Sr	7500	14,15	$2.519 \pm 0.002^{\ddagger}$	$1.49349 \pm 0.00001$	$1.45 \pm 0.16$
286992225	2553-480-1	11.24	A9 SrEu	7490	23	No signature	$2.69128 \pm 0.00002$ $2.72616 \pm 0.00001$	$0.32 \pm 0.03$ $0.54 \pm 0.03$
302602874	2488-1241-1	10.86	A6 SrEu	8160	21	$3.0900 \pm 0.0003$	$2.283125 \pm 0.000002$	$2.12 \pm 0.02$
<i>Candidate roAp</i>								
9171107	2479-429-1	9.36	Unknown	6770	20	No signature	$1.50971 \pm 0.00004$ $1.52902 \pm 0.00002$ $1.53502 \pm 0.00002$	$0.053 \pm 0.009$ $0.108 \pm 0.009$ $0.094 \pm 0.009$
91224991	191380	8.49	F8	6710	14	No signature	$1.159876 \pm 0.000002$	$0.961 \pm 0.0081$
275642037	4330-716-1	9.52	F0	7710	18	$2.8745 \pm 0.0003$	$1.13943 \pm 0.00003$ $1.16939 \pm 0.00002$	$0.08 \pm 0.01$ $0.17 \pm 0.01$
298052991	4554-625-1	8.8	A2	7600	14, 21, 21, 26	$10.5852 \pm 0.0001$	$2.217569 \pm 0.000003$ $2.246466 \pm 0.000002$ $2.291038 \pm 0.000001$	$0.023 \pm 0.004$ $0.038 \pm 0.004$ $0.051 \pm 0.004$
405892692	3567-1092-1	8.99	F0	7230	14, 15	$0.3786 \pm 0.0001^{\dagger}$	$1.17082 \pm 0.00002$	$0.038 \pm 0.006$
429251527	2185-478-1	10.20		7920	15	No signature	$1.77798 \pm 0.00001$	$0.45 \pm 0.02$
445796153	34740	7.22	A0p	8510	19, 26	$5.5721 \pm 0.0004$	$0.601908 \pm 0.000002$ $0.675849 \pm 0.000002$	$0.022 \pm 0.003$ $0.035 \pm 0.003$

not include an error on the measurement. After removing the rotation signal from the data, we analysed the high frequency peak ( $1.44176 \pm 0.00001$  mHz;  $124.5678 \pm 0.0009$  d<sup>-1</sup>), which was also noted by Balona (2022) but the star was not admitted to the roAp star class in that work. Here we rectify that oversight with the identification of a rotationally split sidelobe at  $\nu - \nu_{\text{rot}}$  (at  $S/N \sim 4.5$ ) as is expected from oblique pulsations in roAp stars, which is shown in Fig. 2 by the vertical dotted line. A second sidelobe is identifiable (at  $\nu + \nu_{\text{rot}}$ ), albeit at low  $S/N$  ( $\sim 3.1$ ), indicating the presence of a triplet, and a dipole mode.

With these two sidelobes, we are able to apply the oblique pulsator mode to derive  $\tan i \tan \beta = 0.369 \pm 0.071$ . We also found that the mode is slightly distorted when investigating the phase relations between the mode and the sidelobes. We found that pulsation maximum occurs at rotation phase 0.46 in Fig. 1, hence the pulsation pole coincides with the principal abundance spot that gives rise to the rotational light variations. This spot in turn is likely to coincide with, or be close to, a magnetic pole. Hence this star is typical of the roAp star oblique pulsators in that the pulsation pole lies close to the magnetic pole.





Figure 1. Phase folded light curve of TIC 101624823, phased on a period of  $4.050 \pm 0.001$  d. Binned 50:1.

To further cement its membership of the roAp class, we have obtained a spectrum from LAMOST. While the data are of low  $S/N$ , there is clear presence of Cr II in the spectrum (Fig. B6), with the possible presence of Sr II and Eu II, but at a lower confidence. Therefore, we classified this star as an ApCr(SrEu) star. This shows



**Figure 2.** Amplitude spectra of TIC 101624823. The pulsation mode is identified by the vertical dotted line, with the dashed lines showing the locations of the rotationally split sidelobes.

that mild Ap stars, as classified by their Strömgren-Crawford indices can be roAp stars.

#### 4.1.2 TIC 165052884

TIC 165052884 (HD 51561) has a spectral classification of A5 (Cannon & Pickering 1918b). Stassun et al. (2019) estimate  $T_{\text{eff}} = 6900$  K and  $\log g = 4.18$  ( $\text{cm s}^{-2}$ ). The effective temperature is derived from dereddened *Gaia* Bp-Rp colour. As with the previous star, this value is calculated using the mass and radius estimates. The *Gaia* DR3 (Gaia Collaboration 2022) gives  $T_{\text{eff}} = 6850$  K and  $\log g = 4.22$  ( $\text{cm s}^{-2}$ ).


*TESS* observed TIC 165052884 in sector 20 of Cycle 2, with the data analysed in Balona (2022) who measured a rotation period (using sector 20 and 44 data) of  $P_{\text{rot}} = 17.857$  d and a pulsation frequency of  $2.177$  mHz ( $\nu = 188.069$   $\text{d}^{-1}$ ). However, our analysis cannot reproduce the rotation results of Balona (2022), even if we include the sector 44 data. We further investigated this discrepancy by analysing the SAP data, which does indeed show variability that can be attributed to rotation. However, the signal indicated a rotation period about the same length of the data ( $P_{\text{rot}} \sim 22$  d compared to a data set length of  $\sim 26$  d), so cannot be precisely determined.

The high frequency pulsation mode in TIC 165052884 is found at  $2.176721 \pm 0.000007$  mHz ( $188.0687 \pm 0.0006$   $\text{d}^{-1}$ ; Fig. 3), which is in agreement with Balona (2022). After pre-whitening this signal, there is an indication of a much lower amplitude mode at  $2.17566 \pm 0.00004$  mHz ( $187.977 \pm 0.003$   $\text{d}^{-1}$ ) which is separated by the principal peak by a frequency consistent with half of the presumed rotation frequency ( $\sim 1.06$   $\mu\text{Hz}$ ;  $0.09$   $\text{d}^{-1}$ ;  $10.9$  d). Again, more data are required to confirm this detection.


We admitted TIC 165052884 to the roAp class based on the pulsation and the tentative detection of a rotational sidelobe to the pulsation mode.

#### 4.1.3 TIC 233200244

TIC 233200244 (HD 161846) was classified as A3 in the Henry Draper Catalogue (Cannon & Pickering 1922). From photometry, Pickles & Depagne (2010) give a similar spectral type of A5 V. The TIC gives  $T_{\text{eff}} = 8200 \pm 150$  K, with similar values of 8050 K and 8100 K obtained by McDonald et al. (2017) and Tonry et al. (2018), respectively, while the *Gaia* DR3 lists  $T_{\text{eff}}$  as 8000 K. The star was analysed by Balona (2022) who classified it as a new variable A star, with a rotation period of  $P_{\text{rot}} = 7.299$  d (without an error) and a pulsation frequency of  $1.44$  mHz ( $124.471$   $\text{d}^{-1}$ ), which also



**Figure 3.** Amplitude spectrum of TIC 165052884 with the pulsation mode identified with the vertical dotted line.



**Figure 4.** Phase folded light curve of TIC 233200244, phased on a period of  $7.25002 \pm 0.00006$  d. Binned 200:1.


showed rotational sidelobes. The star also appears in the catalogue by Skarka et al. (2022) who classified it as a rotationally modulated star, with a period of  $7.246$  d, and a candidate roAp star after studying all A and F stars in *TESS*'s northern continuous viewing zone.

*TESS* observed the star in all sectors of Cycle 2 except sector 20. This allows us to derive a precise rotation period of  $7.25002 \pm 0.00006$  d. A folded light curve is shown in Fig. 4. We proceeded to prewhiten this signal from the individual sectors before analysing the pulsation modes in this star, with the resultant amplitude spectrum shown in Fig. 5. We found two pulsation modes at  $1.261462 \pm 0.000002$  mHz ( $108.9903 \pm 0.0001$   $\text{d}^{-1}$ ) and  $1.4390374 \pm 0.0000005$  mHz ( $124.33283 \pm 0.00005$   $\text{d}^{-1}$ ) separated by  $177.576 \pm 0.002$  mHz, which could be three times the large frequency separation<sup>5</sup> ( $\sim 60$   $\mu\text{Hz}$ ), when scaling from the solar value.


The higher frequency mode is flanked by rotational sidelobes split by exactly the rotation frequency, indicating the mode is an  $\ell = 1$  dipole mode. Given their presence, we applied the oblique pulsator model to test the phase relations and determine whether the mode is distorted. We found there is a little distortion to the mode, with the sidelobes having a phase of  $-0.553 \pm 0.030$  rad and the pulsation mode having  $-0.910 \pm 0.031$  rad. The time zero point for this calculation corresponds to rotation phase 0.68 in Fig. 4, i.e., at *TESS* light maximum.

Given the detection of the pulsations and the sidelobes, coupled with a clearly stable rotational light curve, we admitted this star to the class of roAp stars.

<sup>5</sup> defined as the frequency difference between two consecutive modes of the same degree



**Figure 5.** Amplitude spectrum of TIC 233200244 showing the two pulsation modes (marked by the vertical dotted lines) and the rotationally split sidelobes of the higher frequency mode.




**Figure 6.** Phase folded light curve of TIC 259017938, phased on a period of  $5.0226 \pm 0.0005$  d. Binned 100:1.

#### 4.1.4 TIC 259017938

TIC 259017938 (HD 210684) has been classified F0 (Cannon & Pickering 1924). Kervella et al. (2019) revealed the existence of a physical companion orbiting TIC 259017938 from the influence on the star’s proper motion. Assuming  $M_1 = 2.03 \pm 0.10 M_\odot$  (Girardi et al. 2000) and  $R_1 = 2.66 \pm 0.13 R_\odot$  (Kervella et al. 2004), they found the mass of the companion and the radius of its orbit to be  $M_2 = 80_{10}^{23} M_{\text{Jup}}$  and 1.892 au, respectively. Strömgren-Crawford indices are given by Paunzen (2015) as:  $b - y = 0.160$ ,  $m_1 = 0.196$ ,  $c_1 = 0.824$  and  $H\beta = 2.570$ . We derive  $\delta m_1 = 0.066$  and  $\delta c_1 = 0.619$ . These do not indicate an Ap star, however we note that the  $b - y$  and  $H\beta$  indices do not agree. Paunzen (2015) combined two consistent values of  $b - y$ , while only reporting one value of  $H\beta$ . When using  $b - y$  as a proxy for  $H\beta$  (corresponding to 2.786) we found  $\delta m_1 = 0.002$  and  $\delta c_1 = 0.070$  which do not change our conclusion.

*TESS* observed TIC 259017938 during sectors 15 and 16 and revealed the star to be a rotationally and pulsationally variable star (Balona 2022). We determined a rotation period of  $5.0226 \pm 0.0005$  d (Fig. 6), which differs from the previously published value, although that value has no associated error. After removing the signature of rotation, we identified a single pulsation mode in the star at  $1.352116 \pm 0.000006$  mHz ( $116.8228 \pm 0.0005$  d $^{-1}$ ) that showed a rotationally split multiplet (Fig. 7). We again applied the oblique pulsator model to this dipole triplet to investigate the mode distortion and found the phase of the pulsation mode ( $-2.48 \pm 0.05$  rad) differs from the sidelobes ( $-2.17 \pm 0.09$  rad) by  $3\sigma$ , indicating possible slight mode distortion. The pulsation maximum corresponds to rotation phase 0.21 in Fig. 6, i.e., just before *TESS* light maximum.




**Figure 7.** Amplitude spectrum of TIC 259017938 showing the pulsation mode (marked by the vertical dotted line) and the rotationally split sidelobes.

We have also obtained three medium resolution spectra from the LAMOST database to search for chemical peculiarities (Fig. B4). We identified lines of Pr III, Nd II & III and La II amongst others, which indicate the star to be an Ap star, thus providing further evidence to admit the star to the roAp class.


#### 4.1.5 TIC 335457083

TIC 335457083 (HD 48409), was classified as A3 in the Henry Draper Catalogue (Cannon & Pickering 1918b). The TIC gives  $T_{\text{eff}} = 7500 \pm 140$  K and *Gaia* DR3 7500 K with a small uncertainty. The literature values, however, range from 9100 K (Stassun et al. 2018) down to 6900 K (Schofield et al. 2019), with intermediate values of 8700 K, 8750 K and 7650 K (Wright et al. 2003; Ammons et al. 2006; McDonald et al. 2017), respectively. Based on its Strömgren-Crawford indices, of  $b - y = 0.213$ ,  $m_1 = 0.308$ ,  $c_1 = 0.517$  and  $H\beta = 2.838$  (Hauck & Mermilliod 1998), which give  $\delta m_1 = -0.100$  and  $\delta c_1 = -0.332$ , and a large  $\Delta p$  “peculiarity” parameter of 6.779 (defined by Masana et al. (1998)), the star was selected to be observed by the Nainital-Cape Survey (Joshi et al. 2006), but no variability was found in this targeted ground-based photometric study. Previously, Balona (2022) reported a rotation period of 2.907 d and a pulsation frequency of 2.084 mHz ( $180.075$  d $^{-1}$ ) in this star using the *TESS* data.


*TESS* observed TIC 335457083 in sector 20 from which we derived a rotation period of  $2.9139 \pm 0.0001$  d (Fig. 8). At high frequencies, we inferred the presence of four pulsation modes (Fig. 9). In increasing frequency order, we found a doublet and inferred the presence a mode from the observation of two peaks separated by twice the rotation frequency (i.e., a mode at  $2.08819 \pm 0.00002$  mHz;  $180.419 \pm 0.002$  d $^{-1}$ ), a triplet (centred on a mode at  $2.20575 \pm 0.00003$  mHz;  $190.577 \pm 0.002$  d $^{-1}$ ), followed by two single peaks at  $2.24835 \pm 0.00003$  mHz ( $194.258 \pm 0.003$  d $^{-1}$ ) and  $2.29691 \pm 0.00005$  mHz ( $198.453 \pm 0.004$  d $^{-1}$ ). The doublet and triplet show very different ratios of the pulsation mode amplitude and the amplitude of the rotationally split sidelobes. For both modes to be dipole modes, we would require two pulsation axes in this star. Therefore, we assumed the doublet to be a quadrupole mode, and the triplet to be a dipole mode. Under this assumption, we calculated the values of  $i$  and  $\beta$  following Kurtz et al. (1990) such that  $i = 56^\circ$  and  $\beta = 45^\circ$ . This leads to  $i + \beta = 100^\circ$  which is consistent with the partial double wave of the rotation curve, supporting our choice of mode identifications. Of the observed, or inferred modes, there is no regular spacing that can be interpreted as the large frequency separation. However, there are further peaks above the noise in the amplitude spectrum of this star too, but further data are required to reduce the noise level to




**Figure 8.** Phase folded SAP light curve of TIC 335457083, phased on a period of  $2.9139 \pm 0.0001$  d. Binned 50:1.



**Figure 10.** Phase folded SAP light curve of TIC 445493624, phased on a period of  $7.288 \pm 0.002$  d. Binned 50:1.



**Figure 9.** Amplitude spectrum of TIC 335457083 showing the pulsation modes (marked by the vertical dotted lines) and the rotationally split sidelobes (marked with the vertical dashed lines).



**Figure 11.** Amplitude spectrum of TIC 445493624. We have extracted the pulsation mode marked with the vertical dotted line. The other variability clustered around 1.235 mHz is unresolved in this short set of data.

allow confident extraction of those possible mode frequencies. These additional modes may lead to the identification of the large separation frequency in this star.

Given both the strong indication that this star is chemically peculiar from the Strömgren-Crawford photometry, and the presence of modes with rotational splitting, we admitted this star to the class of roAp stars.

#### 4.1.6 TIC 445493624

TIC 445493624 (HD 11948) has several spectral classifications in the literature: F0p (Hoag & Applequist 1965), A5 Am/p (Olsen 1980), A5 SrEu (Renson 1988), A5 SrCrEu (Gomez et al. 1998), all of which indicate this to be a chemically peculiar star. In addition, Strömgren-Crawford indices have also been measured several times (e.g., Perry & Johnston 1982) with values being consistent with:  $b - y = 0.115$ ,  $m_1 = 0.243$ ,  $c_1 = 0.880$  and  $H\beta = 2.873$  (Paunzen 2015), which are also consistent with a chemically peculiar A star (with  $\delta m_1 = -0.041$  and  $\delta c_1 = -0.036$ ). The effective temperature of the star is reported consistently across the literature, e.g.,  $T_{\text{eff}} = 8050$  K (McDonald et al. 2012),  $T_{\text{eff}} = 8250$  K (Joshi et al. 2016) and the TIC providing 8050 K. Kervella et al. (2019) report a mass and radius of  $M = 2.20 M_{\odot}$  and  $R = 2.15 R_{\odot}$ . The *Gaia* DR3 parallax provides a distance of  $180 \pm 1$  pc, which is at odds to the report that TIC 445493624 is a member of the NGC 744 open cluster (Niedzielski & Muciek 1988).


The star has a constant radial velocity in the literature of  $32 \text{ km s}^{-1}$  (de Bruijne & Eilers 2012, Kervella et al. 2019, Brandt 2021), and has a root-mean-squared longitudinal magnetic field of  $\langle B_{\ell} \rangle = 500 \pm$

50 G with extreme values of the longitudinal component being  $B_{\ell} - 550$  and  $+300$  G (Romanyuk & Kudryavtsev 2008).

The star was searched for pulsations during the Nainital-Cape survey (Joshi et al. 2006), but none was found. Since then the *TESS* data have been used by Balona (2022) to identify a rotation period of 7.333 d, and a pulsation frequency of 1.225 mHz ( $105.857 \text{ d}^{-1}$ ). In that study, the star was classified as a possible roAp star.

*TESS* observed TIC 445493624 during sector 18. To derive the rotation period, we cleaned the SAP data of outlying points (which constituted about 5 per cent of the data) since the signal in the PDC\_SAP data had been altered by the pipeline corrections. We derived a rotation period of  $P_{\text{rot}} = 7.288 \pm 0.002$  d, as shown in Fig. 10. To inspect the high frequency variability, we returned to the PDC\_SAP data and removed the low-frequency peaks to the noise level in the frequency range of interest. Here we found significant evidence of variability (Fig. 11). There are many peaks with amplitudes significantly above the noise level, however many are unresolved. We extracted a single mode, which itself does not resemble the window function, at a frequency of  $1.22522 \pm 0.00003$  mHz ( $105.859 \pm 0.002 \text{ d}^{-1}$ ). Since the frequency resolution for the data set is  $0.4 \mu\text{Hz}$ , which is significantly smaller than the expected small separation in roAp stars, the evidence suggests amplitude and/or frequency variability of the mode. This is not unprecedented in the Ap stars, even on short time scales, (e.g., HD 60435; Balona et al. 2019).

To confirm the chemical peculiarity in this star, we have obtained a spectrum using the HERMES spectrograph (Figs. B1 & B2). We found clear presence of Sr II and Eu II, as well as enhancements of other rare earth elements in the region around  $6150 \text{ \AA}$ . Using the procedures outlined above, we derived the following parameters from



**Figure 12.** Phase folded light curve of TIC 467074220, phased on a period of  $5.4195 \pm 0.0004$  d. Binned 50:1.

the spectrum:  $T_{\text{eff}} = 8370 \pm 90$  K,  $\log g = 4.0 \pm 0.3$  ( $\text{cm s}^{-2}$ ) and  $v \sin i = 14 \pm 2$   $\text{km s}^{-1}$ .

With the evidence we have gathered, and that in the literature, we admitted TIC 445493624 to the class of roAp stars.

#### 4.1.7 TIC 467074220

TIC 467074220 (HD 40142) has several classifications in the literature: Ap Sr-Eu (Cowley & Cowley 1965); F0p (Nelson & Kreidl 1993); F0 (Cannon & Pickering 1918b). Its estimated effective temperature is  $T_{\text{eff}} = 7200$  K from *Gaia* DR2 (Gaia Collaboration et al. 2018; Gaia Collaboration 2018) and 6950 K from (McDonald et al. 2012), while Shang et al. (2022) determined 7580 K from LAMOST spectra. *Gaia* DR2 also provided  $\log L_{\odot} = 1.37$ . Hauck & Mermilliod (1998) provide Strömgren–Crawford indices of:  $b - y = 0.216$ ,  $m_1 = 0.175$ ,  $c_1 = 0.869$  and  $H\beta = 2.814$ , giving  $\delta m_1 = 0.031$  and  $\delta c_1 = 0.064$ , indicating a normal star.


This star has a magnetic field first reported by (Romanyuk & Kudryavtsev 2008) with a root mean value of  $\langle B_{\ell} \rangle = 700 \pm 50$  G with 3 magnetic field observations from the 6-m telescope of the Special Astrophysical Observatory of the Russian Academy of Sciences. The extremes measured were  $-780$  G and  $+780$  G.

Pulsations have been searched for in this star before by Nelson & Kreidl (1993) using ground-based *V* and *B* photometric data, but with none found. No other searches have been published.

*TESS* observed this star during sector 19. The data show a clear rotation signature with a period of  $P_{\text{rot}} = 5.4195 \pm 0.0004$  d (Fig. 12). This is the first reporting of this signature in the literature. Further, we detected two pulsation modes (Fig. 13) at  $0.70462 \pm 0.00006$  mHz ( $60.879 \pm 0.005$   $\text{d}^{-1}$ ) and  $0.72476 \pm 0.00003$  mHz ( $62.619 \pm 0.002$   $\text{d}^{-1}$ ). These modes are similar in frequency to  $\delta$  Sct modes, but there are only two modes detected which is unusual for  $\delta$  Sct stars. At this slightly higher frequency, there are more examples of roAp stars with similar pulsations in the literature (e.g., HD 177765; Alentiev et al. (2012); Holdsworth (2016)).

We have extracted a spectrum of TIC 467074220 from the LAMOST archive to confirm the spectral classification (Fig. B9). There is obvious over enhancements of Sr II and Cr II, with an indication of Eu II.

With a combination of the rotational modulation in the star, the detected magnetic field and observed chemical peculiarities, we admitted this star to the class of roAp.



**Figure 13.** Amplitude spectrum of TIC 467074220. The two pulsation modes marked with vertical dotted lines.

## 4.2 roAp stars previously discovered by *TESS*

In this section, we present the roAp stars detected in *TESS* data by our method but which have already been published in the literature.

### 4.2.1 TIC 21024812


TIC 21024812 (HD 14522) was originally identified as a CP2 star by Schneider (1986), and Renson & Manfroid (2009) classified it as A2 SrEu. Muñoz Bermejo et al. (2013) estimated the star’s effective temperature using principal component analysis techniques to compare it to a homogeneous library of stellar spectra, and derived  $T_{\text{eff}} = 7750$  K, in general agreement with the TIC value of  $8050 \pm 130$  K (Stassun et al. 2019) and the *Gaia* DR3 value of 7950 K (Gaia Collaboration 2020). *Gaia* DR3 also derived a distance of 282 pc,  $\log g = 3.93$  ( $\text{cm s}^{-2}$ ) and  $[\text{Fe}/\text{H}] = -0.11$ , while the TIC estimated the stellar radius to be  $2.49 \pm 0.08 R_{\odot}$  and mass to be  $1.96 \pm 0.30 M_{\odot}$ .

Bernhard et al. (2020) made use of archival photometric data from a variety of ground-based surveys to estimate the rotational period to be  $26.36 \pm 0.01$  d, with an amplitude in *V* of about 0.01 mag, while Joshi et al. (2006) conducted a 1.4-hr search for pulsations using a three-channel photometer on the ARIES 1.04-m Sampurnanand telescope, but found no variability.

TIC 21024812 was reported as an roAp star by Balona (2022) who derived a rotation period of  $P_{\text{rot}} = 13.158$  d and a pulsation frequency of 1.103 mHz ( $95.265$   $\text{d}^{-1}$ ) using *TESS* sector 18 data. However, our more detailed analysis of the same sector 18 observations suggests that the rotation period is about twice that reported. We used the SAP light curve, which showed significant variability consistent with stellar rotation, to find that the signal is non-repeating over the light curve duration of 24.356 d. This is consistent with the rotation period derived by Bernhard et al. (2020) of  $26.36 \pm 0.01$  d. Further observations of this star are required to constrain the rotation period with precision.

The short period variability seen in the amplitude spectrum of this star (Fig. 14) is seen at a frequency of  $1.10259 \pm 0.00002$  mHz ( $95.263 \pm 0.002$   $\text{d}^{-1}$ ) with the indication of a sidelobe on the higher frequency side of the pulsation. The peak in the amplitude spectrum appears broad which may be caused by amplitude/phase modulation, or the presence of unresolved sidelobes.

Further to the classification of TIC 21024812 being a chemically peculiar star in the literature, we have two sources of spectra of this star: LAMOST and HERMES. The LAMOST spectrum was obtained in its low resolution mode, and can be seen in Fig. B3, while the HERMES spectrum was recorded at a resolution of  $\sim 85000$



**Figure 14.** Amplitude spectrum of TIC 21024812, with the pulsation mode indicated with the vertical dotted line, and the suspected positive rotational sidelobe with the dashed line. The peak is seen to be broad, indicating amplitude/phase modulation, or further unresolved sidelobes.

and is shown in Figs. B1 & B2. From the LAMOST spectrum, we classified this star as A8 SrEu(Cr). Using the procedures outlined above, we derived the following parameters from the HERMES spectrum:  $T_{\text{eff}} = 8250 \pm 70$  K,  $\log g = 4.0 \pm 0.3$  ( $\text{cm s}^{-2}$ ) and  $v \sin i = 8 \pm 3$   $\text{km s}^{-1}$ .

#### 4.2.2 TIC 25676603


TIC 25676603 (TYC 2194-2347-1) is a slowly rotating Fp star, with particularly strong absorption in the Sr II 4077 Å line and  $v \sin i < 1.1$   $\text{km s}^{-1}$  (North & Duquennoy 1991). There are not many mentions of this star in the literature where detailed studies have been conducted. The *Gaia* DR3 gives the effective temperature of  $T_{\text{eff}} = 6650$  K and a  $\log g = 4.1$  ( $\text{cm s}^{-2}$ ). These are consistent with the spectral types of Ap SrEu (Bidelman 1966) and the Fp class.

TIC 25676603 was first reported as an roAp star by Balona (2022) who claimed to identify the rotation period to be  $P_{\text{rot}} = 5.495$  d and a pulsation mode at a frequency of 1.820 mHz ( $157.205$   $\text{d}^{-1}$ ). However, our analysis of the *TESS* sector 15 data do not show a clear signal of rotation. We analysed the SAP data to investigate if a long rotation period has been removed by the PDC pipeline. Indeed, we found some variability that is consistent with a long rotation period, as would be expected from the previous measurement of the upper limit on  $v \sin i$ . We therefore concluded that the rotation period is longer than the current *TESS* data set, and the value published by Balona (2022) should be disregarded as noise or of instrumental origin.


We identified three pulsation modes in the PDC\_SAP sector 15 data for this star, at frequencies of  $1.787321 \pm 0.000027$  mHz ( $154.4245 \pm 0.0024$   $\text{d}^{-1}$ ),  $1.819500 \pm 0.000002$  mHz ( $157.2048 \pm 0.0002$   $\text{d}^{-1}$ ) and  $1.846724 \pm 0.000008$  mHz ( $159.5569 \pm 0.0007$   $\text{d}^{-1}$ ). All three peaks show structure (Fig. 15), suggesting that the rotation period is not fully resolved in 27 d of *TESS* data, confirming our previous conclusion. The three modes are not separated by the same frequency spacing ( $32.18 \pm 0.03$   $\mu\text{Hz}$  and  $27.22 \pm 0.01$   $\mu\text{Hz}$ ), but since the modes are not fully resolved this view may change with the addition of future observations.

#### 4.2.3 TIC 26833276

TIC 26833276 (HD 10682) is not well studied in the literature. The only classification giving an indication that this is a chemically peculiar star is from Clausen & Jensen (1979) with F0 Vp Sr. However, this is at odds with the classification of A3 (Cannon & Pickering



**Figure 15.** Amplitude spectrum of TIC 25676603, with the pulsation modes indicated with the vertical dotted lines.




**Figure 16.** Amplitude spectra of TIC 26833276 calculated from the two *TESS* orbits when it was observed. The amplitude spectrum with positive amplitude is from orbit 41, with two pulsation modes around 1.6 mHz. The amplitude spectrum with negative amplitude is from orbit 42, showing the emergence of two further modes at high frequency (around 2.8 mHz).

1918a). Catalogue values of the effective temperature range from 6750 K (McDonald et al. 2012) to 8700 K (Wright et al. 2003), while *Gaia* DR3 gives 7300 K (Gaia Collaboration 2022). The hottest temperature is based on the spectral type alone, while the others are derived from measurements. Therefore we suggest the effective temperature is closer to 7000 K which supports the classification of an F0 star.

Balona (2022) classified the star as an roAp star given the presence of a pulsation mode at a frequency of 1.583 mHz ( $136.786$   $\text{d}^{-1}$ ), as well as a rotation period of 10.417 d, as seen in the *TESS* sector 17 data. However, a detailed view of the same *TESS* data here does not support the conclusion of the rotation period. This is another case where the PDC pipeline has altered the rotation signal since it appears to be longer than the data set. When inspecting the SAP data, it is clear there is a partial rotation curve that does not repeat. We therefore conclude that the rotation period is greater than the sector length, i.e.,  $>25$  d, and the previous value presented in the literature should be ignored.

The pulsations in this star are irrefutable. There are two groups of two modes; the lower frequency pair appear at  $1.58317 \pm 0.00002$  mHz ( $136.786 \pm 0.001$   $\text{d}^{-1}$ ) and  $1.60714 \pm 0.00003$  mHz ( $138.857 \pm 0.002$   $\text{d}^{-1}$ ) and are present in all of the sector 17 observations. The two modes at higher frequency are transient in nature and only appear in the second orbit of sector 17 (Fig. 16). These modes are at  $2.74058 \pm 0.00004$  mHz ( $236.786 \pm 0.003$   $\text{d}^{-1}$ ) and  $2.80741 \pm 0.00004$  mHz ( $242.560 \pm 0.003$   $\text{d}^{-1}$ ). We are unsure of the nature of the high-frequency pair.

This star has LAMOST MRS data (Fig. B5) available in the archive which we analysed to identify chemical peculiarities and to resolve



**Figure 17.** Phase folded light curve of TIC 72392575, phased on a period of  $3.9016 \pm 0.0008$  d. Binned 50:1.

the discrepancy between the spectral types. We found an overabundance of Nd III and La II, amongst others, confirming the star to be chemically peculiar. By fitting the  $H\alpha$  line, we determine an effective temperature of  $7200 \pm 200$  K. Therefore, we can confirm the star is an F0p chemically peculiar star.

#### 4.2.4 TIC 72392575

TIC 72392575 (HD 225578) has an uncertain spectral classification; it is given as ApSr: by Cowley & Cowley (1964), while it is classified as an A5 star in the Henry Draper Catalog and Extension (Cannon & Pickering 1923). More recently, Renson & Manfroid (2009) gave the spectral classification for this star as A5 Sr.

Sikora et al. (2019) and Stassun et al. (2019) estimate the effective temperature and  $\log g$  values to be 7550 K and  $4.14$  ( $\text{cm s}^{-2}$ ), respectively. The effective temperature is determined from the dereddened Bp-Rp colour, while the  $\log g$  value is derived from the given mass and radius. Other estimations of  $T_{\text{eff}}$  are 7650 K (Ammons et al. 2006) and 7850 K (Anders et al. 2019).


Balona (2022) listed this star as a new roAp star discovered using *TESS* data, and also classified it as an  $\alpha^2$  CVn variable with a rotation period of  $P_{\text{rot}} = 3.922$  d. He also gave an estimate of the luminosity  $\log L/L_{\odot} = 1.07$  using *Gaia* EDR3 parallaxes (Gaia Collaboration et al. 2016, 2021) in conjunction with interstellar reddening corrections from Gontcharov (2017) and a bolometric correction calibration by Pecaut & Mamajek (2013).

A previous search for pulsational variability did not yield a positive detection down to 1.6 mmag in the amplitude spectrum (Paunzen et al. 2012). However, Balona (2022) found roAp variability in the *TESS* light curve, with the highest amplitude frequency at 1.573 mHz ( $135.944 \text{ d}^{-1}$ ). There are no magnetic field measurements for this star in the literature.


Our analysis of the *TESS* light curve, from sector 14, confirms the rotation period as  $3.9016 \pm 0.0008$  d (Fig. 17), and the single high frequency pulsation at  $1.57343 \pm 0.00003$  mHz ( $135.944 \pm 0.003 \text{ d}^{-1}$ ; Fig. 18). Despite the short rotational period of this star, we do not detect rotational sidelobes to the pulsation, suggesting a low value of  $i$  or  $\beta$  for this star. However, there is a slight power excess at slightly higher frequency which may be a signature of a rotational sidelobe, but this is unclear.

#### 4.2.5 TIC 118247716

TIC 118247716 (HD 12519) has several spectral classifications in the literature: A5 (Cannon & Pickering 1918a), A4pSrEuCr (with a comment of ‘composite?’; Clausen & Jensen 1979) and A2pSrSi



**Figure 18.** Amplitude spectrum of TIC 72392575, showing the pulsation mode. At slightly higher frequency there is a power excess which might be a rotational sidelobe.




**Figure 19.** Amplitude spectrum of TIC 118247716, with three modes clearly identifiable. Further modes may be present close to the noise level.

(Grenier et al. 1999). Stassun et al. (2019) estimate  $T_{\text{eff}} = 7050$  K, and  $\log g = 4.04$  ( $\text{cm s}^{-2}$ ). The effective temperature value is derived from a dereddened Bp-Rp colour. The surface gravity is calculated from the reported mass and radius of the star. *Gaia* DR3 estimates an effective temperature of 7100 K and  $\log g = 4.11$  ( $\text{cm s}^{-2}$ ) (Gaia Collaboration 2022). There are two consistent radial velocity estimates for this star:  $-5.9 \pm 3.3 \text{ km s}^{-1}$  (Gontcharov 2006) and more recently  $-8.9 \pm 0.6 \text{ km s}^{-1}$  from the *Gaia* DR3 data (Gaia Collaboration 2022).

This star was targeted as part of the Nainital-Cape survey (Joshi et al. 2006) which searched for photometric variability. The star was included in the target list given its Strömgren-Crawford indices of  $b - y = 0.238$ ,  $m_1 = 0.289$ ,  $c_1 = 0.546$  and  $H\beta = 2.803$  (Paunzen 2015), giving  $\delta m_1 = -0.086$  and  $\delta c_1 = -0.239$ , being indicative of an Ap star, and its  $\Delta p$  peculiarity parameter of 5.989 (Masana et al. 1998).

TIC 118247716 was first classified as an roAp star using *TESS* data (Balona 2022). We corroborate this with the detection of three pulsation modes in the typical roAp frequency range (Fig. 19):  $1.94247 \pm 0.00004$  mHz ( $167.829 \pm 0.003 \text{ d}^{-1}$ ),  $1.97093 \pm 0.00002$  mHz ( $170.2884 \pm 0.001 \text{ d}^{-1}$ ), and  $2.03280 \pm 0.00002$  mHz ( $175.634 \pm 0.002 \text{ d}^{-1}$ ). There appear to be further modes in this star, but close to the noise level. Thus, we did not extract them, but suggest this star is revisited when more data are available. We see no rotationally split sidelobes to the pulsations which is consistent with the lack of rotational modulation in the light curve.

This star has a LAMOST low resolution spectrum available in the archive, which we used to confirm its spectral type of A7pSrEuCr (Fig. B7). This cooler classification is justified by the lack of con-



**Figure 20.** Amplitude spectrum of TIC 120532285

vincing Si lines in the spectrum, which are associated with hotter CP stars, and that the Balmer lines are well matched with that of an A7 normal star. This cooler classification also agrees better with the literature effective temperatures.


#### 4.2.6 TIC 120532285

TIC 120532285 (HD 213258) is a high proper motion star (Gaia Collaboration 2020), with a measured radial velocity of  $-86.8 \text{ km s}^{-1}$  (North & Duquennoy 1991). It is classified as an A3 star by Cannon & Pickering (1924), and an F str  $\lambda 4077$  star<sup>6</sup> by Bidelman (1985). Gaia DR2 estimates an effective temperature in the range 6750 K to 7350 K, a radius range of  $1.75 - 2.09 R_{\odot}$ , and luminosity range of  $8.07-8.12 L_{\odot}$  (Gaia Collaboration et al. 2018). Paunzen (2015) provides Strömgren-Crawford indices of  $b - y = 0.222$ ,  $m_1 = 0.187$  and  $c_1 = 0.659$ , giving  $\delta m_1 = -0.009$  and  $\delta c_1 = 0.046$ , indicating a mild peculiarity. Most recently, Mathys et al. (2023) measured a mean magnetic field modulus of this star to be  $\langle B \rangle \sim 3.8 \text{ kG}$ , whilst also concluding that the star must have a rotation period of the order 50 yr. They also suggested that the star could be host to a substellar companion with an orbital period suspected to be shorter than 100 d. They were also the first authors to report this star as an roAp star, with three pulsation modes, after analysing TESS data.

TESS observed TIC 120532285 during sector 16. The light curve showed no signature of rotation, as expected given the conclusions of Mathys et al. (2023), but does show high frequency variability as previously reported. We detect three modes in the amplitude spectrum (Fig. 20) at frequencies of  $2.171017 \pm 0.000011 \text{ mHz}$  ( $187.5759 \pm 0.0009 \text{ d}^{-1}$ ),  $2.198746 \pm 0.000008 \text{ mHz}$  ( $189.9716 \pm 0.0007 \text{ d}^{-1}$ ), and  $2.226270 \pm 0.000006 \text{ mHz}$  ( $192.3497 \pm 0.0006 \text{ d}^{-1}$ ). The separation between the modes (in increasing frequency) is  $27.73 \pm 0.01 \mu\text{Hz}$  and  $27.52 \pm 0.01 \mu\text{Hz}$  which are plausibly half of the large frequency separation, given the stellar parameters in the literature. We note a fourth, but not significant, peak in the amplitude spectrum that also follows the separation pattern, at the lower frequency side of the extracted three peaks.

#### 4.2.7 TIC 129820552

TIC 129820552 (TYC 2322-1440-1) is classified in the literature as an Ap SrEu star (Bidelman 1998). The Gaia collaboration, via DR2, provide the following parameters:  $T_{\text{eff}}$  in the range 7200 – 7400 K; a



**Figure 21.** Amplitude spectrum of TIC 129820552 with the pulsation mode identified with the vertical dotted line, with the dashed lines indicating the rotationally split sidelobes.

radius in the range of  $1.63 - 1.72 R_{\odot}$ , and a luminosity range of  $7.06 - 7.41 L_{\odot}$  (Gaia Collaboration et al. 2018). Previously, Netopil et al. (2017) determined the rotation period of the star to be 6.221 d. This star also appeared in the work of Balona (2022) who provided a rotation period of 6.211 d and identified a high frequency pulsation signal at a frequency of 2.013 mHz ( $173.928 \text{ d}^{-1}$ ) based on the TESS observations.


Using those same data, obtained in sector 18, we derived a rotation period of  $6.230 \pm 0.002 \text{ d}$  which differs from the previous determinations (although no errors are given on previous work). We were also able to identify the pulsation mode in this star at a frequency of  $2.01305 \pm 0.00002 \text{ mHz}$  ( $173.928 \pm 0.001 \text{ d}^{-1}$ ; Fig. 21) and note that there are four rotational sidelobes present in the amplitude spectrum. By selecting the appropriate zero-point in time, we were able to fit the quintuplet such that the phases of the peaks were all in agreement, indicating this is a non-distorted quadrupole mode, and found  $\tan i \tan \beta = 2.95 \pm 0.61$ , which implies  $i \approx 78^\circ$  and  $\beta \approx 32^\circ$ , or vice versa.

#### 4.2.8 TIC 394860395


TIC 394860395 (TYC 2666-85-1) has been classified as Ap SrEu type peculiar star (Bidelman 1985). Several studies have been carried out to reveal its magnetic structure. Kudryavtsev et al. (2006) calculated the longitudinal component of the magnetic field from spectral observations of the depression at 5200 Å. Their measurements indicated that the star has displayed a magnetic field with the observed  $B_{\ell}$  values ranging from  $-520$  to  $+540 \text{ G}$ . Bychkov et al. (2009) calculated the averaged quadratic effective magnetic field as  $\langle B_{\ell} \rangle = 380 \pm 111 \text{ G}$  by using the metallic spectral lines. Recently, Romanyuk et al. (2020) measured the longitudinal magnetic field and radial velocity values by using the data taken in 2012. Accordingly, they found  $B_{\ell}(z) = -340 \pm 40 \text{ G}$ ,  $B_{\ell}(r) = -380 \pm 20 \text{ G}$  and  $V_R = -32.5 \pm 3.5 \text{ km s}^{-1}$  at JD 2456140.484 and  $B_{\ell}(z) = 100 \pm 50 \text{ G}$ ,  $B_{\ell}(r) = 20 \pm 30 \text{ G}$ ,  $B_{\ell}(h) = -300 \pm 31$  and  $V_R = -31.4 \pm 3.0 \text{ km s}^{-1}$  at JD 2456233.261.

Paunzen et al. (2013) reported that the star has shown a prominent peak at about 5 mHz in the periodogram of Johnson B photometric data and noted that such a high frequency had never been detected in any roAp star. However, Paunzen et al. (2015) could not find any variability in the given frequency. They gave the effective temperature and interstellar reddening to be  $T_{\text{eff}} = 6800 \pm 400 \text{ K}$  and  $E(B-V)$  of 0.08 mag, respectively. Muñoz Bermejo et al. (2013) performed the principal component analysis to characterise the effective tem-

<sup>6</sup> These stars are spectroscopically similar to Am stars, but with an abnormally strong Sr II 4077 Å line (Bidelman 1981; North 1987)



**Figure 22.** Comparison of the PDC\_SAP light curve (grey) and the SAP light curve (black) for TIC 394860395. Both light curves show variability, but we argue neither is indicative of a rotation period. The dimming in the SAP data are flagged as bad points in the PDC\_SAP data, and are excluded from the final light curve.



**Figure 23.** Amplitude spectrum of TIC 394860395 showing the pulsation modes (marked by the vertical dotted lines) and the rotationally split sidelobes (marked with the vertical dashed lines).

perature from low-resolution spectroscopic data and calculated the temperature value of 7300 K.

This star was reported to be a rotationally variable roAp star by Balona (2022). He measured a rotation period of  $P_{\text{rot}} = 6.667$  d and a pulsation frequency of 1.180 mHz ( $101.949 \text{ d}^{-1}$ ), using *TESS* data. Here we use the *TESS* sector 14 data to refute the previously determined rotation period. There is no clear indication in the PDC\_SAP data that the period above is of an astrophysical nature. We inspected the SAP data and found significant variability, although that does not wholly appear astrophysical either. We show both light curves in Fig. 22.

The pulsation signature in this star is also somewhat complicated (Fig. 23). At first inspection, it appeared there were two modes, but after removing those there remained a third peak exactly between the two previous ones. Therefore, we have taken the pulsation frequency to be the central peak at  $1.18050 \pm 0.00005$  mHz ( $101.995 \pm 0.004 \text{ d}^{-1}$ ) and deduced that the two outer peaks are in fact sidelobes due to rotation. Those sidelobes indicate that the rotation period is on the order of  $23 \pm 2$  d. This is an example of where caution should be exercised when deducing rotation periods with weak signals in the low frequency range in Ap stars.

#### 4.2.9 TIC 435263600

TIC 435263600 (HD 218439) was first classified as an A2p: star with Sr, Cr, and Si peculiar features (Slettebak & Nassau 1959),

and more recently classified as an A5IVpSrSi star (Abt 1985). The star is a known visual binary (e.g., McAlister et al. 1987, 1990; Balega et al. 1994; Germain et al. 1999; Scardia et al. 2006), with a close companion ranging in distance from the primary between 0.47 arcsec to 0.62 arcsec (depending on the epoch), with a magnitude difference of 0.5 mag (Holden 1976). The mass of the primary is  $1.6 M_{\odot}$  (Kervella et al. 2019), with the orbital elements of:  $P_{\text{orb}} = 415 \pm 21$  yrs,  $e = 0.51 \pm 0.05$ , and the inclination of  $i = 165^{\circ} \pm 3^{\circ}$  (Ling 2010).

Strömgren-Crawford indices were measured by Hauck & Mermilliod (1998) to be:  $b - y = 0.187$ ,  $m_1 = 0.138$ ,  $c_1 = 1.008$  and  $H\beta = 2.841$ , giving  $\delta m_1 = 0.070$  and  $\delta c_1 = 0.154$ . These indices suggest the star is perhaps metal deficient, but we note that a binary companion will affect the values and they should not be used to draw conclusions. These indices also suggest a temperature of about 8200 K using relations of Moon & Dworetzky (1985), while an effective temperature of 8300 K was measured through the intensity of the Ca II K line (Floquet 1981).


Variability in this star was first reported by Balona (2022) who analysed the available *TESS* data. He classified it as an  $\alpha^2$  CVn star with both  $\gamma$  Dor and roAp pulsations. We corroborate his conclusion here, although note the complexities in the amplitude spectrum of the combined sector 17 and 24 *TESS* data. The top panel of Fig. 24 shows the low frequency variability in this star. We assumed the highest amplitude peak, which also has a harmonic, corresponds to the rotation period of the star ( $P_{\text{rot}} = 3.0739 \pm 0.0002$  d). We then assumed the remaining peaks, above 0.01 mHz are g-mode pulsations. However, given the star is a known binary it is unclear as to the source of the variability, or indeed if one or more of the peaks in the amplitude spectrum correspond to an orbital period. After removal of the dominant peaks, significant power remained in this low-frequency range. A longer timebase of observations is required to resolve these peaks.

At high frequency (Fig. 24 middle and lower panels) we identified clear signs of pulsations commensurate with those seen in roAp stars. In this frequency range there is significant frequency and amplitude modulation of the pulsations. We show in the bottom panel of Fig. 24 the amplitude spectra of the two sectors of data, which demonstrate this. It is possible to see the variability on shorter timescales too, on the order of 13-d *TESS* orbits or shorter. Such a phenomenon is becoming commonplace in the roAp stars given the long timebase, high precision datasets that are now available (e.g., Holdsworth 2021).

Finally, for this star we have retrieved a spectrum from the HERMES archive. There were two epochs of data, separated by 5 d, which showed very different signatures. Fig. B8 shows the spectra in the blue spectral region and  $H_{\alpha}$ . The change in the line profiles suggests there is a close companion with an orbital period much shorter than the period of 415 yr provided by Ling (2010), perhaps making this a triple system, thus we recommend this star be spectroscopically monitored to study the system in detail. Using the procedures outlined above, we derived the following parameters from the spectrum where primarily only one star is seen:  $T_{\text{eff}} = 7950 \pm 100$  K,  $\log g = 4.0 \pm 0.5$  ( $\text{cm s}^{-2}$ ) and  $v \sin i = 65 \pm 9$   $\text{km s}^{-1}$ . We note that these values should be treated cautiously given the binary nature of the star.

### 4.3 Known roAp stars prior to *TESS* launch

In this section, we present an analysis of the roAp stars which were known prior to the launch of the *TESS* mission. As with the new roAp star discoveries, we present only new information on stars which have already been discussed in the literature. There are 23 stars which are known to be roAp stars in the northern ecliptic hemisphere. Of these,



**Figure 24.** Amplitude spectra of TIC 435263600 showing (top) the low frequency range that indicated either rotational variation, g-mode pulsations, an orbital companion or a combination these phenomena. The middle panel shows the pulsation modes (marked by the vertical dotted lines). All modes show signs of frequency variability making precise determination of the frequencies difficult. The bottom panel shows the amplitude spectra for the same frequency range, but plotting the amplitude spectra for sector 17 (positive) and 24 (negative) data separately.

16 were observed by *TESS* in 2-min cadence in Cycle 2. Of the 7 that were not observed, 6 did not fall on a chip, while one star  $\beta$  CrB (TIC 383521659; HD 137909) was only observed at 30-min cadence in FFIs of sector 24.

#### 4.3.1 TIC 26418690

TIC 26418690 (KIC 11296437) was discovered to be an roAp star by Hey et al. (2019) after searching for high frequency pulsations in *Kepler* long cadence data using super-Nyquist asteroseismology (Murphy et al. 2013). The star was later studied by Murphy et al. (2020) who showed it to be the first  $\delta$  Sct-roAp hybrid star. We refer the reader to the latter study for a detailed analysis. The rotation period reported here, from the analysis of data from sectors 14 and 15, is in agreement with the more precise value determined from *Kepler* observations, and the pulsation mode is recovered at the expected frequency (Table 1). However, the rotational sidelobes are of too low

amplitude to be recovered in the *TESS* data. Fig. C1 shows the amplitude spectrum of this star, in the two regions of known pulsational variability. The spectral type of this star, A9 EuCr (Table 1), is from Hey et al. (2019).

The non-detection of the low amplitude signals in the *TESS* data is not surprising. Using a pulsating eclipsing binary system, Antoci et al. (2019) estimated that the redder passband of *TESS* accounts for an approximate 25 per cent reduction in the relative amplitude when compared to data from the *Kepler* mission. While temperature dependent, this, coupled with the higher noise level in *TESS* data compared to *Kepler* data, explains the non-detection of previously observed peaks.

#### 4.3.2 TIC 26749633

TIC 26749633 (KIC 11031749) is another roAp star discovered by Hey et al. (2019) in long cadence *Kepler* data. This star showed no rotation signal in the two *TESS* sectors in which it was observed, sectors 14 and 15, as was the case with the *Kepler* observations. There is a tentative detection of the pulsation mode at  $1.37275 \pm 0.00003$  mHz ( $118.605 \pm 0.003$  d $^{-1}$ ) with an amplitude of  $0.17 \pm 0.04$  mmag, as shown in Fig. C2. This tentative detection is likely a result of the reduced sensitivity of the *TESS* data over *Kepler* data, as discussed above. The spectral type of this star, F1 SrCrEu (Table 1), is from Hey et al. (2019).


#### 4.3.3 TIC 27395746

TIC 27395746 (KIC 11409673) is a further roAp star discovered by Hey et al. (2019). This is another target that does not show the rotation or pulsation signal as reported from the *Kepler* observations. However, with knowledge of the rotation period,  $P_{\text{rot}} = 1.06365 \pm 0.00002$  d, from Hey et al. (2019), we are able to conclude we detect a harmonic of the rotation frequency of the star, but without that prior knowledge, would have dismissed the signal as instrumental given both its low frequency and marginal significance, hence we have not listed it in Table 1. The combination of the angles  $i$  and  $\beta$  in this star result in a low amplitude mode with sidelobes of the pulsation having higher amplitude. Therefore, we are unable to detect the pulsation mode, but identified the  $\nu - \nu_{\text{rot}}$  sidelobe at a frequency of  $2.49995 \pm 0.00003$  mHz ( $215.996 \pm 0.002$  d $^{-1}$ ) with an amplitude of  $0.18 \pm 0.04$  mmag, as shown in Fig. C3. The spectral type of this star, A9 SrEu (Table 1), is from Hey et al. (2019).

What is noted in the sector 14 and 15 light curve of TIC 27395746 is the presence of eclipse signatures with a period of  $10.27 \pm 0.01$  d (Fig. 25) which are not present in the *Kepler* data. Inspection of the target pixel file (TPF) revealed a background eclipsing binary star included in the standard photometric aperture. While this star does appear in many catalogues concerned with eclipsing binary stars in the *Kepler* data (e.g., Conroy et al. 2014), although those sources quote the rotation period as the orbital period and have thus misclassified the star based on its surface spots.

#### 4.3.4 TIC 77128654

TIC 77128654 (HD 97127) was discovered in SuperWASP data to be an roAp star (Holdsworth et al. 2014a), and classified as an F3 SrEu(Cr). The amplitude spectrum of the sector 22 data for this star does not show any signature of rotation, but does reveal two significant pulsation frequencies (Fig. C4). The dominant mode frequency agrees with that in the literature ( $1.233855 \pm 0.000004$  mHz;



**Figure 25.** Phase folded light curve of TIC 27395746 showing an eclipse-like signature with a period of  $10.27 \pm 0.01$  d. The origin of the eclipse is unclear.

$106.6050 \pm 0.0003$  d $^{-1}$ ) while the second lower amplitude mode at  $1.248661 \pm 0.000031$  mHz ( $107.8843 \pm 0.0027$  d $^{-1}$ ) is newly reported here. The separation of these two modes is  $14.806 \pm 0.003$   $\mu$ Hz which is about one quarter of the expected large frequency separation for this star.

#### 4.3.5 TIC 123231021

TIC 123231021 (KIC 7582608) was discovered to be an roAp star through a search of the SuperWASP photometric archive (Holdsworth et al. 2014a), and later studied in detail by Holdsworth et al. (2014b). This star is known to show significant frequency/phase variability in its sole pulsation mode (Holdsworth et al. 2014b; Holdsworth 2021) which makes precise determination of the pulsation mode frequency difficult.

The *TESS* observations of this star, obtained in sectors 14 and 16, add no additional information over that already presented in the literature (Fig. C5). Each individual sector of data spans just over 1 rotation cycle of the star with the sectors separated by 300 d. This leads to a less precise, and inaccurate, determination of the rotation period in the *TESS* data of  $19.805 \pm 0.005$  d (compared to  $20.4401 \pm 0.0005$  d from Holdsworth et al. 2014b). The *TESS* data also do not resolve the rotationally split sidelobes observed in this star with *Kepler* data. Therefore, we refer the reader to Holdsworth et al. (2014b) for a detailed discussion on this star. The spectral type of this star, A7 EuCr (Table 1), is from Holdsworth et al. (2014a).

#### 4.3.6 TIC 158216369

TIC 158216369 (KIC 7018170) is another roAp star identified by Hey et al. (2019) in LC *Kepler* data. The sectors 14 and 26 data do not show signs of variability due to spots (as was the case with the *Kepler* data) nor pulsation (Fig. C6). Hey et al. (2019) suggested an intrinsic amplitude of 0.5 mmag for the dominant mode in this star which is above the 0.28 mmag upper limit we place on variability here. However, with the redder *TESS* bandpass, it is not surprising that the mode is not detected above the noise here. The spectral type of this star, F2 (SrCr)Eu (Table 1), is from Hey et al. (2019).

#### 4.3.7 TIC 158271090

TIC 158271090 (KIC 10195926) was discovered to be an roAp star in the *Kepler* field by Kurtz et al. (2011). *TESS* observed the star during sectors 14, 15 and 16. Analysis of the *TESS* light curve at low frequency allowed for the determination of the rotation period to be  $5.6845 \pm 0.0002$  d which is in agreement with that in the literature.

*Kepler* observations revealed two pulsation modes in this star, however the *TESS* observations only showed the highest frequency mode (which also has the highest amplitude) to be above the noise level (Fig. C7). The pulsation mode detected in this star is similar to that in TIC 27395746 where the geometry leads to higher amplitude sidelobes than the pulsation signal. Therefore, the pulsation frequency listed in Table 1 is derived from the average of the rotationally split sidelobes. The spectral type of this star, F0 Sr (Table 1), is from Gray et al. (2016).

#### 4.3.8 TIC 158275114

TIC 158275114 (KIC 8677585) was classified as an roAp star by Balona et al. (2011a) from 10 d of commissioning data from *Kepler*. It was further studied by Balona et al. (2013), with Holdsworth (2021) analysing all the *Kepler* data. The *Kepler* data showed two low frequency modes with disputed origins, and many modes in the typical range for roAp stars, all of which show frequency variability. The *TESS* data analysed here, from sectors 14 and 26, do not show the same low frequency variability as the *Kepler* data did, although the signal in the *Kepler* data is significantly below the noise of the *TESS* data at the same frequencies. There is a new ‘signal’ in the sector 14 data at a frequency of 0.012 mHz ( $1.00$  d $^{-1}$ ), however this is not present in the SPOC reduced 30-min cadence data for the same sector, nor the subsequent observations of this star, so we class this as an instrumental (or pipeline) artefact.

We detected two pulsation modes in the combined data for this star, significantly fewer than that seen in the *Kepler* data (Fig. C8). The two modes were detected at  $1.5853813 \pm 0.0000009$  mHz ( $136.97694 \pm 0.00008$  d $^{-1}$ ) and  $1.6217146 \pm 0.0000019$  mHz ( $140.11614 \pm 0.00016$  d $^{-1}$ ), where the separation of the two modes is likely to be the large frequency separation, as previously reported. In Table 1, the A5p spectral classification is from (Macrae 1952).


#### 4.3.9 TIC 159392323

TIC 159392323 (TYC 3547-2692-1) was classified as an roAp star by Holdsworth (2015) through a survey of SuperWASP photometry, who also classified the star as F3p SrEuCr. There have been no further studies of this star in the literature. The *TESS* sector 15 data confirm the previous pulsation in this star at a frequency of  $1.49021 \pm 0.00002$  mHz ( $128.754 \pm 0.001$  d $^{-1}$ ), and allowed for the detection of two new modes at  $1.47658 \pm 0.00005$  mHz ( $127.576 \pm 0.004$  d $^{-1}$ ) and  $1.49610 \pm 0.00003$  mHz ( $129.263 \pm 0.003$  d $^{-1}$ ). As with the previous literature, we did not detect a rotation period for this star.


#### 4.3.10 TIC 169078762

TIC 169078762 (HD 225914) was listed in the Henry Draper Catalogue and Extension (Cannon & Pickering 1923) as an A7 spectral type star, but found to be both chemically peculiar, A5Vp SrCr(Eu), and pulsating (in *Kepler* data) by Smalley et al. (2015). They performed a detailed analysis of this object and reported the pulsation mode at 0.7112 mHz ( $61.45$  d $^{-1}$ ) with an amplitude of 62.6  $\mu$ mag using the 4-yr long *Kepler* photometry, and measured a precise rotation period of  $P_{\text{rot}} = 5.209358 \pm 0.000009$  d.

The analysis of high-resolution spectra revealed the following stellar parameters:  $T_{\text{eff}} = 8100 \pm 200$  K,  $\log g = 4.0 \pm 0.2$  (cm s $^{-2}$ ),  $[\text{Fe}/\text{H}] = +0.31 \pm 0.24$  and  $v \sin i = 14.6 \pm 1.6$  km s $^{-1}$  (Smalley et al. 2015). Line profile variations caused by rotation are also evident. Lines of Sr, Cr, Eu, Mg and Si are strongest when the star is brightest



**Figure 26.** Amplitude spectrum for TIC 159392323. The vertical dotted lines identify the pulsation modes detected in the *TESS* data. The two lower amplitude modes are newly detected.



**Figure 27.** Amplitude spectrum for TIC 171988782. The vertical dotted lines identify the pulsation modes detected in the star.

(in the *Kepler* bandpass), while Y and Ba vary in antiphase with the other elements. The abundances of rare earth elements are only modestly enhanced compared to other roAp stars of similar  $T_{\text{eff}}$  and  $\log g$  (Smalley et al. 2015).

*TESS* observed the star in sector 14 at 2-min cadence. Both the previously determined rotation period and pulsation frequency (Fig. C9) are confirmed in these data.

#### 4.3.11 TIC 171988782

TIC 171988782 (HD 258048) was first classified as an F8 star (Kharchenko 2001) but Holdsworth et al. (2014a) showed it to be F4p EuCr(Sr) using low resolution spectroscopy. They determined an effective temperature of  $6200 \pm 280$  K using the IRFM, and 6600 K from Balmer line fitting. Holdsworth et al. (2014a) also classified the star to be an roAp star given the presence of a significant peak in the amplitude spectrum of SuperWASP photometry at 1.962 mHz ( $169.54 \text{ d}^{-1}$ ) with an amplitude of 1.49 mmag. Jayasinghe et al. (2018) identified TIC 171988782 as a W UMa type binary based on a period found in the ASAS-SN data. However, this period was not found in the SuperWASP data, nor in the present *TESS* data.

*TESS* observed the star in sector 20 of Cycle 2. The principal mode identified in the SuperWASP data is confirmed at  $1.962238 \pm 0.000005$  mHz ( $169.5374 \pm 0.0005 \text{ d}^{-1}$ ) with a new secondary mode found at  $1.975095 \pm 0.000023$  mHz ( $170.6482 \pm 0.0020 \text{ d}^{-1}$ ; Fig. 27). There is no indication of the rotation period in this star.

#### 4.3.12 TIC 264509538

TIC 264509538 (KIC 10685175) was discovered as a roAp star using super-Nyquist analysis on *Kepler* Long Cadence data (Hey et al. 2019). That work provided the first estimate for the principle pulsation frequency of 2.783 mHz ( $240.45189 \text{ d}^{-1}$ ) as well as the rotation period of  $P_{\text{rot}} = 3.1020 \pm 0.0001$  d. These values were reviewed by a new analysis based on *TESS* data by Shi et al. (2020), where the fundamental pulsation mode and rotational period were found to be 2.217 mHz ( $191.5151 \text{ d}^{-1}$ ) and  $3.1028 \pm 0.0048$  d, respectively. In that work the star was found to be an oblique quintuplet pulsator with sidelobes split by exactly the rotation frequency. The discrepancy between the two pulsation frequencies is a result of the incorrect selection of a Nyquist alias by Hey et al. (2019), an issue that is discussed in their work.

Hey et al. (2019) obtained the effective temperature ( $T_{\text{eff}} = 8000 \pm 300$  K) from LAMOST spectra, and a luminosity (scaled with the solar luminosity) of  $\log L/L_{\odot} = 0.896 \pm 0.022$  from *Gaia* DR2. A mass of  $M = 1.65 \pm 0.25 M_{\odot}$  was also estimated through an interpolation over stellar tracks based on the models of Cunha et al. (2013).

Since the *TESS* data, from sectors 14 and 15 (Fig. C10), have already been analysed, we do not repeat the process here, but refer the reader to Shi et al. (2020).

#### 4.3.13 TIC 272598185

TIC 272598185 (KIC 10483436) was discovered to be an roAp star by Balona et al. (2011b) through the analysis of *Kepler* SC data. In that work, it was concluded that the star pulsates with a single mode at a frequency of 1.35 mHz ( $116.91 \text{ d}^{-1}$ ), however the *TESS* data analysed here, from sectors 14 and 15, showed no signs of variability with this frequency (Fig. C11) which is most likely the result of the unfavourable longer wavelength observations of *TESS*. The rotation period presented in Table 1 is more precise than that presented by Balona et al. (2011b) since the combined *TESS* observations analysed here have a longer baseline than the 27-d *Kepler* dataset previously used.

#### 4.3.14 TIC 273777265

TIC 273777265 (KIC 6631188) was discovered to be an roAp star by Hey et al. (2019) through the analysis of *Kepler* LC data. The sector 14 and 15 *TESS* data showed low frequency variability which is attributed to rotation of the star. However, the derived rotation period is half of the value recorded in the literature, suggesting we detected only a harmonic of the rotation period of the star. The pulsation frequency found in the *TESS* data agreed with that in the literature within  $2.5 \sigma$  ( $1.49349 \pm 0.00001$  mHz;  $129.038 \pm 0.001 \text{ d}^{-1}$ ; Fig. C12). Interestingly for this object, the amplitude presented in Table 1 is larger than the predicted intrinsic amplitude in the *Kepler* bandpass which is unexpected due to the redder wavelength coverage of *TESS*. The spectral type of this star, F0 Sr (Table 1), is from Hey et al. (2019).

#### 4.3.15 TIC 286992225

TIC 286992225 (TYC 2553-480-1) was discovered by Holdsworth et al. (2014a) in SuperWASP data to be an roAp star. In that paper a single pulsation frequency of 2.726 mHz ( $235.54 \text{ d}^{-1}$ ) was found, which is one of the highest frequencies found for roAp stars. The authors classified the spectrum as A9p SrEu, and they found an effective temperature ranging from

7100 ± 200 K (from Balmer lines) to 7500 ± 400 K (from SED analysis). Moreover analysis of the H $\alpha$  profile seemed to indicate the presence of a core-wing anomaly (e.g., Cowley et al. 2001).

Analysis of *TESS* data for this star is presented in Mathys et al. (2022). The data two modes separated by 66  $\mu$ Hz, which the authors deduced that to be half of the large separation. TIC 286992225 is also classified as a super-slowly rotating Ap (ssrAp) star, i.e., an Ap star with a rotation period greater than 50 d.

We add no further analysis for this star with the *TESS* data (sector 23; Fig. C13), so refer the reader to Mathys et al. (2022) for full details.

#### 4.3.16 TIC 302602874

TIC 302602874 (TYC 2488-1241-1) was, upon its discovery as a variable star, classified as A6p SrEu (Holdsworth et al. 2014a). The authors found  $T_{\text{eff}}$  to be 7800 ± 200 K (from Balmer lines) and 8300 ± 500 K (from SED fitting). The latter estimate is in line with  $T_{\text{eff}} = 8200$  K provided in the TIC.

This star was the subject of its own study in Holdsworth et al. (2018c), in which the pulsation mode was concluded to be a distorted quadrupole mode. The *TESS* data analysed here (from sector 21; Fig C14) add no further information for this star; we refer the reader to Holdsworth et al. (2018c) for full details.

## 5 CANDIDATE ROAP STARS


In this section, we provide information on a number of stars that were flagged by the search teams as possible roAp stars. These stars are classed as candidate roAp stars since a combination of their spectral classification not indicating chemical peculiarity, their light curve does not show  $\alpha^2$  CVn rotational variability, their pulsation signature does not show signs of multiplets split by the rotation frequency, their presumed pulsation signal is close to a signal to noise limit of 4.0, or there is indication they are a member of a binary system. The stars listed in this section would benefit from high-resolution spectroscopy to search for peculiarities and a magnetic field.

#### 5.0.1 TIC 9171107

TIC 9171107 (TYC 2479-429-1) has limited information available in the literature. The TIC gives an effective temperature of 6800 K and  $\log g$  of 4.2 ( $\text{cm s}^{-2}$ ) while *Gaia* EDR3 gives  $T_{\text{eff}} = 6800$  K and  $\log g$  of 3.7 ( $\text{cm s}^{-2}$ ) (Gaia Collaboration 2020). No pulsation detections have been reported.

TIC 9171107 was observed in *TESS* sector 20 and showed no sign of rotation in its light curve. However, there are three significant short period signals in the amplitude spectrum of the light curve that we attribute to pulsation: 1.50971 ± 0.00004 mHz (130.439 ± 0.003 d $^{-1}$ ), 1.52902 ± 0.00002 mHz (132.107 ± 0.002 d $^{-1}$ ) and 1.53502 ± 0.00002 mHz (132.625 ± 0.002 d $^{-1}$ ; Fig. 28). The two lowest frequency modes are separated by 19.3  $\mu$ Hz, while the two highest frequency modes are separated by 6.0  $\mu$ Hz. The former separation may be half of the large frequency separation, while the latter separation may be the small frequency separation<sup>7</sup>.

There are no spectroscopic data of this object in the literature, and no Strömgen-Crawford indices that are available to classify the star



**Figure 28.** Amplitude spectrum of TIC 9171107, with the pulsation modes indicated with the vertical dotted lines.

as chemically peculiar. Therefore, we placed this star as a candidate roAp star until such data are available.

#### 5.0.2 TIC 275642037


TIC 275642037 (TYC 4330-716-1), has no published spectral type. However, from photometry Ofek (2008) and Pickles & Depagne (2010) both infer F0 V. The TIC gives  $T_{\text{eff}} = 7700 \pm 400$  K, while 7200 K was obtained by McDonald et al. (2017) and 7550 K by Ammons et al. (2006). The *Gaia* DR3 flags this star as photometrically variable as "DSCT|GDOR|SXPHE", with a *g*-band amplitude of 5.9 mmag at a single frequency of 8.1  $\mu$ Hz (0.6965 d $^{-1}$ , 1.436 d). *Gaia* DR3 data also showed a variable radial velocity of 4.56 ± 0.84 km s $^{-1}$ , with an amplitude of 10.67 km s $^{-1}$  and a spectral broadening of 44 ± 10 km s $^{-1}$ , which is a proxy for  $v \sin i$  (Frémat et al. 2023). Other than this, there are no other mentions of any chemical peculiarities, rotational variability or pulsations in the literature.

*TESS* observed this star during sector 18. There is clear low-frequency variability in this star that could either be attributed to rotation in the presence of stable spots, or an orbiting companion. The *Gaia* DR3 low-frequency result is half of the period we detected here ( $P_{\text{rot}} = 2.8745 \pm 0.0003$  d; Fig. 29), which is the dominant signal in an amplitude spectrum. The pulsations we detected (Fig. 30) at frequencies of 1.13943 ± 0.00003 mHz (98.447 ± 0.003 d $^{-1}$ ) and 1.16939 ± 0.00002 mHz (101.035 ± 0.001 d $^{-1}$ ) are stable in both amplitude and phase over the observation period. They are also separated by 30  $\mu$ Hz, which could plausibly be half the large separation. Given the short period rotation/orbital signature is unexpected in an roAp star it remains a candidate until the star is spectroscopically observed to determine any chemical peculiarity.


#### 5.0.3 TIC 298052991

TIC 298052991 (TYC 4554-625-1) has been classified as an A2 star by Kervella et al. (2019). The same authors also performed an analysis to characterise the presence of a stellar/substellar mass around the star and found a physical companion orbiting it. They gave the mass, assuming a semi-major axis of 1 au as  $M_2 = 12_5^6 M_{\text{Jup}}$ , using a primary mass and radius of  $M_1 = 1.60 \pm 0.08 M_{\odot}$  (Girardi et al. 2000) and  $R_1 = 1.50 \pm 0.08 R_{\odot}$  (Kervella et al. 2004). McDonald et al. (2012) derived the stellar temperature and luminosity to be  $T_{\text{eff}} = 9550$  K and  $L = 9.6 L_{\odot}$  from the comparison of the  $\chi^2$  statistic between observed photometric data and a set of synthetic stellar spectra. Balona (2022) presented an analysis of the *TESS* data from sectors


<sup>7</sup> defined as the difference between two odd or even degree modes that have a radial overtone difference of one




**Figure 29.** Phase folded light curve of TIC 275642037, phased on a period of  $2.8745 \pm 0.0003$  d. Binned 50:1



**Figure 31.** Phase folded SAP light curve of TIC 298052991, phased on a period of  $10.5852 \pm 0.0001$  d. Binned 200:1



**Figure 30.** Amplitude spectrum of TIC 275642037 showing the pulsation modes (marked by the vertical dotted lines).



**Figure 32.** Amplitude spectrum of TIC 298052991 showing the pulsation modes (marked by the vertical dotted lines).

14, 20, 21, 26, 40 and 41 and found a rotation period of 10.638 d and a pulsation frequency of 2.291 mHz ( $197.951 \text{ d}^{-1}$ ).


During Cycle 2, TIC 298052991 was observed during sectors 14, 20, 21 and 26. For this star, we used the SAP data to derive the rotation period as the signal in the PDC data had been altered by the processing. We derived a period of  $10.5852 \pm 0.0001$  d (Fig. 31), which significantly differs from that previously determined given the use of the SAP data here. To investigate the pulsations, we pre-whitened the individual sectors to remove peaks below 0.116 mHz ( $10 \text{ d}^{-1}$ ) to the noise level of the high-frequency range. This allowed us to detect three pulsation frequencies in this star (Fig. 32;  $2.217569 \pm 0.000003$  mHz,  $191.5979 \pm 0.0002 \text{ d}^{-1}$ ;  $2.246466 \pm 0.000002$  mHz,  $194.0947 \pm 0.0001 \text{ d}^{-1}$ ;  $2.291038 \pm 0.000001$  mHz,  $197.9457 \pm 0.0001 \text{ d}^{-1}$ ), with a tentative detection of a fourth ( $2.185497 \pm 0.000003$  mHz,  $188.8270 \pm 0.0003 \text{ d}^{-1}$ ).

Since there is no evidence of chemical peculiarities or rotational splitting of modes, we are unable to confirm if this star is a bona fide roAp star, thus it remains a candidate for now.

#### 5.0.4 TIC 405892692

TIC 405892692 (BD+49 3179) is a little-studied star. Hill & Schilt (1952) classified this  $V = 9.28$  mag star as F0. The TIC catalogue (Stassun et al. 2019) lists the atmospheric parameters  $T_{\text{eff}} = 7250$  K and  $\log g = 4.24$  ( $\text{cm s}^{-2}$ ), however there are no indications of chemical peculiarity for this star. Balona (2022) claimed the star to be a rotational variable with a pulsation at high frequency, based on TESS observations.


This star was observed in sectors 14 and 15. We detected a low frequency peak of uncertain origin in the amplitude spectrum, cor-



**Figure 33.** Amplitude spectrum of TIC 405892692 showing the detection of a pulsation mode (marked by the vertical dotted line).

responding to a period of  $P = 0.3786 \pm 0.0001$  d. This is a short rotation period for an F0 star, so we postulated that this is in fact a signature of a binary, or higher order, system. This is tentatively supported by the Gaia DR3 binary probability flag being essentially unity (Gaia Collaboration 2022). The pulsation signature we detected is at a frequency of  $1.17082 \pm 0.00002$  mHz ( $101.159 \pm 0.001 \text{ d}^{-1}$ ; Fig. 33), which agrees with that previously reported.

Since there is little information available on this star, and the lack of any convincing evidence that it is chemically peculiar, this remains a candidate roAp star.



**Figure 34.** Amplitude spectrum of TIC 429251527 showing the pulsation mode marked by the vertical dotted line.

### 5.0.5 TIC 429251527


TIC 429251527 (TYC 2185-478-1) is another star with very little information in the literature. This  $V = 10.5$  mag star has no spectral classification nor any other dedicated studies available. The atmospheric parameters,  $T_{\text{eff}} = 7900$  K and  $\log g = 4.66$  ( $\text{cm s}^{-2}$ ), are provided in the TIC (Stassun et al. 2019), while the *Gaia* DR3 data suggested a compatible value of  $T_{\text{eff}} = 7950$  K.

*TESS* observed the star in sector 15. The data did not show any rotation signature, but did show a pulsation with significant amplitude at  $1.77798 \pm 0.00001$  mHz ( $153.6178 \pm 0.0008$   $\text{d}^{-1}$ ; Fig. 34). This is the first reporting of the pulsation in this star. Given the presence of only one mode, which does not show rotational splitting, and the lack of information in the literature, we cannot confirm this star to be an roAp star, and thus leave this as a candidate for now.


### 5.0.6 TIC 445796153

TIC 445796153 (HD 34740) has several spectral classifications in the literature: Am (A3-A7) (Bertaud & Floquet 1967); A0p or A2V (Percy 1975); A0p (Cannon & Pickering 1918b). However, the peculiarity was marked as doubtful by Renson & Manfroid (2009). The estimated effective temperature from *Gaia* DR3 is  $T_{\text{eff}} = 10500$  K (Gaia Collaboration 2022), this, coupled with the spectral classifications, places this star at the hot end of the roAp range. Hauck & Mermilliod (1998) measured Strömgren-Crawford indices of this star to be:  $b - y = 0.073$ ,  $m_1 = 0.201$ ,  $c_1 = 0.937$  and  $H\beta = 2.817$ , giving  $\delta m_1 = 0.005$  and  $\delta c_1 = 0.127$ . However, there is disparity between the measured  $H\beta$  and  $b - y$  values, with the former giving an equivalent  $b - y$  of 0.129, redder than the observed value. The star is listed as a multiple system (Mason et al. 2001) with a  $\delta m_v = 2.96$  mag, which, based on the *Gaia*  $T_{\text{eff}}$  suggests a late B star with an early F companion. This may explain the discrepancies in the indices. The *TESS* data for TIC 445796153 were analysed by Balona (2022) to label the star as a possible  $\alpha^2$  CVn star (with a period of  $P_{\text{rot}} = 5.576$  d) with high-frequency  $\delta$  Sct modes.

Here, we used the *TESS* sector 19 and 26 data to confirm the rotation period of  $P_{\text{rot}} = 5.5721 \pm 0.0004$  d (Fig. 35). We also detected two pulsation modes (Fig. 36) that are in the frequency range where both  $\delta$  Sct and roAp pulsation modes can be found (e.g., TIC 356088697,  $0.646584 \pm 0.000008$  mHz; Paper I):  $0.601908 \pm 0.000002$  mHz ( $52.0049 \pm 0.0002$   $\text{d}^{-1}$ ) and  $0.675849 \pm 0.000002$  mHz ( $58.3934 \pm 0.0001$   $\text{d}^{-1}$ ). It is unusual for a  $\delta$  Sct star to exhibit so few modes, especially at this amplitude. However, we cannot rule out the possibility that these modes are indeed  $\delta$  Sct in nature, as that requires an investigation into the pulsation driving which differs be-



**Figure 35.** Phase folded light curve of TIC 445796153, phased on a period of  $5.5721 \pm 0.0004$  d. Binned 100:1



**Figure 36.** Amplitude spectrum of TIC 445796153. The two pulsation modes marked with vertical dotted lines.

tween the two classes and is beyond the scope of this work. We therefore class TIC 445796153 as a candidate roAp star, especially given the lack of convincing spectral classification of the star being chemically peculiar. Spectroscopic monitoring to provide a binary solution, coupled with time-resolved spectra will help identify the origin of the pulsations in this binary system. Efforts should also be made to exclude the possibility of a contaminating  $\delta$  Sct star.

### 5.0.7 TIC 91224991

TIC 91224991 (HD 191380) has a spectral classification in the literature of F8 (Cannon & Pickering 1923). Stassun et al. (2019) estimate the effective temperature and  $\log g$  to be 6700 K and  $3.97$  ( $\text{cm s}^{-2}$ ), respectively. The effective temperature is determined from the dereddened Bp-Rp colour, while the  $\log g$  value is derived using the estimated mass and radius. Another estimation of  $T_{\text{eff}}$  is given by Ammons et al. (2006) as 6650 K, while Anders et al. (2019) derived  $T_{\text{eff}} = 6960$  K based on narrow-band photometry.

Balona (2022) categorised this star as a rapidly oscillating A (roA) star, based on the *TESS* data. He estimated a luminosity  $\log L/L_{\odot} = 0.96$  using *Gaia* EDR3 parallaxes (Gaia Collaboration et al. 2016, 2021) in conjunction with interstellar reddening corrections from Gontcharov (2017) and a bolometric correction calibration by Pecaut & Mamajek (2013). Balona (2022) found a characteristic frequency  $1.160$  mHz ( $100.213$   $\text{d}^{-1}$ ) and also one harmonic, but does not give a rotation frequency. This star was also observed using long baseline interferometry by Cruzalèbes et al. (2019), who found an angular diameter corresponding to a radius of  $R = 2.13 R_{\odot}$ .

This star was added to the *TESS* 2-min target list as a result of a tentative detection of variability in KELT data (Pepper et al. 2007) that




Figure 37. Amplitude spectrum of TIC 91224991

was made available to the TASOC WG4 members. The *TESS* data, from sector 14, clearly showed the presence of high frequency variability. We determined the frequency to be  $1.159876 \pm 0.000002$  mHz ( $100.2133 \pm 0.0002$  d<sup>-1</sup>; Fig. 37), and note the presence of the second harmonic at twice this frequency. Our results are consistent with those of Balona (2022). There are no rotational sidelobes to the pulsation peak in the amplitude spectrum, which is expected given the lack of a rotation signal in the *TESS* data.

## 6 THE REMAINING ROAP STARS

To provide a complete sample of roAp stars, we must include those that have not been observed by *TESS* and those that have been observed or discovered beyond the Cycle 1 and Cycle 2 observations. Therefore, in Table 2 we present these stars. Where there are now *TESS* data available, we follow the procedures outlined above to provide the rotation and pulsation properties. For each star, we present their amplitude spectrum in the range of the pulsations in Appendix D, and for stars that have yet to be observed, we provide the literature information.

After accounting for the stars already presented as roAp stars in Paper I, this work, and that of Jayaraman et al. (2021) who reported TIC 198781841 and TIC 229960986 to be roAp stars, there are 27 additional stars listed in Balona (2022) as roAp stars, or candidate roAp stars. Of these 27, 11 appeared in Holdsworth et al. (2014a) as either Am or marginal Am (Am:) stars. Four of these 11 stars have no *TESS* 2-min cadence data so it is unclear how Balona (2022) decided a different variability type than the original classification, four further stars showed clear  $\delta$  Sct pulsations and one is a known binary with a compact pulsator. Of the two final Am(:) stars, TIC 22113439 is listed as a double lined spectroscopic binary in LAMOST DR9 data (Zheng et al. 2023) which explained the low frequency variability combined with the  $\delta$  Sct pulsations, and careful examination of the light curve of TIC 22132451 showed that the low-frequency variability changed phase over the observations (sectors 40, 52 and 53), which is not expected of an  $\alpha^2$  CVn variable. This, coupled with the pulsations being of a  $\delta$  Sct nature, means we rejected this as an roAp star.

Beyond the Am(:) stars, we considered TIC 427400331. This star is listed as a Be star by Labadie-Bartz et al. (2022) who commented on the presence of Slowly Pulsating B (SPB) star g-modes, and noted the unusual presence of high-frequency variability. We checked the high-frequency modes and found they did not show oblique pulsation, despite a confirmed rotation period. Two low resolution spectra, in the Be Star Spectra (BeSS) database did not show strong absorption lines typical of Ap stars, and the *Gaia* DR3 Re-normalised Unit

Weight Error (RUWE) value is  $>25$ , strongly indicating a binary star. We therefore rejected this star as an roAp star. We rejected a further eight stars on the grounds of either them being obvious  $\delta$  Sct stars or having no signal at the claimed frequency. Finally, two stars of the 27 roAp stars in Balona (2022) are listed as candidates in our work (TIC 410163387 in Paper I, and TIC 405892692 here), given there are no additional data products available, we did not include these stars as roAp in Table 2. The five remaining stars are included as roAp stars in Table 2.

In other work, Mathys et al. (2020) and Mathys et al. (2022) noted 10 candidate roAp stars between them that were detected as part of a systematic search of the *TESS* data for super slowly rotating Ap stars. Nine stars, one of which is the same as a candidate in Balona (2022), are rejected as roAp stars here due to the lack of a detection of an appropriate peak in the amplitude spectra. One star, TIC 72802368 from Mathys et al. (2020), remains a candidate as a signal at the reported frequency ( $1.207677 \pm 0.000001$  mHz) is detected in two (sectors 36 and 62) of the three available sectors.

Below, we provide comments on all stars in Table 2 with *TESS* data:

- *TIC 33604636* – There is clear signs of residual power in the amplitude spectrum after the removal of the pulsation frequency. It is not clear if there are further modes present, or amplitude/frequency variability at play (cf. Fig D1).
- *TIC 36576010* – There may be additional modes around 1.79 mHz that are not fully resolved (cf. Fig D2).
- *TIC 198781841* – The modes in this star are split by  $\sim 35.8$  mHz, which is plausibly half the large frequency separation (see Jayaraman et al. 2021) (cf. Fig D3).
- *TIC 229960986* – Application of the oblique pulsator model for this star resulted in a  $\tan i \tan \beta$  value of  $6.10 \pm 0.43$ , and the conclusion that the mode is a slightly distorted dipole mode due to unequal phases between the pulsation mode and the sidelobes (cf. Fig D4).
- *TIC 453826702* – The three modes identified in this star are separated by an average of  $\sim 33$   $\mu$ Hz which is plausibly half of the large frequency separation for this star. After removing the three identified modes, there remains excess power of unknown origin in the same frequency range (cf. Fig D5).
- *TIC 456673854* – The amplitude spectrum of this star showed, in increasing frequency, a dipole mode, a quadrupole mode and another dipole mode. The dipole modes form doublets with only the sidelobes visible while the quadrupole mode showed the pulsation and the  $\pm 2\nu_{\text{rot}}$  sidelobes. With the application of the oblique pulsator model, we found that all three modes have  $\tan i \tan \beta$  in agreement ( $7.57 \pm 3.38$  for the first dipole,  $10.91 \pm 3.84$  for the quadrupole, and  $5.29 \pm 2.57$  for the second dipole), with the phase relations suggesting the first two modes are non-distorted, while the second dipole mode is suggested to be a distorted mode. This star will benefit from an in-depth study (cf. Fig D6).
- *TIC 630844439* – The amplitude spectrum of this star showed the presence of both rotation and g-mode pulsations. It is noted in the literature as a binary star, so we postulate that the g-modes are from a  $\gamma$  Dor companion to an roAp star. The high-frequency peaks form a triplet, suggesting a single dipole mode in the roAp star. Applying the oblique pulsator model to the triplet resulted in  $\tan i \tan \beta = 2.48 \pm 0.27$ , and the indication that the mode is slightly distorted (cf. Fig D7).
- *TIC 96855460* – The SAP data were used for this star to provide a lower limit on the rotation period. No new information is gleaned about the pulsation mode (cf. Fig D8).

**Table 2.** Details of the roAp stars not observed in Cycles 1 or 2. The columns provide the TIC identifier and star name, the TIC v8.1 *TESS* magnitude, the spectral type, the effective temperature, as provided in the TIC, the sectors in which *TESS* observed the target, where applicable. The final three columns provide the stellar rotation period, the pulsation frequency(ies) (note that sidelobes that arise from oblique pulsation are not listed), and the pulsation amplitude(s) seen in the *TESS* data. A “\*” indicates excess power remains in the amplitude spectrum after extracting the modes listed. Frequencies marked with a † are derived from sidelobes. A ‡ denotes that the SAP data were used to derive the value. For the stars not observed by *TESS* the superscript after the TIC ID give the data reference, and we neglect providing the amplitude since various different techniques and/or filters were used, thus making them non-comparable to the *TESS* results.

TIC	HD/TYC name	<i>TESS</i> mag	Spectral type	$T_{\text{eff}}^{\text{TIC}}$ (K)	Sectors	$P_{\text{rot}}$ (d)	Pulsation frequency (mHz)	Pulsation amplitude (mmag)
<i>roAp stars discovered in TESS data</i>								
33604636	42605	8.78	Ap SrEuCr	8280	33	$2.7793 \pm 0.0011$	$1.25733 \pm 0.00002^*$	$0.068 \pm 0.007$
36576010	216018	7.45	A9Vp SrCrEu	7750	42	No signature	$1.92919 \pm 0.00003$	$0.045 \pm 0.006$
198781841	2134-154-1	10.58		7730	40,53,54	No signature	$1.8913785 \pm 0.0000005$	$0.25 \pm 0.01$
							$1.9271666 \pm 0.0000003$	$0.40 \pm 0.01$
							$1.9629854 \pm 0.0000009$	$0.15 \pm 0.01$
229960986	4221-240-1	10.38		8150	40,41,47-55	$2.093996 \pm 0.000003$	$2.2179825 \pm 0.0000009$	$0.082 \pm 0.006$
453826702	22032	8.81	Ap SrEuCr	8100	31	$4.8525 \pm 0.0007$	$1.57518 \pm 0.00003^*$	$0.051 \pm 0.007$
							$1.60998 \pm 0.00001$	$0.131 \pm 0.007$
							$1.64107 \pm 0.00003$	$0.061 \pm 0.007$
456673854	59702	8.56	A8V	7770	34	$5.438 \pm 0.001$	$3.13157 \pm 0.00004^\dagger$	<0.03
							$3.16928 \pm 0.00002$	$0.1132 \pm 0.0083$
							$3.20693 \pm 0.00004^\dagger$	<0.02
630844439	9348-1029-1	8.44			27,28	$1.6618 \pm 0.0003$	$3.36021 \pm 0.00001$	$0.13 \pm 0.01$
<i>Known roAp stars prior to TESS launch</i>								
96855460	185256	9.55	Fp SrEu	7100	27	~ 24	$1.62766 \pm 0.00003$	$0.13 \pm 0.01$
100196783	193756	8.98	Ap SrCrEu	7590	67	$3.0096 \pm 0.0005$	$1.28405 \pm 0.00001$	$0.199 \pm 0.008$
163587609	101065	7.53	F8/G0p	6125	37	No signature	$1.315020 \pm 0.000017$	$0.065 \pm 0.005$
							$1.369901 \pm 0.000010$	$0.116 \pm 0.005$
							$1.372881 \pm 0.000001^*$	$0.958 \pm 0.005$
							$1.393128 \pm 0.000010$	$0.112 \pm 0.005$
							$1.427743 \pm 0.000039$	$0.028 \pm 0.005$
293265536	4-562-1	9.93	A9p SrEu(Cr)	7640	42	No signature	$1.73910 \pm 0.00001$	$0.32 \pm 0.01$
299000970	176232	5.69	A7p ArEu	7520	53,54	No signature	$1.336785 \pm 0.000013$	$0.012 \pm 0.001$
							$1.396832 \pm 0.000003$	$0.045 \pm 0.001$
							$1.417638 \pm 0.000018$	$0.009 \pm 0.001$
							$1.427100 \pm 0.000005$	$0.029 \pm 0.001$
							$1.447892 \pm 0.000006$	$0.027 \pm 0.001$
							$1.508482 \pm 0.000024$	$0.006 \pm 0.001$
318007796	190290	9.61	Ap EuSr	7560	27,39	$4.09070 \pm 0.00002$	$2.2307867 \pm 0.0000015$	$0.049 \pm 0.007$
							$2.2723808 \pm 0.0000002$	$0.307 \pm 0.007$
354619745‡	201601	4.48	A9Vp SrCrEu	7490	55	No signature	$1.311230 \pm 0.000009$	$0.044 \pm 0.002$
							$1.338619 \pm 0.000001^*$	$0.317 \pm 0.002$
							$1.366022 \pm 0.000003$	$0.124 \pm 0.002$
383521659‡	137909	3.40	F2Vp SrEuSi		51	~ 18.3	No signal in <i>TESS</i> data	< 0.005
407929868‡	24355	9.43	A7/F0	7830	42,43,44	$27.724 \pm 0.008$	$2.5961335 \pm 0.0000003$	$2.056 \pm 0.006$
420687462	122970	7.93	F0	6800	50	No signature	$1.476848 \pm 0.000035$	$0.053 \pm 0.007$
							$1.477817 \pm 0.000015$	$0.120 \pm 0.007$
							$1.502519 \pm 0.000005$	$0.290 \pm 0.006$
445543326	12098	7.91		7760	58	$5.4816 \pm 0.0008$	$2.12206 \pm 0.00003$	$0.062 \pm 0.007$
							$2.12636 \pm 0.00004$	$0.040 \pm 0.007$
							$2.30602 \pm 0.00003$	$0.054 \pm 0.007$
<i>Known roAp stars not observed by TESS</i>								
153101639 <sup>a</sup>	–	11.87	A7Vp SrEu(Cr)	7510		$3.6747 \pm 0.0005$	$1.7585027 \pm 0.0000002$	
335303863 <sup>b</sup>	137949	6.45	F0Vp EuGdSr	7400		None published	$2.01475317 \pm 0.00000007$	
411247704 <sup>c</sup>	196470	9.49	Fp SrEu	7450		None published	1.544	
413938178 <sup>d</sup>	148593	8.71	Ap SrEuCr	7390		None published	$1.56 \pm 0.01$	
432223926 <sup>e</sup>	134214	7.17	F2Vp SrCrEu	7040		None published	$2.9496 \pm 0.0004$	
465996299 <sup>f</sup>	177765	8.87	Ap SrEuCr	7420		None published	$0.702580 \pm 0.000006$	
							$0.713382 \pm 0.000015$	
							$0.717106 \pm 0.000010$	

<sup>a</sup>Holdsworth et al. (2018b); <sup>b</sup>Holdsworth et al. (2018d); <sup>c</sup>Martinez et al. (1990); <sup>d</sup>Kochukhov et al. (2013); <sup>e</sup>Kreidl et al. (1994); <sup>f</sup>Holdsworth (2016)

- *TIC 100196783* – This star showed strong g-mode pulsation superimposed on the rotation signature. This is the first time this is mentioned in the literature. While there is no confirmation that this star is a binary, the *Gaia* RUWE suggests the star is a binary, however we are unsure how the pulsation may affect the calculation of this value. The *TESS* data provide no new information on the roAp pulsation (cf. Fig D9).

- *TIC 163587609* – While there are five pulsation modes detected in the star, there is no clear indication of the large frequency separation. Three modes are separated by  $\sim 55 \mu\text{Hz}$ , but this value is in disagreement with  $64 \mu\text{Hz}$  from [Mkrtychian et al. \(2008\)](#). It is unclear which, if either, of these large separation values is correct. There is also excess power remaining in the amplitude spectrum around the principal peak after it was removed (cf. Fig D10).

- *TIC 293265536* – No new information is provided by the *TESS* data (cf. Fig D11).

- *TIC 299000970* – With the identification of six pulsation modes, we were able to identify two recurrent separations of  $\sim 60 \mu\text{Hz}$  and  $\sim 30 \mu\text{Hz}$ , which we interpreted as the large frequency separation, and half of it. This is different from the value adopted by [Huber et al. \(2008\)](#) of  $50 \mu\text{Hz}$ , based on MOST photometry (cf. Fig D12).

- *TIC 318007796* – The *TESS* data showed both modes in this star to be multiplets thus allowing for the determination that the lowest frequency mode is a dipole mode, with the other being a quadrupole mode, which suggests a large frequency separation of  $\sim 83 \mu\text{Hz}$ . Applying the oblique pulsator model resulted in  $\tan i \tan \beta$  values for the dipole and quadrupole of  $4.01 \pm 0.61$  and  $3.51 \pm 0.27$ , respectively, with the phase variations relations for each mode suggesting they are both distorted (cf. Fig D13).

- *TIC 354619745* – We list three pulsation modes in this star, but note there is residual power remaining in the amplitude spectrum after they are removed. The three modes are each separated by  $27.4 \mu\text{Hz}$ , which is presumably half of the large frequency separation. The principal mode showed a harmonic at twice its frequency, which is joined by a second peak whose frequency is the sum of the two highest amplitude modes. This non-linear interaction is only the third occurrence noted in the literature (after 33 Lib ([Holdsworth et al. 2018d](#)) and  $\alpha$  Cir [Paper I](#); cf. Fig D14).

- *TIC 383521659* – We did not detect the pulsation mode in this star in the *TESS* data. The pulsation was detected spectroscopically with an amplitude of about  $30 \text{ m s}^{-1}$  ([Kurtz et al. 2007](#)), and is suspected to be below the *TESS* noise level (cf. Fig D15).

- *TIC 407929868* – The *TESS* data of this star showed a highly distorted quadrupole mode, but did not add additional information over the literature (cf. Fig D16).

- *TIC 420687462* – The *TESS* data do not add new information to the literature for this star (cf. Fig D17).

- *TIC 445543326* – This star showed a dipole triplet with a close in frequency second mode, and a third mode at higher frequency. Applying the oblique pulsator model to the dipole triplet resulted in a  $\tan i \tan \beta$  value of  $2.28 \pm 0.30$ , with the phase relations showing this mode to be very distorted. A more detailed analysis of this star is required (cf. Fig D18).

## 7 CONCLUSIONS

This work constitutes the second phase of analysing the *TESS* data in the search for high-frequency pulsations in Ap stars. We have used the 2-min cadence Cycle 2 data of all hot ( $T_{\text{eff}} > 6000 \text{ K}$ ) stars in the Northern Ecliptic hemisphere, amounting to over 36 000 stars. This has enabled us to: identify seven new roAp stars unreported as such

in the literature, present a detailed analysis of a further 9 roAp stars previously identified as new in the *TESS* data, show that four of 16 well established roAp stars do not show variability in the analysed *TESS* data, and present 7 candidate roAp stars where insufficient spectral data exist to admit the star to the roAp class.

Three of the seven new roAp stars are multiperiodic pulsators, thus offering the opportunity for a more complete asteroseismic analysis. Furthermore, of these seven stars, five were previously not identified as chemically peculiar stars. This has important ramifications for the continued search for roAp stars in *TESS* data, and indeed other data sets. We have shown that catalogues of CP stars are, understandably, incomplete and as such, the search for high-frequency pulsations should include stars previously not identified as CP.

All of these new roAp stars are rotationally variable stars, and of the 32 roAp stars presented here, almost 70 per cent (22 stars) show variability due to spots. When we combine the results from this paper, [Paper I](#), and all roAp stars in the literature, we found 63 per cent (71 of 112 roAp stars) showed rotation signatures in their light curves. While some periods are still to be determined, many of the stars showed periods less than 10 d. A future investigation will explore the potential correlations between rotation rates and pulsation properties (Cunha et al., in prep.), with the ssrAp stars showing a higher incidence of roAp stars than is found for all Ap stars ([Mathys et al. 2020, 2022](#)).

We have identified a few examples of short timescale mode variability, which while not as extreme as in some cases, show that the modes in the roAp stars can be unstable on the order of days to weeks. A significant example here is *TIC 26833276* where we see changes in mode amplitude between *TESS* orbits (cf., Fig. 16). More significant is *TIC 435263600*, which is complex in both its pulsation spectrum (Fig 24) and its stellar spectrum (Fig. B8). With a presumed close companion, this star provides a prime opportunity to investigate whether interactions between roAp stars and companions affect their pulsation spectra.

As with [Paper I](#), we note the lack of detection of roAp pulsations in the hottest of the Ap stars. The theoretical blue edge of the roAp instability strip reaches to temperatures of about 10 000 K whereas the hottest known pulsator based on the TIC temperature lies at  $T_{\text{eff,TIC}} \sim 8540 \text{ K}$  (*TIC 119327278*; [Holdsworth et al. 2021](#)). This presents a significant challenge for the driving of pulsation in the hottest Ap stars, showing the need to further develop theoretical models of these stars.

Our results show that the incidence of roAp stars among the Ap stars is low. Before the launch of *TESS*, the TASOC WG4 proposed to observe all stars listed as Ap in the [Renson & Manfroid \(2009\)](#) catalogue with 2-min cadence during the nominal mission. This amounted to 1381 stars proposed, of which 1285 have been observed to date. Of the 112 stars presented across our two papers, 73 appear in that catalogue. This implies that the incidence of roAp stars in the known Ap star population is just 5.5 per cent. This raises the fundamental question of: why do only some Ap stars pulsate? We must consider that, as we have shown, some pulsation amplitudes are below the *TESS* detection limit, but that is only a fraction of the known roAp stars. Thanks to our work, we now have a statistically significant sample of roAp stars with which we can investigate this conundrum, and drive forward our theoretical understanding of these complex stars.

These works, [Paper I](#) and the current paper, represent the most comprehensive and collaborative observational study of the roAp stars observed during the nominal mission of *TESS*. In this collective work, we have identified 19 new roAp stars, and presented an analysis of 87 more. Our approach has resulted in a homogeneous sample of

roAp stars in both the detection methods used and their analysis techniques and software. This resource will stand as the definitive collection of roAp stars observed during Cycles 1 & 2 of the *TESS* mission, with other roAp stars noted in the literature from later *TESS* Cycles 4, 5 and 6.

## ACKNOWLEDGEMENTS

DLH acknowledges financial support from the Science and Technology Facilities Council (STFC) via grant ST/M000877/1. MSC acknowledges the support of Fundação para a Ciência e Tecnologia FCT/MCTES, Portugal, through national funds by these grants UIDB/04434/2020, UIDP/04434/2020.FCT, 2022.03993.PTDC and CEECIND/02619/2017. VA was supported by a research grant (00028173) from VILLUM FONDEN.

DMB gratefully acknowledges a senior postdoctoral fellowship from Research Foundation Flanders (FWO; grant number: 1286521N), and the Engineering and Physical Sciences Research Council (EPSRC) of UK Research and Innovation (UKRI) in the form of a Frontier Research grant under the UK government's Horizon Europe funding guarantee (grant number: [EP/Y031059/1]).

A.H. acknowledges support from the Science and Technology Facilities Council (STFC) grant No. ST/X000915/1.

ÅS and ZsB acknowledge the financial support of the KKP-137523 'SeismoLab' Èlvonal grant of the Hungarian Research, Development and Innovation Office (NKFIH).

This research was supported by the KKP-137523 'SeismoLab' Èlvonal grant of the Hungarian Research, Development and Innovation Office (NKFIH) and by the MW-Gaia COST Action (CA18104).

ZsB acknowledges the support by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences. IS acknowledges a partial financial support from BFNI via grant KII-06-H58/30. JCS acknowledges support from the Spanish State Research Agency (AEI) Projects No. PID2019-107061GB-C64: "Contribution of the UGR to the PLATO2.0 space mission. Phases C/D-1". MS acknowledges the support by Inter-transfer grant no LTT-20015. LFM acknowledges the financial support from the UNAM under grant PAPIIT IN117323. E. Ziaali acknowledges Iran National Science Foundation (INSF) for its total support under grant No. 4002562. SJM was supported by the Australian Research Council (ARC) through Future Fellowship FT210100485.

This paper includes data collected by the *TESS* mission, which are publicly available from the Mikulski Archive for Space Telescopes (MAST). Funding for the *TESS* mission is provided by NASA's Science Mission directorate. Funding for the *TESS* Asteroseismic Science Operations Centre is provided by the Danish National Research Foundation (Grant agreement no.: DNR106), ESA PRODEX (PEA 4000119301) and Stellar Astrophysics Centre (SAC) at Aarhus University. We thank the *TESS* team and staff and TASC/TASOC for their support of the present work. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France.

Guoshoujing Telescope (the Large Sky Area Multi-Object Fiber Spectroscopic Telescope LAMOST) is a National Major Scientific Project built by the Chinese Academy of Sciences. Funding for the project has been provided by the National Development and Reform Commission. LAMOST is operated and managed by the National Astronomical Observatories, Chinese Academy of Sciences.

Some of the data in this work are observations obtained with the HERMES spectrograph, which is supported by the Research Foundation - Flanders (FWO), Belgium, the Research Council of KU Leuven, Belgium, the Fonds National de la Recherche Scientifique

(F.R.S.-FNRS), Belgium, the Royal Observatory of Belgium, the Observatoire de Genève, Switzerland, and the Thüringer Landessternwarte Tautenburg, Germany, and were assembled by the Mercator Telescope, operated on the island of La Palma by the Flemish Community, at the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias.

This work has made use of the BeSS database, operated at LESIA, Observatoire de Meudon, France: <http://basebe.obspm.fr>.

## REFERENCES

- Abt H. A., 1985, *ApJS*, **59**, 95
- Aerts C., Christensen-Dalsgaard J., Kurtz D. W., 2010, *Asteroseismology*. Springer Science and Business Media, doi:10.1007/978-1-4020-5803-5
- Alecian G., Stift M. J., 2010, *A&A*, **516**, A53
- Alentiev D., Kochukhov O., Ryabchikova T., Cunha M., Tsymbal V., Weiss W., 2012, *MNRAS*, **421**, L82
- Ammons S. M., Robinson S. E., Strader J., Laughlin G., Fischer D., Wolf A., 2006, *ApJ*, **638**, 1004
- Anders F., et al., 2019, *A&A*, **628**, A94
- Andrae R., et al., 2018, *A&A*, **616**, A8
- Antoci V., et al., 2019, *MNRAS*, **490**, 4040
- Balega I. I., Balega Y. Y., Belkin I. N., Maximov A. F., Orlov V. G., Pluzhnik E. A., Shkhagosheva Z. U., Vasyuk V. A., 1994, *A&AS*, **105**, 503
- Balmforth N. J., Cunha M. S., Dolez N., Gough D. O., Vauclair S., 2001, *MNRAS*, **323**, 362
- Balona L. A., 2022, *MNRAS*, **510**, 5743
- Balona L. A., et al., 2011a, *MNRAS*, **410**, 517
- Balona L. A., et al., 2011b, *MNRAS*, **413**, 2651
- Balona L. A., et al., 2013, *MNRAS*, **432**, 2808
- Balona L. A., Holdsworth D. L., Cunha M. S., 2019, *MNRAS*, **487**, 2117
- Barceló Forteza S., Michel E., Roca Cortés T., García R. A., 2015, *A&A*, **579**, A133
- Barceló Forteza S., Moya A., Barrado D., Solano E., Martín-Ruiz S., Suárez J. C., García Hernández A., 2020, *A&A*, **638**, A59
- Bernhard K., Hümmerich S., Paunzen E., 2020, *MNRAS*, **493**, 3293
- Bertaud C., Floquet M., 1967, *Journal des Observateurs*, **50**, 425
- Bidelman W. P., 1966, *Vistas in Astronomy*, **8**, 53
- Bidelman W. P., 1981, *AJ*, **86**, 553
- Bidelman W. P., 1985, *AJ*, **90**, 341
- Bidelman W. P., 1998, *PASP*, **110**, 270
- Bigot L., Dziembowski W. A., 2002, *A&A*, **391**, 235
- Blanco-Cuaresma S., 2019, *MNRAS*, **486**, 2075
- Blanco-Cuaresma S., Soubiran C., Heiter U., Jofré P., 2014, *A&A*, **569**, A111
- Bowman D. M., Kurtz D. W., 2018, *MNRAS*, **476**, 3169
- Brandt T. D., 2021, *ApJS*, **254**, 42
- Bychkov V. D., Bychkova L. V., Madej J., 2009, *MNRAS*, **394**, 1338
- Bychkov V. D., Bychkova L. V., Madej J., 2016, *MNRAS*, **455**, 2567
- Cannon A. J., Pickering E. C., 1918a, *Annals of Harvard College Observatory*, **91**, 1
- Cannon A. J., Pickering E. C., 1918b, *Annals of Harvard College Observatory*, **92**, 1
- Cannon A. J., Pickering E. C., 1919, *Annals of Harvard College Observatory*, **94**, 1
- Cannon A. J., Pickering E. C., 1922, *Annals of Harvard College Observatory*, **97**, 1
- Cannon A. J., Pickering E. C., 1923, *Annals of Harvard College Observatory*, **98**, 1
- Cannon A. J., Pickering E. C., 1924, *Annals of Harvard College Observatory*, **99**, 1
- Castelli F., Kurucz R. L., 2003, in Piskunov N., Weiss W. W., Gray D. F., eds, Vol. 210, *Modelling of Stellar Atmospheres*. p. A20 (arXiv:astro-ph/0405087), doi:10.48550/arXiv.astro-ph/0405087
- Clausen J. V., Jensen K. S., 1979, in IAU Colloq. 47: *Spectral Classification of the Future*. p. 479

- Conroy K. E., Prša A., Stassun K. G., Orosz J. A., Fabrycky D. C., Welsh W. F., 2014, *AJ*, **147**, 45
- Cowley A. P., Cowley C. R., 1964, *PASP*, **76**, 119
- Cowley A. P., Cowley C. R., 1965, *PASP*, **77**, 184
- Cowley C. R., Hubrig S., Ryabchikova T. A., Mathys G., Piskunov N., Mittemayer P., 2001, *A&A*, **367**, 939
- Crawford D. L., 1975, *AJ*, **80**, 955
- Crawford D. L., 1979, *AJ*, **84**, 1858
- Cruzalèbes P., et al., 2019, *MNRAS*, **490**, 3158
- Cui X.-Q., et al., 2012, *Research in Astronomy and Astrophysics*, **12**, 1197
- Cunha M. S., 2006, *MNRAS*, **365**, 153
- Cunha M. S., Gough D., 2000, *MNRAS*, **319**, 1020
- Cunha M. S., Alentiev D., Brandão I. M., Perrauc K., 2013, *MNRAS*, **436**, 1639
- Cunha M. S., et al., 2019, *MNRAS*, **487**, 3523
- Deal M., Cunha M. S., Keszthelyi Z., Perrauc K., Holdsworth D. L., 2021, *A&A*, **650**, A125
- Floquet M., 1981, *A&A*, **101**, 176
- Frémat Y., et al., 2023, *A&A*, **674**, A8
- Freyhammer L. M., Kurtz D. W., Cunha M. S., Mathys G., Elkin V. G., Riley J. D., 2008, *MNRAS*, **385**, 1402
- Gaia Collaboration 2018, *VizieR Online Data Catalog*, p. I/345
- Gaia Collaboration 2020, *VizieR Online Data Catalog*, p. I/350
- Gaia Collaboration 2022, *VizieR Online Data Catalog*, p. I/355
- Gaia Collaboration et al., 2016, *A&A*, **595**, A1
- Gaia Collaboration et al., 2018, *A&A*, **616**, A1
- Gaia Collaboration et al., 2021, *A&A*, **649**, A1
- Germain M. E., Douglass G. G., Worley C. E., 1999, *AJ*, **117**, 1905
- Girardi L., Bressan A., Bertelli G., Chiosi C., 2000, *A&AS*, **141**, 371
- Gomez A. E., Luri X., Grenier S., Figueras F., North P., Royer F., Torra J., Mennessier M. O., 1998, *A&A*, **336**, 953
- Gontcharov G. A., 2006, *Astronomy Letters*, **32**, 759
- Gontcharov G. A., 2017, *Astronomy Letters*, **43**, 472
- Gray R. O., et al., 2016, *AJ*, **151**, 13
- Grenier S., et al., 1999, *A&AS*, **137**, 451
- Gruberbauer M., et al., 2011, *A&A*, **530**, A135
- Hauck B., Mermilliod M., 1998, *A&AS*, **129**, 431
- Hey D. R., et al., 2019, *MNRAS*, **488**, 18
- Hill S. J., Schilt J., 1952, Contributions from the Rutherford Observatory of Columbia University New York, **32**, 1
- Hoag A. A., Applequist N. L., 1965, *ApJS*, **12**, 215
- Holden F., 1976, *PASP*, **88**, 325
- Holdsworth D. L., 2015, PhD thesis, Keele University
- Holdsworth D. L., 2016, *Information Bulletin on Variable Stars*, **6185**, 1
- Holdsworth D. L., 2021, *Frontiers in Astronomy and Space Sciences*, **8**, 31
- Holdsworth D. L., et al., 2014a, *MNRAS*, **439**, 2078
- Holdsworth D. L., Smalley B., Kurtz D. W., Southworth J., Cunha M. S., Clubb K. I., 2014b, *MNRAS*, **443**, 2049
- Holdsworth D. L., Kurtz D. W., Smalley B., Saio H., Handler G., Murphy S. J., Lehmann H., 2016, *MNRAS*, **462**, 876
- Holdsworth D. L., et al., 2018a, *MNRAS*, **473**, 91
- Holdsworth D. L., Saio H., Bowman D. M., Kurtz D. W., Sefako R. R., Joyce M., Lambert T., Smalley B., 2018b, *MNRAS*, **476**, 601
- Holdsworth D. L., Saio H., Sefako R. R., Bowman D. M., 2018c, *MNRAS*, **480**, 2405
- Holdsworth D. L., Cunha M. S., Shibahashi H., Kurtz D. W., Bowman D. M., 2018d, *MNRAS*, **480**, 2976
- Holdsworth D. L., et al., 2021, *MNRAS*, **506**, 1073
- Huber D., et al., 2008, *A&A*, **483**, 239
- Hubrig S., Järvinen S. P., Alvarado-Gómez J. D., Ilyin I., Schöller M., 2023, *MNRAS*, **526**, L83
- Jayaraman R., Kurtz D. W., Handler G., Rappaport S., Ricker G., 2021, *Research Notes of the American Astronomical Society*, **5**, 268
- Jayasinghe T., et al., 2018, *MNRAS*, **477**, 3145
- Jenkins J. M., et al., 2016, in Chiozzi G., Guzman J. C., eds, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series Vol. 9913, Software and Cyberinfrastructure for Astronomy IV. p. 99133E, doi:10.1117/12.2233418
- Joshi S., Mary D. L., Martinez P., Kurtz D. W., Girish V., Seetha S., Sagar R., Ashoka B. N., 2006, *A&A*, **455**, 303
- Joshi S., et al., 2016, *A&A*, **590**, A116
- Kervella P., Thévenin F., Di Folco E., Ségransan D., 2004, *A&A*, **426**, 297
- Kervella P., Arenou F., Mignard F., Thévenin F., 2019, *A&A*, **623**, A72
- Khalack V., Kobzar O., Lenz P., 2021, in MOBSTER-1 virtual conference: Stellar Variability as a Probe of Magnetic Fields in Massive Stars. p. 7, doi:10.5281/zenodo.5504415
- Kharchenko N. V., 2001, *Kinematika i Fizika Nebesnykh Tel*, **17**, 409
- Kochukhov O., Alentiev D., Ryabchikova T., Boyko S., Cunha M., Tsymbal V., Weiss W., 2013, *MNRAS*, **431**, 2808
- Kreidl T. J., Kurtz D. W., Schneider H., van Wyk F., Roberts G., Marang F., Birch P. V., 1994, *MNRAS*, **270**, 115
- Kudryavtsev D. O., Romanyuk I. I., Elkin V. G., Paunzen E., 2006, *MNRAS*, **372**, 1804
- Kupka F., Dubernet M. L., VAMDC Collaboration 2011, *Baltic Astronomy*, **20**, 503
- Kurtz D. W., 1982, *MNRAS*, **200**, 807
- Kurtz D. W., 1985, *MNRAS*, **213**, 773
- Kurtz D. W., Shibahashi H., Goode P. R., 1990, *MNRAS*, **247**, 558
- Kurtz D. W., Elkin V. G., Mathys G., 2007, *MNRAS*, **380**, 741
- Kurtz D. W., et al., 2011, *MNRAS*, **414**, 2550
- Kurucz R. L., 1993, SYNTHE spectrum synthesis programs and line data
- Kurucz R. L., 2005, *Memorie della Societa Astronomica Italiana Supplementi*, **8**, 14
- Labadie-Bartz J., Carciofi A. C., Henrique de Amorim T., Rubio A., Luiz Figueiredo A., Ticiani dos Santos P., Thomson-Paressant K., 2022, *AJ*, **163**, 226
- LeBlanc F., Monin D., Hui-Bon-Hoa A., Hauschildt P. H., 2009, *A&A*, **495**, 937
- Ling J. F., 2010, *AJ*, **139**, 1521
- Liu C., et al., 2020, arXiv e-prints, p. arXiv:2005.07210
- Lüftinger T., Kochukhov O., Ryabchikova T., Piskunov N., Weiss W. W., Ilyin I., 2010, *A&A*, **509**, A71
- Luo A. L., et al., 2015, *Research in Astronomy and Astrophysics*, **15**, 1095
- Macrae D. A., 1952, *ApJ*, **116**, 592
- Martinez P., Kurtz D. W., Kauffmann G., Jonson A. C., 1990, *Information Bulletin on Variable Stars*, **3506**, 1
- Martinez P., Kurtz D. W., Kauffmann G. M., 1991, *MNRAS*, **250**, 666
- Masana E., Jordi C., Maitzen H. M., Torra J., 1998, *A&AS*, **128**, 265
- Masana E., Jordi C., Ribas I., 2006, *A&A*, **450**, 735
- Mason B. D., Wycoff G. L., Hartkopf W. I., Douglass G. G., Worley C. E., 2001, *AJ*, **122**, 3466
- Mathys G., 2017, *A&A*, **601**, A14
- Mathys G., Kurtz D. W., Holdsworth D. L., 2020, *A&A*, **639**, A31
- Mathys G., Kurtz D. W., Holdsworth D. L., 2022, *A&A*, **660**, A70
- Mathys G., Khalack V., Kobzar O., LeBlanc F., North P. L., 2023, *A&A*, **670**, A72
- McAlister H. A., Hartkopf W. I., Hutter D. J., Franz O. G., 1987, *AJ*, **93**, 688
- McAlister H., Hartkopf W. I., Franz O. G., 1990, *AJ*, **99**, 965
- McDonald I., Zijlstra A. A., Boyer M. L., 2012, *MNRAS*, **427**, 343
- McDonald I., Zijlstra A. A., Watson R. A., 2017, *MNRAS*, **471**, 770
- Michaud G., Megessier C., Charland Y., 1981, *A&A*, **103**, 244
- Mkrtichian D. E., Hatzes A. P., Saio H., Shobbrook R. R., 2008, *A&A*, **490**, 1109
- Moon T. T., Dworetzky M. M., 1985, *MNRAS*, **217**, 305
- Muñoz Bermejo J., Asensio Ramos A., Allende Prieto C., 2013, *A&A*, **553**, A95
- Murphy S. J., Shibahashi H., Kurtz D. W., 2013, *MNRAS*, **430**, 2986
- Murphy S. J., Hey D., Van Reeth T., Bedding T. R., 2019, *MNRAS*, **485**, 2380
- Murphy S. J., Saio H., Takada-Hidai M., Kurtz D. W., Shibahashi H., Takata M., Hey D. R., 2020, *MNRAS*, **498**, 4272
- Nelson M. J., Kreidl T. J., 1993, *AJ*, **105**, 1903
- Netopil M., Paunzen E., Hümmelich S., Bernhard K., 2017, *MNRAS*, **468**, 2745
- Niedzielski A., Muciek M., 1988, *Acta Astron.*, **38**, 225
- North P., 1987, *A&A*, **186**, 191
- North P., Duquennoy A., 1991, *A&A*, **244**, 335

- Ofek E. O., 2008, *PASP*, **120**, 1128  
 Olsen E. H., 1980, *A&AS*, **39**, 205  
 Paunzen E., 2015, *A&A*, **580**, A23  
 Paunzen E., Netopil M., Rode-Paunzen M., Handler G., Božić H., Ružđjak D., Sudar D., 2012, *A&A*, **542**, A89  
 Paunzen E., et al., 2013, *Information Bulletin on Variable Stars*, **6058**, 1  
 Paunzen E., Netopil M., Rode-Paunzen M., Handler G., Božić H., 2015, *A&A*, **575**, A24  
 Paunzen E., et al., 2018, *Research in Astronomy and Astrophysics*, **18**, 135  
 Pecaut M. J., Mamajek E. E., 2013, *ApJS*, **208**, 9  
 Pepper J., et al., 2007, *PASP*, **119**, 923  
 Percy J. R., 1975, *AJ*, **80**, 698  
 Perry C. L., Johnston L., 1982, *ApJS*, **50**, 451  
 Pickles A., Depagne É., 2010, *PASP*, **122**, 1437  
 Raskin G., et al., 2011, *A&A*, **526**, A69  
 Renson P., 1988, *A&AS*, **76**, 127  
 Renson P., Manfroid J., 2009, *A&A*, **498**, 961  
 Ricker G. R., et al., 2015, *Journal of Astronomical Telescopes, Instruments, and Systems*, **1**, 014003  
 Romanyuk I. I., Kudryavtsev D. O., 2008, *Astrophysical Bulletin*, **63**, 139  
 Romanyuk I. I., Moiseeva A. V., Semenko E. A., Kudryavtsev D. O., Yakunin I. A., 2020, *Astrophysical Bulletin*, **75**, 294  
 Rózański T., Niemczura E., Lemiesz J., Posiłek N., Rózański P., 2022, *A&A*, **659**, A199  
 Saio H., Gautschi A., 2004, *MNRAS*, **350**, 485  
 Sbordone L., Bonifacio P., Castelli F., Kurucz R. L., 2004, *Memorie della Societa Astronomica Italiana Supplementi*, **5**, 93  
 Scardia M., Prieur J. L., Panseccchi L., Argyle R. W., Sala M., Ghigo M., Koechlin L., Aristidi E., 2006, *MNRAS*, **367**, 1170  
 Schneider H., 1986, *A&A*, **161**, 203  
 Schofield M., et al., 2019, *ApJS*, **241**, 12  
 Shang L.-H., et al., 2022, *ApJS*, **259**, 63  
 Shi F., Kurtz D., Saio H., Fu J., Zhang H., 2020, *ApJ*, **901**, 15  
 Shi F., Kurtz D. W., Holdsworth D. L., Saio H., Cunha M. S., Zhang H., Fu J., Handler G., 2021, *MNRAS*, **506**, 5629  
 Shibahashi H., 1979, *PASJ*, **31**, 87  
 Sikora J., Wade G. A., Power J., Neiner C., 2019, *MNRAS*, **483**, 2300  
 Skarka M., et al., 2022, *A&A*, **666**, A142  
 Slettebak A., Nassau J. J., 1959, *ApJ*, **129**, 88  
 Smalley B., et al., 2015, *MNRAS*, **452**, 3334  
 Stassun K. G., et al., 2018, *AJ*, **156**, 102  
 Stassun K. G., et al., 2019, *AJ*, **158**, 138  
 Tassoul M., 1980, *ApJS*, **43**, 469  
 Tonry J. L., et al., 2018, *ApJ*, **867**, 105  
 Wright C. O., Egan M. P., Kraemer K. E., Price S. D., 2003, *AJ*, **125**, 359  
 Zhao G., Zhao Y.-H., Chu Y.-Q., Jing Y.-P., Deng L.-C., 2012, *Research in Astronomy and Astrophysics*, **12**, 723  
 Zheng Z., Cao Z., Deng H., Mei Y., Tan L., Wang F., 2023, *ApJS*, **266**, 18  
 de Bruijne J. H. J., Eilers A. C., 2012, *A&A*, **546**, A61


## APPENDIX A: AUTHOR AFFILIATIONS

## APPENDIX B: SPECTRAL PLOTS


Here we provide the spectral plots used to derive new or updated spectral classifications. Information of the sources can be found in Sec. 3 We also provide, in Table B1, a log of all spectroscopic observations related to this work.

**Table B1.** Log of the spectroscopic observations collected for stars presented in this paper. The resolution refers to the canonical resolving power of the instrument in the configuration used. For full details see Section 3.


TIC	Spectrograph	Resolution	Nobs
21024812	HERMES	85 000	2
21024812	LAMOST	1 800	1
26833276	LAMOST	7 500	1
101624823	LAMOST	1 800	1
118247716	LAMOST	1 800	1
259017938	LAMOST	7 500	9
435263600	HERMES	85 000	2
445493624	HERMES	85 000	2
467074220	LAMOST	1 800	1




**Figure B1.** HERMES spectra of TIC 445493624 and TIC 21024812 in the blue region.




**Figure B2.** HERMES spectra of TIC 445493624 and TIC 21024812 in the red region.




**Figure B3.** LAMOST low resolution spectrum of TIC 21024812.




**Figure B4.** Segment of the LAMOST medium resolution spectrum of TIC 259017938.





**Figure B5.** Segment of the LAMOST medium resolution spectrum of TIC 26833276.




**Figure B6.** LAMOST low resolution spectrum of TIC 101624823 showing the clear presence of Cr II, with indications of Sr II and Eu II indicated.



**Figure B7.** LAMOST low resolution spectrum of TIC 118247716 showing the clear presence of Sr II, Eu II and Cr II.



**Figure B8.** HERMES spectra of TIC 435263600 showing the binary nature of the system. The two epochs are plotted in black and red, offset in intensity for clarity. The top panel shows the region classically used for spectral classification (note the significant change in the Ca II K line profile), while the bottom panel shows the region around  $H_{\alpha}$ . It is easy to see how multiple different classifications can arise from observing the star at different epochs.




**Figure B9.** LAMOST low resolution spectrum of TIC 467074220 showing the clear presence of Sr II and Cr II with an indication of Eu II.


### APPENDIX C: AMPLITUDE SPECTRA OF ROAP STARS KNOWN PRIOR TO THE LAUNCH OF TESS

### APPENDIX D: AMPLITUDE SPECTRA OF ROAP STARS OBSERVED BEYOND CYCLE 2


This paper has been typeset from a  $\text{\TeX}/\text{\LaTeX}$  file prepared by the author.




**Figure C1.** Amplitude spectrum for TIC 26418690 in the regions of previously reported variability. The top panel shows the low frequency range where  $\delta$  Sct modes have been reported (indicated by the arrows), but are below the *TESS* detection limit. The bottom panel shows the *TESS* detection of the roAp pulsation in this star.




**Figure C2.** Amplitude spectrum for TIC 26749633 showing the tentative detection of the known pulsation mode in the *TESS* data.




**Figure C3.** Amplitude spectrum for TIC 27395746 showing the detection of the  $\nu - \nu_{\text{rot}}$  sidelobe (indicated by the dashed vertical line) of the unseen pulsation mode (with a frequency indicated by the arrow).




**Figure C4.** Amplitude spectrum for TIC 77128654. The vertical dotted lines identify the pulsation modes.




**Figure C5.** Amplitude spectrum for TIC 123231021. The vertical dotted lines identify the pulsation mode provided in the literature. The fact that the mode is split here demonstrates the significant frequency variability present in this star.




**Figure C6.** Amplitude spectrum for TIC 158216369. The arrows identify the locations of the pulsation modes listed in the literature. Their amplitudes are below the noise level in the *TESS* data.




**Figure C7.** Amplitude spectrum for TIC 158271090. The vertical dotted line identifies the pulsation mode, with the dashed lines identifying the rotationally split sidelobes. The arrow identifies the location of a second mode in this star which is below the noise level in the *TESS* data.




**Figure C10.** Amplitude spectrum for TIC 264509538. The vertical dotted line identifies the pulsation mode, with the dashed lines indicating the rotationally split sidelobes.




**Figure C8.** Amplitude spectrum for TIC 158275114. The vertical dotted lines identify the pulsation modes detected in the *TESS* data, while the arrows indicate previously known pulsation modes.




**Figure C11.** Amplitude spectrum for TIC 272598185. The arrow indicates the previously known pulsation mode, which is not detected in the *TESS* data.




**Figure C9.** Amplitude spectrum for TIC 169078762. The vertical dotted line identifies the pulsation mode detected in the *TESS* data.




**Figure C12.** Amplitude spectrum for TIC 273777265. The vertical dotted line identifies the pulsation mode detected in the *TESS* data.




**Figure C13.** Amplitude spectrum for TIC 286992225. The vertical dotted lines identify the pulsation modes detected in the *TESS* data; the lowest frequency mode is a new detection.




**Figure C14.** Amplitude spectrum for TIC 302602874. The vertical dotted line identifies the pulsation mode, while the dashed lines identify the rotationally split sidelobes.




**Figure D1.** Amplitude spectrum for TIC 33604636. The vertical dotted line identifies the pulsation mode. After the removal of this peak, there is residual power remaining in this frequency range.




**Figure D2.** Amplitude spectrum for TIC 36576010. The vertical dotted line identifies the pulsation mode.




**Figure D3.** Amplitude spectrum for TIC 198781841. The vertical dotted lines identify the pulsation modes.




**Figure D4.** Amplitude spectrum for TIC 229960986. The vertical dotted line identifies the pulsation mode, with the dashed lines indicating the rotational sidelobes.




**Figure D5.** Amplitude spectrum for TIC 453826702. The vertical dotted lines identify the pulsation modes.




**Figure D8.** Amplitude spectrum for TIC 96855460. The vertical dotted line identifies the pulsation mode. The distorted shape is likely due to close, unresolved, sidelobes.




**Figure D6.** Amplitude spectrum for TIC 456673854. The vertical dotted line identifies the pulsation mode. We conclude this star to have, in increasing frequency, a dipole mode, a quadrupole mode, and a dipole mode. We have inferred the presence of the dipole modes the sidelobes.




**Figure D9.** Amplitude spectrum for TIC 100196783. The vertical dotted line identifies the pulsation mode.




**Figure D7.** Amplitude spectrum for TIC 630844439. The vertical dotted line identifies the pulsation mode, with the dashed lines showing the rotational sidelobes.




**Figure D10.** Amplitude spectrum for TIC 163587609. The vertical dotted lines identify the pulsation modes.




**Figure D11.** Amplitude spectrum for TIC 293265536. The vertical dotted line identifies the pulsation mode.




**Figure D14.** Amplitude spectrum for TIC 354619745. The vertical dotted lines identify the pulsation modes.




**Figure D12.** Amplitude spectrum for TIC 299000970. The vertical dotted lines identify the pulsation modes.




**Figure D15.** Amplitude spectrum for TIC 383521659 around the frequency range where pulsations were previously reported in spectroscopic data (Kurtz et al. 2007). There is no indication of variability in the *TESS* data.




**Figure D13.** Amplitude spectrum for TIC 318007796. The vertical dotted lines identify the pulsation modes, with the dashed lines showing the rotational sidelobes.



**Figure D16.** Amplitude spectrum for TIC 407929868. The vertical dotted line identifies the pulsation mode, while the dashed lines show the rotational sidelobes. In this view, the window function of the mode is hiding some of the lower-amplitude sidelobes.



**Figure D17.** Amplitude spectrum for TIC 420687462. The vertical dotted lines identify the pulsation modes.



**Figure D18.** Amplitude spectrum for TIC 445543326. The vertical dotted lines identify the pulsation modes, while the dashed lines show the rotational sidelobes.